

UChicago REU Notes Week 3

Zih-Yu Hsieh

July 3, 2026

Contents

1 Operad: Alternative Definition and Relation to Monad (ND, look at algebras)	1
2 Review: 2-Categories	5
3 Permutative Operads, $\mathcal{F}$-spaces	6

1 Operad: Alternative Definition and Relation to Monad (ND, look at algebras)

For the last section, we've seen an associative non- Σ operad, to produce the version in operad (equipped with symmetric group actions), we need to consider another version. But for this, I think there are some technicalities in the category theory, so we'll assume $\mathcal{V}$ is monoidally cocomplete.

An alternative assumption can be made that internal hom functor exists. Out of curiosity, are the two equivalent definition? Also, are there any symmetric monoidal category where it doesn't commute with coproduct?

Example 1.1. Let $\mathcal{M}(j) := \kappa[\Sigma_j] := \coprod_{\psi \in \Sigma_j} \kappa_{\psi}$, coproduct of $\#\Sigma_j$ copies of κ (each κ_{ψ} is the unit κ , subscript is for labeling convenience). It has a natural right Σ_j -action, where for each $\sigma \in \Sigma_j$, the identity $\kappa_{\psi\sigma^{-1}} \xrightarrow{\sim} \kappa_{\psi}$ induces an isomorphism on coproduct $\tilde{\sigma} \in \text{Aut}(\mathcal{M}(j))$:

$$\begin{array}{ccc}
 \kappa_{\psi} & \xrightarrow{\sim} & \kappa_{\psi\sigma} \\
 \downarrow \iota_{\psi} & & \downarrow \iota_{\psi\sigma} \\
 \coprod_{\psi \in \Sigma_j} \kappa_{\psi} & \xrightarrow[\exists! \tilde{\sigma}]{} & \coprod_{\psi \in \Sigma_j} \kappa_{\psi\sigma}
 \end{array}$$

This construction enforces $(\sigma\tilde{\sigma}') = \tilde{\sigma}' \circ \tilde{\sigma}$, which is a group homomorphism $\Sigma_j^{\text{op}} \rightarrow \text{Aut}_{\mathcal{V}}(\mathcal{M}(j))$.

There is a natural unit morphism $\eta : \kappa \rightarrow \kappa[\Sigma_1] = \kappa$ via identity.

Also, each identity morphism $\kappa_{\psi} \rightarrow \kappa$ induces a trivialization $t : \kappa[\Sigma_j] \rightarrow \kappa$. Combining the action $\psi \in \Sigma_k$ (by identifying concatenation of blocks $(j_{\psi^{-1}(1)}, \dots, j_{\psi^{-1}(k)}) = \{1, \dots, j\}$ in order, also inducing inclusion $\Sigma_{j_s} \hookrightarrow \Sigma_j$), we get the following morphism:

$$\kappa_{\psi} \otimes \left(\bigotimes_{s=1}^k \kappa[\Sigma_{j_s}] \right) \xrightarrow{\sim} \bigotimes_{s=1}^k \kappa[\Sigma_{j_{\psi^{-1}(s)}}] \xrightarrow{t^{k-1}} \kappa[\Sigma_{j_{\psi^{-1}(s)}}] \rightarrow \kappa[\Sigma_j]$$

Running through all $\psi \in \Sigma_k$, this induces a morphism $\gamma : \coprod_{\psi \in \Sigma_k} \left[\kappa_\psi \otimes \left(\bigotimes_{s=1}^k \kappa[\Sigma_{j_s}] \right) \right] \rightarrow \kappa[\Sigma_j]$, and use the fact $\otimes$ commutes with colimits, this is equivalent to a morphism $\gamma : \kappa[\Sigma_k] \otimes \left(\bigotimes_{s=1}^k \kappa[\Sigma_{j_s}] \right) \rightarrow \kappa[\Sigma_j]$.

This is helpful when defining an alternative definition.

Use Σ to denotes the category of finite sets, each $n \in \mathbb{N}$ has $\mathbf{n} := \{1, \dots, n\}$ (and $\mathbf{0}$ denotes the emptyset), with homsets being all possible isomorphisms (so $n \neq m$ there's no morphism between them, while the endomorphism is consists of permutations). Each object $\mathbf{n}$ has a canonical Σ_n -action. This is called a *permutation category*.

Definition 1.2. We define a Σ -object in the monoidal category $\mathcal{V}$ to be a functor $F : \Sigma^{\text{op}} \rightarrow \mathcal{V}$, which each $F(n)$ is equipped with a natural right Σ_n -action.

A morphism between Σ -objects will be natural transformations, which by definition each morphism at $\mathbf{n}$ is a Σ_n -equivariant morphism, as all $\sigma \in \Sigma_n$ (as a right action, equivalently an isomorphism) satisfies the following condition:

$$\begin{array}{ccc} F(\mathbf{n}) & \xrightarrow{\sigma} & F(\mathbf{n}) \\ \mu_n \downarrow & & \downarrow \mu_n \\ G(\mathbf{n}) & \xrightarrow{\sigma} & G(\mathbf{n}) \end{array}$$

this forms a *category of Σ -objects* for $\mathcal{V}$.

Now, consider the following operation:

Definition 1.3. Given F, G two Σ -objects of $\mathcal{V}$, define the *substitution product* $G \circ F : \Sigma^{\text{op}} \rightarrow \mathcal{V}$ as follow:

$$G \circ F(\mathbf{j}) := \coprod_{j_1 + \dots + j_k = j} G(\mathbf{k}) \otimes_{\kappa[\Sigma_k]} \left[\left(\bigotimes_{r=1}^k F(\mathbf{j}_r) \right) \otimes_{\kappa[\Sigma_{j_1} \times \dots \times j_k]} \kappa[\Sigma_j] \right]$$

Let's assume we know these are associative due to the coequalizer property :)

But roughly speaking, an operad is a Σ -object, that is also a monoid under the above product. (There are more details to work out...)

Talking about monads relevant to an operad $\mathcal{C}$ over $\mathcal{V}$, we define it as follow:

Definition 1.4. Define a functor $C : \mathcal{V} \rightarrow \mathcal{V}$ as follow:

$$\coprod_{j \geq 0} \mathcal{C}(j) \otimes \kappa[\Sigma_j] \otimes X^j \rightrightarrows \coprod_{j \geq 0} \mathcal{C}(j) \otimes X^j \xrightarrow{\text{coeq}} C(X)$$

Where given any $\psi \in \Sigma_j$, define the top morphism using $\mathcal{C}(j) \otimes \kappa_\psi \otimes X^j \xrightarrow{\sigma \otimes \text{id}} \mathcal{C}(j) \otimes X^j$ (action on $\mathcal{C}(j)$), which induces left action morphism $\mathcal{C}(j) \otimes \kappa[\Sigma_j] \otimes X^j \xrightarrow{\ell} \mathcal{C}(j) \otimes X^j$, then using coproduct universality to induce the morphism.

The bottom morphism is defined using $\mathcal{C}(j) \otimes X^j \xrightarrow{\text{id} \otimes \sigma^{-1}} \mathcal{C}(j) \otimes X^j$ (by sending the i th index X to the $\sigma^{-1}(i)$ th index X), then induce the coproduct morphism using similar logic (and label it as $\mathcal{C}(j) \otimes \kappa[\Sigma_j] \otimes X^j \xrightarrow{r} \mathcal{C}(j) \otimes X^j$). This is formally saying: A right action on $\mathcal{C}(j)$ transfers to an inverse left action on X^j .

Given a morphism $\psi : X \rightarrow Y$, the morphism $\text{id} \otimes \text{id} \otimes \psi^j : \mathcal{C}(j) \otimes \kappa[\Sigma_j] \otimes X^j \rightarrow \mathcal{C}(j) \otimes \kappa[\Sigma_j] \otimes Y^j$, and $\text{id} \otimes \psi^j : \mathcal{C}(j) \otimes X^j \rightarrow \mathcal{C}(j) \otimes Y^j$ but satisfies the following:

$$\begin{array}{ccc} \mathcal{C}(j) \otimes \kappa[\Sigma_j] \otimes X^j & \xrightarrow{\ell, r} & \mathcal{C}(j) \otimes X^j \\ \text{id} \otimes \text{id} \otimes \psi^j \downarrow & & \downarrow \text{id} \otimes \psi^j \\ \mathcal{C}(j) \otimes \kappa[\Sigma_j] \otimes Y^j & \xrightarrow{\ell, r} & \mathcal{C}(j) \otimes Y^j \end{array}$$

As including $\mathcal{C}(j) \otimes \kappa_\sigma \otimes X^j$, the top right route provides the morphism $\mathcal{C}(j) \otimes \kappa_\sigma \otimes X^j \xrightarrow{\sigma \otimes \text{id} \otimes \text{id}} \mathcal{C}(j) \otimes X^j \xrightarrow{\text{id} \otimes \psi^j} \mathcal{C}(j) \otimes Y^j$, equivalent to $\mathcal{C}(j) \otimes X^j \xrightarrow{\sigma \otimes \psi^j} \mathcal{C}(j) \otimes Y^j$, and by chasing through the coproduct property, we realize the following diagram commutes:

$$\begin{array}{ccc} \mathcal{C}(j) \otimes \kappa_\sigma \otimes X^j & \longrightarrow & \mathcal{C}(j) \otimes \kappa[\Sigma_j] \otimes X^j \\ \text{id} \otimes \text{id} \otimes \psi^j \downarrow & & \downarrow \text{id} \otimes \text{id} \otimes \psi^j \\ \mathcal{C}(j) \otimes \kappa_\sigma \otimes Y^j & \longrightarrow & \mathcal{C}(j) \otimes \kappa[\Sigma_j] \otimes Y^j \end{array}$$

Combining the two squares, do some diagram chasing, we realize the original square is commuting (with a pinch of coproduct property on $\kappa[\Sigma_j]$).

Finally, these induce the morphism between coproducts, and they of course commute with the group action by construction (as permuting ψ^j is equivalent to pre-permuting with ψ^j , or compose ψ^j with post-permuting). So, composing with the coequalizing morphism of Y , the desired diagram factors through coequalizing morphism of X , inducing a morphism $C(X) \rightarrow C(Y)$. Such induced morphism is functorial, hence C is indeed a functor.

To show C is a monad, we need to construct two natural transformations:

First, define $\eta' : \text{Id}_V \rightarrow C$ by taking the natural unit morphism $\eta : \kappa \rightarrow \mathcal{C}(1)$, and consider the sequence $\eta'_X : X \cong \kappa \otimes X \rightarrow \mathcal{C}(1) \otimes X \rightarrow \coprod_{j \geq 0} \mathcal{C}(j) \otimes X^j \rightarrow C(X)$. This “transformation” is natural, because the morphism after applying C is induced in the same manner.

To define the multiplication transformation, there are some slight work to do: Consider applying the construction to the first one, we get the following (quite chaotic) isomorphism:

$$\coprod_{k \geq 0} \mathcal{C}(k) \otimes \left(\coprod_{j \geq 0} \mathcal{C}(j) \otimes X^j \right)^k \cong \coprod_{k \geq 0} \coprod_{j_1 + \dots + j_k = j} \mathcal{C}(k) \otimes \mathcal{C}(j_1) \otimes \dots \otimes \mathcal{C}(j_k) \otimes X^j$$

This is because the k -fold monoidal product of the gigantic coproduct records all possible k -combinations of $\mathcal{C}(j_i)$, and the product on the copies of X precisely is $X^{j_1 + \dots + j_k}$. Apply the operadic structure morphism γ on each entry of the coproduct, we get the following morphism:

$$\overline{\mu}'_X : \coprod_{k \geq 0} \mathcal{C}(k) \otimes \left(\coprod_{j \geq 0} \mathcal{C}(j) \otimes X^j \right)^k \cong \coprod_{k \geq 0} \coprod_{j_1 + \dots + j_k = j} \mathcal{C}(k) \otimes \left(\bigotimes_{r=1}^k \mathcal{C}(j_r) \right) \otimes X^j \xrightarrow{\sqcup \gamma \otimes \text{id}} \coprod_{j \geq 0} \mathcal{C}(j) \otimes X^j \rightarrow C(X)$$

Which, there are some truly loooong diagram chase to check this factors through $C^2(X)$ (by checking it factors through $\mathcal{C}(k) \otimes (C(X))^2$ on each summand $\mathcal{C}(k) \otimes \left(\coprod_{j \geq 0} \mathcal{C}(j) \otimes X^j \right)^k$, then check again it satisfies the equivariance diagram; there are a lot of nuqnces about when to use the fact $\otimes$ commutes with colimits). This defines $\mu : C^2 \rightarrow C$.

By skipping some details, let's conclude that μ' is associative (intuitively, it's deduced from the associativity of tensor product; if the morphism before the factorization satisfies certain property, then passing through coequalizer it should still satisfy it). Also, by definition η' does serve as a unit (by the fact $\kappa \otimes C(X) \rightarrow \mathcal{C}(1) \otimes C(X) \rightarrow C^2(X) \rightarrow C(X)$ has passed through the collapsing map $\mathcal{C}(1) \otimes C(X) \rightarrow C(X)$, which cancels the unit, and reform identity).

Now, to say the correspondance between the operad and the monad, we compare their categories of algebras:

Theorem 1.5. *Given an operad $\mathcal{C}$ and its corresponding monad C , their categories of algebras are equivalent.*

Proof. To do. □

2 Review: 2-Categories

This section follows Emily Riehl's *Categorical Homotopy Theory*.

To properly define notions needed afterward, we need a concrete way of viewing categories enriched over small categories. This more generally follows categories enriched over symmetric monoidal categories, which are categories whose hom functor (or some similarly-behaved functor) are endowed with extra structures relevant to a monoidal category.

Notation 2.1. We'll use $(\mathcal{V}, \times, *)$ to denote a symmetric monoidal category here. If monoidally closed, we'll use $\mathcal{V}(-, -)$ to denote the internal hom functor.

Definition 2.2. A category $\mathcal{D}$ is *enriched over* $\mathcal{V}$, or called a $\mathcal{V}$ -category, if it consists of the following data:

- For each pair $x, y \in \mathcal{D}$, there is a *Hom-object* $\mathcal{D}(x, y) \in \mathcal{V}$.
- For each $x \in \mathcal{D}$, a *unit morphism* $\text{id}_x : * \rightarrow \mathcal{D}(x, x)$ in $\mathcal{V}$.
- For each triple $x, y, z \in \mathcal{D}$, a *composition morphism* $\circ : \mathcal{D}(y, z) \times \mathcal{D}(x, y) \rightarrow \mathcal{D}(x, z)$ in $\mathcal{V}$.

such that the following diagram commutes when given all $x, y, z, w \in \mathcal{D}$:

- *Associativity of composition:*

$$\begin{array}{ccc} \mathcal{D}(z, w) \times \mathcal{D}(y, z) \times \mathcal{D}(x, y) \times \circ & \xrightarrow{1} & \mathcal{D}(z, w) \times \mathcal{D}(x, z) \\ \circ \times 1 \downarrow & & \downarrow \circ \\ \mathcal{D}(y, w) \times \mathcal{D}(x, y) & \xrightarrow{\circ} & \mathcal{D}(x, w) \end{array}$$

- *Existence of Two-sided Units:*

$$\begin{array}{ccc} \mathcal{D}(x, y) \times * & \xrightarrow{1 \times \text{id}_x} & \mathcal{D}(x, y) \times \mathcal{D}(x, x) \\ \searrow & & \downarrow \circ \\ & & \mathcal{D}(x, y) \end{array} \quad \begin{array}{ccc} \mathcal{D}(y, y) \times \mathcal{D}(x, y) & \xleftarrow{\text{id}_y \times 1} & * \times \mathcal{D}(x, y) \\ \downarrow \circ & & \swarrow \\ \mathcal{D}(x, y) & & \end{array}$$

These conditions are mimicing the idea of homsets in a given category. For instance, with this definition by choosing $\mathcal{D}(-, -) := \text{Hom}_{\mathcal{D}}(-, -)$, we immediately realize all categories are enriched over **Set**. Since more general categories don't have concrete descriptions of objects using sets, we instead need a generalization so that the analogy of "hom functors" can land in more general categories.

3 Permutative Operads, $\mathcal{F}$ -spaces

Need to know:

- 2-categories
- work out details for $\mathcal{M}$ and the permutative operad $\mathcal{P} = \mathcal{E}\mathcal{M}$ (right now there is some basic idea about how it works).