

Math CS 122A HW4

Zih-Yu Hsieh

February 8, 2025

1

Question 1 Ahlfors Pg. 96 Problem 2:

Map the region between $|z| = 1$ and $|z - \frac{1}{2}| = \frac{1}{2}$ on a half plane.

Pf:

Consider the following transformation $g : \mathbb{C} \cup \{\infty\} \rightarrow \mathbb{C} \cup \{\infty\}$:

$$f(z) = \frac{z+1}{z-1} \cdot \frac{-i-1}{-i+1}, \quad g(z) = e^{\pi f(z)}$$

First, if consider the points $-i, -1, 1$ respectively on $|z| = 1$, linear transformation f maps the following:

$$f(-i) = \frac{-i+1}{-i-1} \cdot \frac{-i-1}{-i+1} = 1, \quad f(-1) = \frac{-1+1}{-1-1} \cdot \frac{-i-1}{-i+1} = 0, \quad f(1) = \infty$$

(Note: Since $f(1)$ is not defined under \mathbb{C} , it gets map to ∞).

Because the orientation of $|z| = 1$ is $-i$ to -1 to 1 , going clockwise, and the orientation of the image is 1 to 0 to ∞ , which on the right side is the half plane with positive imaginary parts. Hence, the right of $|z| = 1$ under this orientation (which is the interior of $|z| = 1$) gets mapped to the half plane $\text{Im}(z) > 0$.

Insert Image Here

Now, consider the points $\frac{1}{2}(1-i), 0, 1$ on $|z - \frac{1}{2}| = \frac{1}{2}$, linear transformation f maps the following:

$$\begin{aligned} f\left(\frac{1}{2}(1-i)\right) &= \frac{\left(\frac{1}{2} - \frac{1}{2}i\right) + 1}{\left(\frac{1}{2} - \frac{1}{2}i\right) - 1} \cdot \frac{-i-1}{-i+1} = \frac{(1-i) + 2}{(1-i) - 2} \cdot \frac{-i-1}{-i+1} = \frac{3-i}{-1-i} \cdot \frac{-1-i}{1-i} \\ &= \frac{3-i}{1-i} = \frac{(3-i)(1+i)}{(1-i)(1+i)} = \frac{3+1-i+3i}{2} = \frac{4+2i}{2} = 2+i \\ f(0) &= \frac{1}{-1} \cdot \frac{-i-1}{-i+1} = -\frac{-(1+i)^2}{(1-i)(1+i)} = \frac{2i}{2} = i, \quad f(1) = \infty \end{aligned}$$

So, since the three points gets mapped to $(2+i), i, \infty$ respectively, and linear transformation maps circle to circle, hence this is a circle passing through ∞ , or a straight line passing through i and $(2+i)$, which is the line $\text{Im}(z) = 1$. Then, with the orientation $\frac{1}{2}(1-i)$ to 0 to 1 , the image has orientation $(2+i)$ to i to ∞ , which the left side is the half plane $\text{Im}(z) < 1$. Hence, the left of $|z - \frac{1}{2}| = \frac{1}{2}$ under this orientation (the exterior of $|z - \frac{1}{2}| = \frac{1}{2}$) gets mapped to the half plane $\text{Im}(z) < 1$.

Insert Image Here

With the above statements, all points in the region between $|z| = 1$ and $|z - \frac{1}{2}| = \frac{1}{2}$ are in the interior of $|z| = 1$, and in the exterior of $|z - \frac{1}{2}| = \frac{1}{2}$. So, they are the intersection of $Im(z) > 0$ and $Im(z) < 1$.

Which, $\pi f(z)$ represents the region $0 < Im(z) < \pi$.

So, for all z_0 in the given open region, $z_0 = a + bi$, where $a \in \mathbb{R}$, and $0 < b < \pi$. So:

$$e^{z_0} = e^{a+bi} = e^a \cdot e^{ib}, \quad b \in (0, \pi)$$

Hence, e^{z_0} satisfies $\arg(e^{z_0}) = b \in (0, \pi)$, and $|e^{z_0}| = e^a > 0$, hence the image of the region $0 < Im(z) < \pi$ is in the half plane $Im(z) > 0$ (in fact, the image is the whole half plane, since the choice of $a \in \mathbb{R}$ and $b \in (0, \pi)$ are arbitrary, hence $e^a \in (0, \infty)$ could be any value in the given region).

Eventually, since $\pi f(z)$ maps the region between $|z| = 1$ and $|z - \frac{1}{2}| = \frac{1}{2}$ onto the region $0 < Im(z) < 1$, while e^z maps this new region onto the half plane $Im(z) > 0$, then the composition $e^{\pi f(z)}$ maps the desired region to the half plane $Im(z) > 0$.

Question 2 Ahlfors Pg. 97 Problem 5:

Map the inside of the right-hand branch of the hyperbola $x^2 - y^2 = a^2$ on the disk $|w| < 1$ so that the focus corresponds to $w = 0$ and the vertex to $w = -1$.

Pf:

WLOG, assume $a > 0$ (Note: $a < 0$ can be replaced with $(-a)$ instead). Under this configuration, the vertex is when $y = 0$, or $x = a$ for the right hand branch (the vertex is $z = a$). Also, the focus is given by $(ka, 0)$ with $k = \sqrt{1 + \frac{b'^2}{a'^2}}$ when given the hyperbola $\frac{x^2}{a'^2} - \frac{y^2}{b'^2} = 1$, which under this configuration, $a' = b' = a$, hence $k = \sqrt{2}$ (so the focus is $z = \sqrt{2}a$).

(Note 2: under the requirement, the focus and vertex needs to be two distinct points, hence $a \neq 0$).

Map of z^2 :

Notice that for all $z \in \mathbb{C}$, since $z = x + iy$ for some $x, y \in \mathbb{R}$, then $z^2 = (x^2 - y^2) + i \cdot 2xy$.

If take the plane $\operatorname{Re}(z) > 0$ (where $x > 0$), the map is injective: Suppose $z^2 = z_1^2$ for $z, z_1 \in \mathbb{C}$, then $z^2 - z_1^2 = (z - z_1)(z + z_1) = 0$, hence $z = z_1$ or $z = -z_1$. However, if restrict onto the plane $\operatorname{Re}(z) > 0$, then $z = -z_1$ is impossible for all values on this plane, hence $z = z_1$, showing it's injective.

Now, consider the inside of the right-hand branch of the hyperbola $x^2 - y^2 = a^2$, which is restricted by the condition $x^2 - y^2 \geq a^2$: For all $z = x + iy$ in the given region, $x^2 - y^2 \geq a^2$; hence, $z^2 = (x^2 - y^2) + i \cdot 2xy$ is in the half plane $\operatorname{Re}(w) \geq a^2$. Also, for all w in the half plane $\operatorname{Re}(w) \geq a^2$ ($a^2 > 0$), since it is in the domain of \sqrt{z} (which is in $\mathbb{C} \setminus \{x \in \mathbb{R} \mid x \leq 0\}$), then there exists $z = x + iy$ with $z^2 = w$, hence $\operatorname{Re}(w) = \operatorname{Re}(z^2) = (x^2 - y^2) \geq a^2$, showing that z is in the given region.

Hence, we can conclude that the function z^2 restricting onto the inside of the right-hand branch of the given hyperbola (with condition $x^2 - y^2 \geq a^2$), it is an injective function mapping the region onto the half plane $\operatorname{Re}(z) \geq a^2$.

Mapping the Half Plane $\operatorname{Re}(z) \geq a^2$ onto the Unit Disk:

Consider the following linear transformation:

$$f(w) = 1 - \frac{2a^2}{w}$$

For the points w_0 on the line $\operatorname{Re}(w) = a^2$, $w_0 = a^2 + iv$ for some $v \in \mathbb{R}$, hence the following is true:

$$f(w_0) = 1 - \frac{2a^2}{w_0} = \frac{w_0 - 2a^2}{w_0} = \frac{(a^2 + iv) - 2a^2}{a^2 + iv} = \frac{-a^2 + iv}{a^2 + iv} = \frac{-(a^2 - iv)}{a^2 + iv} = -\frac{\bar{w}_0}{w_0}$$

Hence, $|f(w_0)| = \left| -\frac{\bar{w}_0}{w_0} \right| = \frac{|\bar{w}_0|}{|w_0|} = 1$, the boundary of the half plane gets mapped to the boundary of the unit disk $|w| < 1$;

Also, for all points w_1 in the plane $\operatorname{Re}(w) > a^2$ (let $w = u + iv$ for $u, v \in \mathbb{R}$, hence $u > a^2$), there are two cases to consider. The following is what w_1 gets mapped to:

$$f(w_1) = 1 - \frac{2a^2}{w_1} = \frac{w_1 - 2a^2}{w_1} = \frac{(u - 2a^2) + iv}{u + iv}$$

First, if $u \leq 2a^2$, notice that since $0 \leq |u - 2a^2| = (2a^2 - u) < (2a^2 - a^2) = a^2 < u$, then, $|(u - 2a^2) + iv| = \sqrt{(u - 2a^2)^2 + v^2} < \sqrt{u^2 + v^2} = |w_1|$, hence $|f(w_1)| = \frac{|(u - 2a^2) + iv|}{|u + iv|} < 1$.

Else, if $u > 2a^2$, then since $0 < (u - 2a^2) < u$, then again $|(u - 2a^2) + iv| = \sqrt{(u - 2a^2)^2 + v^2} < \sqrt{u^2 + v^2} = |w_1|$, hence $|f(w_1)| = \frac{|(u-2a^2)+iv|}{|u+iv|} < 1$ is still true.

So, we can conclude that the half plane $Re(w) \geq a^2$ gets mapped to the unit disk $|w| = 1$, and since this is a linear transformation, the map is bijective.

Mapping Inside of Hyperbola to Unit Disk:

If Compose the two functions above, consider the following transformation $\bar{f}(z) = f(z^2) = 1 - \frac{2a^2}{z^2}$: First, for all z in the inside of the given branch of hyperbola (in the region $x^2 - y^2 \geq a^2$), z^2 appears in the half plane $Re(w) \geq a^2$, and there is a one-to-one correspondence between the two regions under the map; furthermore, since f maps the half plane $Re(w) \geq a^2$ to the unit disk $|w| \leq 1$, and is also a one-to-one correspondence, then the composition $f(z^2)$ maps the interior of the hyperbola to the unit disk.

Also, computing the following, we get:

$$\bar{f}(a) = 1 - \frac{2a^2}{a^2} = 1 - 2 = -1, \quad \bar{f}(\sqrt{2}a) = 1 - \frac{2a^2}{(\sqrt{2}a)^2} = 1 - \frac{2a^2}{2a^2} = 1 - 1 = 0$$

Which, since given the right branch of hyperbola $x^2 - y^2 = a^2$, $z_0 = a$ is the vertex and $z_1 = \sqrt{2}a$ is the focus, then the vertex gets mapped to -1 , and the focus gets mapped to 0 , hence this conformal map satisfies the given condition.

Insert Image Here

3

Question 3 Ahlfors Pg. 78 Problem 4:

Show that any linear transformation which transforms the real axis into itself can be written with real coefficient.

Pf:

Try to show that because of injectivity and surjectivity, 3 distinct real numbers must be mapped to $1, 0, \infty$ respectively.

Question 4 Ahlors Pg. 80 Problem 3:

If the consecutive vertices z_1, z_2, z_3, z_4 of a quadrilateral lie on a circle, prove that

$$|z_1 - z_3| \cdot |z_2 - z_4| = |z_1 - z_2| \cdot |z_3 - z_4| + |z_2 - z_3| \cdot |z_1 - z_4|$$

and interpret the result geometrically.

Pf:

Ptolemy's Theorem (try to look over that)

5

Question 5 Ahlfors Pg. 83 Problem 4:

Find the linear transformation which carries the circle $|z| = 2$ into $|z + 1| = 1$, the point -2 into the origin, and the origin into i .

Pf:

The symmetric point of the origin is some point, then try to unravel the mapping.

6

Question 6 Ahlfors Pg. 84 Problem 1:

If z_1, z_2, z_3, z_4 are points on a circle, show that z_1, z_3, z_4 and z_2, z_3, z_4 determine the same orientation if and only if $(z_1, z_2, z_3, z_4) > 0$.

Pf:

Not sure if it's rigorous enough, but can argue that z_1, z_2 stay on the same side iff they both get mapped to positive numbers.

7

Question 7 Ahlfors Pg. 88 Problem 6:

Find all circles which are orthogonal to $|z| = 1$ and $|z - 1| = 4$.

Pf:

Textbook Pg. 87, 88 were talking about this (about under conformal linear transformation, all the circles orthogonal to the two should correspond to a family of circles, all mapped to similar lines.)