

Math 111B HW2

Zih-Yu Hsieh

January 15, 2025

1

Question 1 Let $f : (a, b) \rightarrow \mathbb{R}$ be differentiable on (a, b) .

Prove : if $\forall x \in (a, b)$, $f'(x) \neq 0$, then f is one-to-one on (a, b) .

Give an example showing that the converse statement is in general not true.

Pf:

Suppose $\forall x \in (a, b)$, $f'(x) \neq 0$:

(1) $f'(x)$ is strictly less than or greater than 0 on (a, b) :

We'll prove by contradiction: Suppose $f'(x)$ is neither strictly less than 0 nor strictly greater than 0 on (a, b) , then there exists $x_0, x_1 \in (a, b)$, with $f'(x_0) \leq 0$ and $f'(x_1) \geq 0$, and by the assumption that $f'(x) \neq 0$, the strict inequality $f'(x_0) < 0$ and $f'(x_1) > 0$ is applied. (This also implies $x_0 \neq x_1$, since derivatives are different at the two points).

Recall that for function $f : [a, b] \rightarrow \mathbb{R}$ be differentiable on (a, b) , if $a < c < d < b$ and $f'(c) \neq f'(d)$, for any λ strictly in between $f'(c)$ and $f'(d)$ (either $f'(c) < \lambda < f'(d)$ or $f'(c) > \lambda > f'(d)$), there exists $x \in (c, d)$ with $f'(x) = \lambda$.

Then, first suppose $x_0 < x_1$: f is differentiable on (a, b) and $f'(x_0) < 0 < f'(x_1)$ implies there exists $x \in (x_0, x_1)$ with $f'(x) = 0$, which contradicts the assumption;

then suppose $x_1 < x_0$: again, f is differentiable on (a, b) and $f'(x_1) > 0 > f'(x_0)$ implies there exists $x \in (x_1, x_0)$ with $f'(x) = 0$, which again contradicts the assumption.

So, the assumption is false, $f'(x)$ must be strictly greater than 0 or less than 0 for all $x \in (a, b)$.

(2) f is strictly increasing or decreasing on (a, b) :

Based on **(1)**, $f'(x)$ is strictly less than 0 or strictly greater than 0.

Suppose $f'(x) > 0$ for all $x \in (a, b)$, then for any $x, y \in (a, b)$ with $x < y$, by the Mean Value Theorem, there exists $c \in (x, y) \subseteq (a, b)$, such that the following is true:

$$\frac{f(y) - f(x)}{y - x} = f'(c) > 0, \quad (f(y) - f(x)) = f'(c)(y - x)$$

Since $(y - x), f'(c) > 0$ by assumption, the $(f(y) - f(x)) = f'(c)(y - x) > 0$, thus $f(y) > f(x)$, showing that f is strictly increasing.

Similarly, suppose $f'(x) < 0$ for all $x \in (a, b)$, with the same x, y above, by Mean Value Theorem, there exists $c \in (x, y) \subseteq (a, b)$, such that the following is true:

$$\frac{f(y) - f(x)}{y - x} = f'(c) < 0, \quad (f(y) - f(x)) = f'(c)(y - x)$$

Since $(y - x) > 0$ and $f'(c) < 0$, then $(f(y) - f(x)) = f'(c)(y - x) < 0$, of $f(y) < f(x)$, showing that f is strictly decreasing.

With the above condition, since f is either strictly increasing or strictly decreasing on $[a, b]$, then for all $x, y \in (a, b)$, $x \neq y$ implies $f(x) \neq f(y)$ (or else it's no longer strictly increasing or decreasing). Thus, f is in fact one-to-one on (a, b) .

Counterexample of Converse:

Let $f : [-1, 1] \rightarrow \mathbb{R}$ be $f(x) = x^3$, which $f'(x) = 3x^2$, which $f'(0) = 0$. Yet, suppose $x, y \in (-1, 1)$ has $x^3 = y^3$, then:

$$x^3 - y^3 = (x - y)(x^2 + xy + y^2) = 0$$

Which, the only real solution is $x = y$ (since if treating y as constant, $y^2 - 4y^2 = -3y^2 \leq 0$; the only time with real solution is when $y = 0$, which implies $x^3 = 0$, or $x = 0$).

So, $f(x) = x^3$ is one-to-one on the region $(-1, 1)$, but still has $f'(0) = 0$, which is a counterexample.

2

Question 2

3

Question 3

4

Question 4

5

Question 5