



Published in final edited form as:

Trends Cancer. 2025 August ; 11(8): 753–769. doi:10.1016/j.trecan.2025.04.011.

Harnessing Technologies to Unravel Gastric Cancer Heterogeneity

Yuefan Wang^{1,*}, Ramon Jin², Joanne Xu³, Ding Chiao Lin¹, Zhenyu Sun¹, Yuanwei Xu¹, Qing K. Li¹, Hui Zhang¹

¹Department of Pathology, Johns Hopkins University School of Medicine, Baltimore, MD 21231, USA

²Division of Oncology and Gastroenterology, Washington University School of Medicine, Saint Louis, MO 63110, USA

³College of Engineering, The Ohio State University, Columbus, OH 43210, USA

Abstract

Gastric cancer arises from complex carcinogenic factor interactions, with limited treatment options due to the lack of targetable driver gene mutations and significant tumor heterogeneity. Recent studies have provided promising novel approaches to improve our understanding of gastric cancer heterogeneity through integrated characterization, combining genomics with emerging technologies. Delineating the molecular changes and targeting specific molecular subtypes will enhance the efficacy of gastric cancer treatment and improve clinical outcomes. This review provides a comprehensive overview of current technologies used in gastric cancer research, highlighting key discoveries and treatment strategies driven by these innovations. Finally, we discuss the emerging technology-guided directions and potential breakthroughs that could enhance the understanding of gastric cancer tumor heterogeneity, ultimately improving diagnosis and treatment.

Keywords

Gastric cancer; tumor heterogeneity; multi-omics technology; single cell analysis

Overview of gastric cancer

Gastric cancer (GC) is one of the most common types of cancers seen across the world, constituting an estimated 1 million new cases and 660,000 deaths in 2022 [1]. GC encompasses all cancers derived primarily from the stomach and can be described in various

*Correspondence: ywang298@jhu.edu (Y. Wang).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Declaration of Interests to the manuscript
The authors declare no competing interests.

ways when considering morphology of the cancer. The most common histomorphological type of GC is adenocarcinomas, however, it is a heterogeneous group of diseases, including many histomorphological subtypes. In addition, cancers from different stomach regions (cardia verse non-cardia), exhibit distinct histomorphologies, disease progression patterns, risk factors, and epidemiological features [2]. GC shows strong regional and ethnic disparities in incidental rates [1,3]. The incidental rates of GC remain high in East Asia, Latin America, and Eastern Europe, with men being twice as likely to be affected as women [1,3]. Specifically, individuals from these high-risk populations have a higher rate of non-cardia GC than non-Hispanic Caucasians in the same area, indicating both environmental and genetic contributions to GC formation [4]. The complexity of this disease makes the 5-year survival rate of GC low, especially for advanced stage and metastatic GC, which has a survival rate of less than 10% [5]. Lauren's histological classification focused on common subtypes of GC has become a widely adopted diagnostic tool for GC [2]. Subsequently, World Health Organization (WHO) Classification of GC has developed a more detailed classification scenario, based upon histomorphological and clinical characteristics of GC [6]. In the WHO classification, most common subtypes of GC have been included, such as previously recognized tubular/intestinal and poorly cohesive (such as signet ring cell) carcinoma, as well as papillary, mucinous, mixed and other subtypes. This knowledge has significantly advanced the understanding of GC biology. Given that GC is a complex and highly heterogeneous disease, leveraging diverse new technologies is essential to better study and understand its pathogenesis.

Recent advances in technologies have supported new findings on the mechanisms behind GC carcinogenesis, as well as introduced innovative strategies for preventing, diagnosing, and treating this disease [7,8]. Therefore, the pursuit and adaptation for ongoing technological innovations is crucial for advancing our understanding of GC heterogeneity and for developing new and improved treatment strategies. This article reviews innovative technologies that have significantly advanced our understanding of gastric cancer. In addition, future perspectives and new advanced technological developments to gain a deeper understanding of GC are evaluated.

New technological advancements to study gastric cancer heterogeneity

GC develops from a complex interplay of carcinogenic factors [2], including intrinsic factors like genomic and epigenomic changes, as well as the tumor microenvironment, and extrinsic factors such as bacterial and viral chronic infections, dysbiosis, and dietary habits. This complexity in carcinogenesis leads to high GC heterogeneity, presenting significant challenges for improving treatment. Given the lack of targetable driver mutations [9], employing diverse technologies and multi-omics analyses are crucial to identifying new therapeutic targets and clinical markers to guide precise and personalized treatment approaches (Figure 1).

Genomic technologies

Genomic analyses have become an invaluable tool in cancer research, revolutionizing our understanding of cancer biology by uncovering novel cancer markers and therapeutic targets.

Beginning with polymerase chain reaction (PCR) and further advanced by next-generation sequencing (NGS) technologies, the landscape of GC research and treatment is evolving [9]. NGS technologies are now widely utilized in genomics, epigenomics, and transcriptomics. In genomics, whole-genome sequencing (WGS) systematically reads entire DNA, revealing copy number variations (CNVs), single nucleotide variants (SNVs), and gene-fusion events [10,11]. Whole-exome sequencing (WES), which focuses on coding region information, efficiently identifies germline and somatic mutations [12]. RNA sequencing (RNA-seq), a transcriptomics technique, reads every base of RNA fragments or complementary DNA (cDNA) from large amounts of transcriptome data. It classifies several subtypes including exonic reads, junction reads, poly(A) end-reads, and non-coding RNA, assembles transcriptomes with or without reference genomes, and evaluates gene expression profiling [11,13].

Epigenetic changes are significant in cancer biology as they can directly affect gene expression to increase the plasticity of cancer cells [14]. Most epigenetic studies focus on DNA methylation and biochemical modifications on chromatin. DNA methylation can be detected by methylation arrays [15]. Chromatin immunoprecipitation (ChIP)-sequencing and the more recent Cleavage Under Targets and Release Using Nuclease (CUT&RUN) assays are used to profile DNA fragments bound by transcription factors or other chromatin-associated proteins [16]. These techniques use antibodies to specifically target and enrich for the DNA-protein interactions at selected genomic sites [17,18]. Genomic technologies have greatly advanced our understanding of GC. However, the functional implications of these genomic findings in GC need additional technologies to be elucidated.

Proteomic and proteogenomic approaches

Proteins perform biological functions based on their abundance, post-translational modifications (PTMs), and interactions within protein complexes, in both homeostatic and pathological states. Alterations in protein levels can indicate disease presence and help guide treatment strategies [19]. PTMs, such as phosphorylation, glycosylation, and ubiquitination, regulate protein function with aberrant modifications frequently implicated in cancer [20,21]. Mapping protein-protein interactions (PPIs) reveals disease-induced disruptions in protein complexes, helping in the identification of potential drug targets [22]. Mass spectrometry (MS) has been commonly utilized for proteomics, enabling the unbiased characterization of these protein-related patterns and has provided detailed insights into their functional roles [23] (Box 1).

To fully understand the functional implications of genomic and epigenomic changes in cancer, it is essential to integrate the analysis of proteins, PTMs, and PPIs with genomic, epigenomic, and transcriptomic data from the same tissue. This comprehensive approach has given rise to the field of proteogenomics[19,24], which provides deeper insights into biological and pathological processes at both the gene and protein levels. While proteogenomic study has been reported in GC, it mainly focused on the early-onset diffuse subtype GC [25]. Expanding proteogenomic research to include all ages and histological subtypes of GC is crucial for a more complete understanding of the disease. Strategies for proteogenomics and multiomics data integration involve leveraging computational tools and

statistical methods to align and harmonize diverse data types, such as genomic, epigenomic, transcriptomic, and proteomic datasets. Key approaches include addressing batch effects, overcoming missing data challenges, and developing robust frameworks for correlating variations across -omics layers, enabling deeper insights into tumor heterogeneity and potential therapeutic targets [24].

Microbiome and metabolites analysis

The Epstein–Barr virus (EBV) and *Helicobacter pylori* are contributors to human GC [26,27]. Pioneered by Woese [28], the 16S rRNA gene, a highly conserved component of ribosomal RNA in all bacteria, is widely used to detect the bacterial composition of various microbiomes. 16S rRNA gene contains both conserved and variable regions, enabling the differentiation between species and taxa [29]. Currently, 16S rRNA gene sequencing is the most common approach to studying gastric bacteria [30]. The gastric mycobiota is characterized by internal transcribed spacer (ITS) and 18S rRNA gene [31]. The viruses can be detected by PCR or real-time PCR (RT-PCR).

With the deep coverage provided by current genomic sequencing technologies, it is now possible to directly retrieve entire microbiome genome information from sequencing data using bioinformatics tools like PathSeq [32]. These tools for microbiome analysis have been used in pan-cancer mycobiome studies in GC [33], as well as in analyzing microbiome composition changes during GC progression [34]. This approach enables the integration of microbiome data within deep multi-omics analyses, providing a significant advantage by adding the microbiome layer to comprehensive multi-omics studies.

Not only do the microbiomes themselves act as factors in tumorigenesis, but their metabolites can contribute to tumor progression [35]. As the metabolites are small molecular weight organic molecules, human metabolites and microbial metabolites can be acquired simultaneously. The detection of metabolites can be achieved by mass spectrometry, sharing the same platform for cancer metabolomic studies [36,37].

Single-cell analyses and spatial omics techniques

Single-modality omics methods, which profile the genome, epigenome, transcriptome, proteome, metabolome, and other emerging omics, have increasingly been applied to single-cell cancer studies [38,39]. Compared to traditional bulk omics studies, where heterogeneity in cell types, cellular stages and dynamics were often omitted, single-cell omics have provided unprecedented insights into the cellular heterogeneity and molecular mechanisms driving cancer formation [40]. Studying intricate changes such as gene expression, genetic variations, epigenetic modifications, and protein expression patterns at the single-cell level is crucial for the understanding of the molecular underpinnings of GC [34,41-45] (Table 1). GC presents significant clinical challenges due to its phenotypic diversity and the complexity of its tumor microenvironment (TME), which contributes to immune evasion and resistance to therapies [46]. Single-cell analyses could provide insights into tumor heterogeneity, genetic alterations, TME, immune responses, prognostic markers, and potential therapeutic targets. When combined with spatial omics, these analyses provide

even deeper insights by revealing the spatial organization and interactions of cellular populations within the tumor.

Over the past 15 years, remarkable advancements (Table 1) in single-cell sequencing (sc-seq) and spatial transcriptomics (ST) have transformed our understanding of GC [44,45]. Early applications of sc-seq focused on transcriptome and single-cell genome analysis of GC cell lines, revealing marked genetic and transcriptional diversity [47]. This led to the identification of several notable mutated genes, providing crucial insights into the genetic alterations underlying GC and highlighting potential therapeutic targets. Furthermore, sc-seq technologies have been instrumental in studying tumor heterogeneity, uncovering substantial genetic and transcriptional diversity within tumor cells [41,42]. In addition, sc-seq can uncover immune and stromal cell states as well as ecotypes during GC progression, providing deeper insights into the entire evolutionary process [48]. Beyond using sc-seq alone, its integration with other technologies has yielded profound implications for tumor progression, treatment response, and the development of personalized therapies. Integrating sc-seq with technologies like CRISPR-based lineage tracing enhances understanding by tracking heritable barcodes across cell lineages under therapeutic pressures [49,50]. Emerging multi-omics approaches, including single-cell and spatially resolved multi-omics, remain in early development in GC [43,45,51]. These technologies integrate genomics, epigenomics, transcriptomics, proteomics, and metabolomics to provide a comprehensive view of cancer progression. Single-cell multi-omics uncovers intratumoral heterogeneity, rare cell populations, and dynamic states, while spatially resolved multi-omics maps the tumor microenvironment and spatial cell relationships. By connecting molecular changes to phenotypic outcomes across temporal and spatial dimensions, this integration fosters a holistic understanding of cancer biology, enabling novel therapeutic strategies and personalized treatments.

Gastric cancer biological insights gained from multi-omics studies

Before the genomic era, protein overexpression, such as HER2, had already been observed in GC [52]. Benefiting from PCR and sequencing (Figure 2), GC somatic mutations include common cancer mutations found across multiple cancer types (such as *TP53*, *KRAS*, *ERBB2*, and *PIK3CA*) as well as more tissue-specific or lineage-specific mutations (such as *CDH1*, *RHOA*, and *CLDN18-ARHGAP* fusions) [9]. The latest comprehensive genomic study identified 77 significantly mutated genes, which was conducted on a multi-ancestry cohort of over 1,300 GC cases [53]. Among these studies [53–55], only *TP53* was mutated in over 50% of the population, but it is considered as a non-targetable driver mutation. The remaining mutations were present in less than 25% of the total population. This highlights the lack of targetable driver mutations in GC, underscoring the limitations of single-target-based treatments for widespread use in this type of cancer.

Using NGS, comprehensive genomic and transcriptomic studies have provided a deeper and systematical understanding of GC. The Cancer Genome Atlas (TCGA) introduced a framework for the molecular characterization of GC, categorizing it into four distinct types [54]: EBV-positive, microsatellite instability (MSI), genomically stable (GS; enriched for Lauren diffuse type with specific somatic mutations), and tumors with

chromosomal instability (CIN). CIN subtypes frequently exhibit *TP53* mutations and tyrosine kinase receptor amplifications (such as *ERBB2* and *EGFR*), whereas mutations in phosphatidylinositol-4,5-bisphosphate 3-kinase catalytic subunit alpha (*PIK3CA*), AT-rich interaction domain-containing protein 1A (*ARID1A*), and hypermethylation are common in MSI and EBV subtypes [54,56]. Additionally, programmed cell death ligand 1/2 (PD-L1/2 or *CD274*) is often overexpressed through gene amplification and structural variation in EBV subgroups, making immune checkpoint inhibitors (ICIs) effective. ICIs are also effective in MSI subtypes due to high-frequency gene mutations and neoantigen expression [57]. In contrast, the GS subtype displays fewer genomic abnormalities leading to lower ICIs efficacy—only 12% in a study of 61 patients with metastatic GC, compared to 85.7% for MSI and 100% for EBV subtypes. Furthermore, the expression of HER2, a potential therapeutic target, is low in the GS subtype [57]. Thus, advanced therapeutic strategies are needed to improve outcomes for the GS subtype.

Diffuse-type GC, which aligns with the GS molecular subtype, features low-frequency structural chromosome aberrations and gene mutations. Key genetic variants include *CDH1* germline mutations and *RHOA* driver gene mutations [56]. While *CDH1* mutations are not viable treatment targets due to their “loss-of-function” nature [58], *RHOA* “gain-of-function” mutations present potential therapeutic opportunities, though challenges in inhibitor development remain [59,60]. Whole-genome and transcriptome analyses have identified *CLDN18-ARHGAP* fusion genes in diffuse-type GC, linked to poorer prognoses and highlighting the ineffectiveness of current chemotherapies [54,61,62]. An emerging approach involves activating T cells to target fusion genes in cancer cells [63].

In the TCGA study, the role of stromal cells may have been underestimated, a gap that was addressed in the subsequent the Asian Cancer Research Group (ACRG) study using transcriptomic data [55]. ACRG classified GC into two distinct types, MSI (best survival rate) and microsatellite stability (MSS). The MSS contains three subtypes: epithelial to mesenchymal transition (EMT, worst survival rate and prognosis), TP53⁺, and TP53⁻ [64]. The ACRG subtypes shared some common features with TCGA subtype [64], such as GS, EBV positive, and CIN subtypes in TCGA were enriched in MSS/EMT, MSS/TP53⁺, and MSS/TP53⁻ subtypes in ACRG [55]. Those subtypes showed distinct frequencies in different gastric anatomic locations [56], indicating GC is a highly heterogeneous disease.

To gain a deeper understanding of the heterogeneity, lineage diversity, and progression of GC, single-cell level analyses were performed. More than 30 distinct cell lineages have been identified in GC, and prognostic markers, such as signature gene panels, *SDC2* overexpression in cancer associated fibroblasts, and specific cell populations, have been discovered [41,42,48]. During GC progression, *OR51E1* was associated with endocrine cells in the early stages of malignant lesions [65], while *SOX9* and 25 other driver genes were identified in intestinal metaplasia [34]. *HES6* further aided in the identification of metaplasia at an early stage [65].

The initial proteomic and proteogenomic characterizations of GC focused on the diffuse type [25,66]. Proteome-based subtyping was found to be linked to diffuse type GC patient survival, with the immune response protein-enriched subtype being associated

with the poorest survival outcomes [66]. Integrated clustering in proteogenomic studies was used to identify functional subtypes-proliferation, immune response, metabolism, and invasionoffering deeper insights beyond those provided by genomic studies [25]. A follow-up study compared the molecular differences between diffuse and intestinal types at the proteomic level, revealing that *ARID1A* mutations had opposite prognostic effects in the two subtypes, with NF- κ B and SWI/SNF complexes identified as key distinguishing factors between them [67].

It is important to note that the current molecular subtyping of GC has had modest impact on the treatment of GC patients. Current molecular subtyping has been largely associated with retrospective clinical data to generate prognostic models [68] that have had minimal real world clinical applicability given limitations of availability, applicability, and timeliness. Clinical NGS testing has mostly been used to identify patients with actionable mutations with available matched therapies rather than molecular subtype classification purposes [69]. However, recognizing the deficient mismatch repair (MMR) and MSI subtype of GC has become clinically relevant due availability of biomarkers (immunohistochemistry for MMR protein expression or NGS testing for MSI status) and prospective data showing benefit of immune checkpoint inhibitors [70].

Multi-omics, especially proteogenomic characterization, is essential for fully understanding the molecular mechanisms in GC, as it connects both coding and non-coding mutations to protein functions and disease progression, offering deeper insights into tumor biology and potential treatments. An upcoming comprehensive CPTAC study will encompass all GC subtypes and stages [71]. These novel efforts may conceivably overcome the translational limitations of the current subtyping efforts by offering new biomarkers to personalize GC care, as well as novel mechanistic insights regarding targetable pathways to design future clinical trials. However, there is currently no evidence to demonstrate that multi-omics data are clinically translatable in terms of availability, applicability, and timeliness.

The microbiome plays a unique role in GC tumorigenesis has been increasingly recognized to be associated with dysbiosis in the stomach [72]. *H. pylori* is recognized as a key causal factor, with 90% of GC cases linked to *H. pylori* infection [73]. However, *H. pylori* is more enriched in normal adjacent tissues compared to the GC tissues [37,74], while the abundance of lactic acid bacteria, *Streptococcus*, and other genera including *Fusobacterium* increases in the GC tissues [30,37,75,76]. Multi-omic analysis of the microbiome and associated GC tissues have shown that *H. pylori* play a crucial role in the initial stages of GC formation, while other bacterial species may drive disease progression by generating carcinogenic N-nitroso compounds, reactive oxygen species, and metabolites that inhibit antitumor immune responses [30,77]. These studies have specifically identified cancer driving processes involve WNT/ β -catenin, MAPK/ERK, NF- κ B, and PI3K/Akt pathways [77,78]. Direct oncologic clinical translation of our basic understanding of the gut microbiome has been difficult as efforts to alter the gut microbiota have been focused on improving therapy and reducing treatment-related toxicity [79]. Testing for *H. pylori* is clinically available, timely, and has demonstrated clear clinical benefit in certain aspects of GC treatment. Specifically, *H. pylori* identification and successful eradication can prevent GC on a population-scale [80,81]. In addition, treatment of this gastric bacterium can prevent recurrence of early GC

[82] and improve survival in GC patients undergoing curative intent surgery [83]. But the implications of infection with *H. pylori* in patients with advanced GC is still controversial and an area of current active research [84,85].

Early detection of gastric cancer

Early detection and effective monitoring of tumor progression are crucial for reducing the burden and mortality of GC. Liquid biopsy is a non-invasive technique for detecting and analyzing circulating tumor cells (CTCs), circulating free DNA (cfDNA), circulating tumor DNA (ctDNA), non-coding RNAs (ncRNAs), and exosomes in biological fluid samples [86]. It holds promising clinical applications for the early diagnosis and screening of GC patients. CTCs, which are tumor cells that have detached from the primary tumor or metastases and entered the patient's bloodstream, represent an easily obtainable cancer specimen [87]. In a study of 116 patients that underwent gastrectomy for GC, the sensitivity and specificity of CTCs in GC detection compared with healthy controls were 85.3% and 90.3%, respectively [88]. A combination of cytokeratin 19 and 20 multiplex PCR assay from CTCs can detect GC with sensitivity and specificity of 87.5% and 94.7%, respectively [89]. These results suggest CTCs could be useful in the diagnosis of GC.

ctDNA can be detected in fluids from GC patients and normal controls. Changes in ctDNA across various pathological states have been successfully utilized as noninvasive biomarkers for liquid biopsies [90,91]. ctDNA levels were reported significantly higher in stage I GC patients (n = 124) when compared their levels with benign gastric disease and normal controls [92]. ctDNA is also valuable for monitoring disease recurrence after surgical procedures, which can be detected in individuals with early- and late-stage GC [93]. Changes in ctDNA had been reported with a sensitivity of 68.9% using a cutoff value of 2700 copies/mL ctDNA ($p < 0.001$) in a cohort of 429 GC patients and 95 healthy controls. In addition, ctDNA sequencing can identify fibroblast growth factor receptor 2 (*FGFR2*) amplification that may be missed by tissue testing in patients with advanced GC, and these patients may benefit from FGFR inhibition [94]. Another study demonstrated a significant association between pre-operatively detectable ctDNA and a higher T stage of the tumor ($p = 0.006$) [95].

Clinically, the benefit of liquid biopsy implementation has not been yet demonstrated for GC patients. CTCs can be predictive of treatment response for metastatic colon, breast, and prostate cancers [96-98] leading to availability of this technology clinically to guide treatment decisions for these advanced solid tumors. For ctDNA, the best prospective clinical data has shown its utility to guide adjuvant treatment decisions for colon cancer patients after surgery [99]. In terms of early cancer detection, the use of liquid biopsy is still being investigated with promising data showing sensitivity and specificity for colon cancer screening [100], as well as multicancer early detection potential [101,102]. Overall, ctDNA has the potential to improve sensitivity and specificity for early GC detection.

Non-coding RNAs (ncRNAs) are a type of RNA molecule with a key regulatory role in protein synthesis and potential biomarkers in GC [103,104]. ncRNAs influence gene expression and various cellular mechanisms, enhancing stability in exosomes and serving as

diagnostic or prognostic biomarkers [105]. Exosomes are lipid bilayer-enclosed extracellular vesicles (EVs) approximately 30-150 nm in size and contain a variety of nucleic acids, proteins, and lipids [106]. Increasing evidence has highlighted the potential diagnostic value of EVs or circulating exosomes for early detection of GC [107,108]. However, the isolation and characterization of exosomes remain technically challenging due to the lack of standardized protocols, affecting reproducibility and comparability of results [109]. Additionally, the stability and quantification of ncRNAs within EVs are highly sensitive to sample collection, processing, and storage conditions, introducing variability into clinical studies [110]. Addressing these limitations is essential to realize the potential of ncRNA and EV-based biomarkers for the early detection and management of GC in clinical practice.

Proteomics is another valuable tool for the early detection of GC [111] as both circulating and tissue proteins can serve as diagnostic markers for GC. Circulating proteins, including secreted proteins from tumor cells and effector/reactive proteins from the immune system, provide important diagnostic values. Given that many of these secreted and effector proteins undergo glycosylation, glycoproteomics analysis could be a powerful approach for early GC detection [112-114]. Altered glycosylation on serum haptoglobin [115], serum IgG [116], and salivary proteins [117], have been reported to associate with GC. Tumor tissue proteomic analysis can help to predict survival and sensitivity to chemotherapy [66]. The glycosylation changes on tissue proteins [112], such as mucins [113,118] and E-cadherin [119,120], can also help to detect GC as both may be early molecular events during gastric carcinogenesis. Specifically, a shift from sialomucins to sulfomucins has been associated with increased risk of GC progression in noncancerous metaplastic lesions in the stomach [121], and loss of E-cadherin expression has been found to be an early molecular event that can drive development of the diffuse histological subtype of GC [119,120,122,123].

In summary, although several single-omics methods have been applied to the early detection of GC, this area remains underexplored, and there are no FDA-approved biomarkers for GC early detection. Multi-omics-based biomarkers have also not been explored yet but represent a promising direction for future research.

Therapeutic Targets

Traditional chemotherapy for GC has relied on targeting rapidly dividing cells using cytotoxic agents. Drugs can disrupt DNA replication and cell division to induce tumor cell apoptosis. However, these therapies often lack specificity and can result in considerable off-target effects. To overcome the limitations of chemotherapy, there has been a paradigm shift towards targeted therapies that aim to selectively eradicate cancer cells while sparing normal tissues [124]. Clinical trials using multi-omics approaches are anticipated to drive significant advancements in the diagnosis and treatment of GC [124].

HER2 is a tyrosine kinase receptor that drives cell proliferation when overexpressed [125]. Anti-HER2 therapy is a pivotal advancement in GC treatment, leveraging targeted approaches to improve patient outcomes [64]. Anti-HER2 therapies, like trastuzumab, disrupt this pathway by binding to HER2 and inhibiting downstream signaling, while antibody-drug conjugates such as trastuzumab deruxtecan deliver chemotherapy directly

to HER2-positive cells [126]. Since HER2's discovery as an oncogene, omics technologies have deepened our understanding of HER2-positive GC, revealing mechanisms of resistance and guiding more precise treatment strategies [127,128]. The integration of advanced omics technologies has greatly enhanced our understanding of HER2-positive GC. Genomic tools like WGS and PCR identified *ERBB2* gene amplification as a key driver of the disease and helped pinpoint patients who may benefit from anti-HER2 therapies. Proteomics has revealed resistance mechanisms, such as elevated extracellular matrix proteins and PI3K-AKT pathway activation [129], suggesting that combining anti-HER2 agents with PI3K-AKT inhibitors could improve treatment outcomes. scRNA-seq has shown how GC tumors adapt at the single-cell level to resist therapy [130], while spatial metabolomics using MS has linked metabolic heterogeneity to better responses to trastuzumab [131]. These findings highlight the critical role of omics and single-cell technologies in refining HER2-targeted therapies and underscore the potential for developing new treatment strategies as these technologies advance.

Anti-PD-1/PD-L1 therapies have reshaped GC treatment by enabling the immune system to target cancer cells. Anti-PD-1/PD-L1 therapies, including pembrolizumab and nivolumab, block the interaction between tumor cells and T cells to restore the immune system's ability to attack tumors [132]. The KEYNOTE-059 trial demonstrated pembrolizumab's efficacy in advanced GC [133], leading to FDA approval for PD-L1-positive cases, while nivolumab was approved in combination with chemotherapy for first-line GC treatment [134]. The advancement of anti-PD-1/PD-L1 therapies in GC has been significantly driven by omics technologies. Genomics studies have identified *CD274* gene amplification in EBV subtype [57], suggesting that these patients may benefit from anti-PD-L1 therapy. Further genomic research on plasma ctDNA may uncover additional predictive factors for immune checkpoint therapy efficacy in GC [135]. Proteomics, using mass spectrometry and machine learning [136], has linked low activity in the complement and coagulation cascades pathway to positive anti-PD-1 responses [137]. scRNA-seq has identified NK cell markers as indicators for predicting therapy outcomes of GC [138]. These integrated omics approaches have enhanced our understanding of biomarkers and molecular mechanisms, helping to refine and improve the efficacy of anti-PD-1/PD-L1 therapies in GC [139]. The response to neoadjuvant chemotherapy has also been characterized using multi-omics approaches [140]. But to date, novel multi-omics technologies have not yielded translatable therapeutic targets or biomarkers clinically for gastric cancer in terms of availability, applicability, and timeliness.

In addition to these FDA-approved targeted therapies for GC, omics studies have expanded our understanding of other potential drug targets currently in clinical trials, such as Claudin 18.2 [62,141]. For instance, whole-transcriptome sequencing has characterized Claudin 18.2 as a significant target in advanced GC in cases where traditional immunohistochemistry assessments are challenging [142]. Moreover, proteomic studies have identified phosphorylation and cleavage activation as the primary post-translational modifications of Trop-2 in cancer [143]. Consequently, omics-based approaches are expected to continue exploring therapeutic targets in GC.

In summary, the development and application of targeted therapies are a significant advancement in the treatment of GC, harnessing the power of biologics to combat cancer cells. The integration of omics technologies, such as genomics, proteomics, and single-cell analysis, has been pivotal in identifying predictive biomarkers and understanding the molecular underpinnings of therapy response. Metabolomics also offers insights into GC heterogeneity and patient responses to treatment [144]. As research progresses, these technologies will continue to enhance our ability to predict, refine, and potentially improve the efficacy of targeted therapies in GC.

Concluding Remarks and Perspectives

GC is one of the leading causes of cancer-related deaths globally and presents unique complexities. Although current studies for GC, such as TCGA genomic profiling, early-onset proteogenomic characterizations, microbiota analyses, and early detection findings, have revealed important discoveries, comprehensive integrative studies that combine these technologies across a broader cohort of GC patients are still necessary. Such comprehensive proteogenomic studies enriched with multi-omics insights would provide deeper insights into tumor heterogeneity, molecular functions, and microbiome-host interaction, uncover more precise biomarkers, and identify potential new therapeutic targets, ultimately advancing personalized treatment approaches and improving survival rates in GC.

In addition to ongoing efforts on the characterization of GC at bulk level, single-cell characterization of GC provides valuable insights into its progression, heterogeneity, and potential early detection markers, as well as the TME. However, integrating multi-omics approaches at the single-cell level remains a significant challenge. Current multi-omics analyses may reach the ten- or hundred-cell resolution level, allowing for a more detailed profiling of specific cell types and lineages. Single-cell multi-omics along with spatial information will provide a more comprehensive landscape of the complex biology of GC, which could inform early detection and treatment strategies in the future. Cell state and ecotype analyses offer a powerful approach to bridge single-cell and bulk data, providing a new direction for advancing GC research.

Emerging biologics, including antibody-drug conjugates (ADCs), and chimeric antigen receptor (CAR)-T therapies, hold tremendous promise for advancing the treatment of GC. Multi-omics-based technologies can be applied into clinical studies, to pinpoint the target molecules, identify the potentially responsive patient population, and evaluate the treatment outcomes. As these innovative technologies continue to evolve, they hold the potential to significantly enhance early detection, enable individualized cancer characterization, and advance the development of precision oncology. But even with all these technological innovations (Box 2), there remain multiple challenges translating these new advances directly to improve clinical outcomes. The goal for these current and future multi-omics investigations of GC is to gain further insights into the intra- and interpatient heterogeneity of GC to develop novel targeted and personalized treatment strategies. However, there is currently limited availability of many of these new technologies in the clinic beyond targeted NGS [145]. In addition, many of these multi-omic studies have used GC samples gathered retrospectively from limited patient populations. It is unclear how these results apply to

all GC patients or whether these multi-omics strategies can be applied prospectively (i.e. applicability) [145]. Thirdly, besides generating multi-omics data [24], novel bioinformatic advances will be needed to interpret the results in an expeditious fashion [146]. In other words, these technologies will need to exhibit timeliness to benefit patients with GC. As such, there are currently few validated predictive and prognostic biomarkers for gastric cancer [64,147,148].

However, there are examples of technologically driven improvements in the treatment of patients with GC. The identification of deficient MMR and MSI subtypes of gastric cancer [54,55,64], has enabled biomarker (immunohistochemistry for MMR protein expression or NGS testing for MSI status) driven identification of a subset of GC patients that preferentially benefit from immune checkpoint inhibitors [149,150]. In addition, the development of zolbetuximab, a monoclonal antibody against the tight-junctional protein, Claudin-18.2, is a recent example of successful translational application of -omics data to predictive biomarker driven treatment [141]. Using an *in silico* bioinformatic approach, Claudin-18.2 was identified as a unique surface protein target in gastric cancer [151], and subsequent clinical trials including two large randomized phase III trials [152,153] led to the recent approval of this drug for a molecular subset of GC that expresses Claudin-18.2 [154]. Both above examples have been able to overcome the challenges of availability, applicability, and timeliness through efficient biomarker driven prospective clinical trial data demonstrating clinical improvements in GC treatment. The technological revolution and evolution driven by the merging of multiple technologies and disciplines, not only promises to improve patient outcomes but also shifts the focus from a “one-size-fits-all” approach to a more personalized, data-driven model. Ultimately, these advancements could pave the way for early detection strategies, more effective therapies, reduced treatment-related toxicities, and enhanced clinical outcomes, ushering in a new era of oncology that prioritizes the unique needs of each patient (see Outstanding questions).

Acknowledgement

This work was supported by National Institutes of Health, National Cancer Institute, the Clinical Proteomic Tumor Analysis Consortium (CPTAC, U24CA271079), the Early Detection Research Network (EDRN, U2CCA271895), and Pancreatic Cancer Detection Consortium (PCDC, U01CA274514). R.U.J. was supported by P30DK052574 and K08DK139376 by the National Institutes of Health. Figures were partly generated by BioRender.com

Reference

1. Bray F. et al. (2024) Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 74, 229–263. 10.3322/caac.21834 [PubMed: 38572751]
2. Smyth EC et al. (2020) Gastric cancer. *Lancet* 396, 635–648. 10.1016/s0140-6736(20)31288-5 [PubMed: 32861308]
3. Siegel RL et al. (2024) Cancer statistics, 2024. *CA Cancer J Clin* 74, 12–49. 10.3322/caac.21820 [PubMed: 38230766]
4. Shah SC et al. (2020) Population-Based Analysis of Differences in Gastric Cancer Incidence Among Races and Ethnicities in Individuals Age 50 Years and Older. *Gastroenterology* 159, 1705–1714.e1702. 10.1053/j.gastro.2020.07.049 [PubMed: 32771406]

5. Thrift AP et al. (2023) Global burden of gastric cancer: epidemiological trends, risk factors, screening and prevention. *Nat Rev Clin Oncol* 20, 338–349. 10.1038/s41571-023-00747-0 [PubMed: 36959359]
6. Nagtegaal ID et al. (2020) The 2019 WHO classification of tumours of the digestive system. *Histopathology* 76, 182–188. 10.1111/his.13975 [PubMed: 31433515]
7. Lei ZN et al. (2022) Signaling pathways and therapeutic interventions in gastric cancer. *Signal Transduct Target Ther* 7, 358. 10.1038/s41392-022-01190-w [PubMed: 36209270]
8. Hirata Y. et al. (2023) Early stage gastric adenocarcinoma: clinical and molecular landscapes. *Nat Rev Clin Oncol* 20, 453–469. 10.1038/s41571-023-00767-w [PubMed: 37264184]
9. Yeoh KG and Tan P (2022) Mapping the genomic diaspora of gastric cancer. *Nat Rev Cancer* 22, 71–84. 10.1038/s41568-021-00412-7 [PubMed: 34702982]
10. Goodwin S. et al. (2016) Coming of age: ten years of next-generation sequencing technologies. *Nat Rev Genet* 17, 333–351. 10.1038/nrg.2016.49 [PubMed: 27184599]
11. Slatko BE et al. (2018) Overview of Next-Generation Sequencing Technologies. *Curr Protoc Mol Biol* 122, e59. 10.1002/cpmb.59 [PubMed: 29851291]
12. Satam H. et al. (2023) Next-Generation Sequencing Technology: Current Trends and Advancements. *Biology (Basel)* 12. 10.3390/biology12070997
13. Kumar KR et al. (2024) Next-Generation Sequencing and Emerging Technologies. *Semin Thromb Hemost.* 10.1055/s-0044-1786397
14. Feinberg AP and Levchenko A (2023) Epigenetics as a mediator of plasticity in cancer. *Science* 379, eaaw3835. 10.1126/science.aaw3835 [PubMed: 36758093]
15. Hattori N. et al. (2023) DNA Methylation Analysis. In *Inflammation and Cancer: Methods and Protocols* (Jenkins BJ, ed), pp. 165–183, Springer US
16. Skene PJ and Henikoff S (2017) An efficient targeted nuclease strategy for high-resolution mapping of DNA binding sites. *Elife* 6. 10.7554/eLife.21856
17. Park PJ (2009) ChIP-seq: advantages and challenges of a maturing technology. *Nat Rev Genet* 10, 669–680. 10.1038/nrg2641 [PubMed: 19736561]
18. Nakato R and Sakata T (2021) Methods for ChIP-seq analysis: A practical workflow and advanced applications. *Methods* 187, 44–53. 10.1016/j.ymeth.2020.03.005 [PubMed: 32240773]
19. Rodriguez H. et al. (2021) The next horizon in precision oncology: Proteogenomics to inform cancer diagnosis and treatment. *Cell* 184, 1661–1670. 10.1016/j.cell.2021.02.055 [PubMed: 33798439]
20. Cao L. et al. (2021) Proteogenomic characterization of pancreatic ductal adenocarcinoma. *Cell* 184, 5031–5052.e5026. 10.1016/j.cell.2021.08.023 [PubMed: 34534465]
21. Geffen Y. et al. (2023) Pan-cancer analysis of post-translational modifications reveals shared patterns of protein regulation. *Cell* 186, 3945–3967.e3926. 10.1016/j.cell.2023.07.013 [PubMed: 37582358]
22. Lu H. et al. (2020) Recent advances in the development of protein-protein interactions modulators: mechanisms and clinical trials. *Signal Transduct Target Ther* 5, 213. 10.1038/s41392-020-00315-3 [PubMed: 32968059]
23. Aebersold R and Mann M (2003) Mass spectrometry-based proteomics. *Nature* 422, 198–207. 10.1038/nature01511 [PubMed: 12634793]
24. Mani DR et al. (2022) Cancer proteogenomics: current impact and future prospects. *Nat Rev Cancer* 22, 298–313. 10.1038/s41568-022-00446-5 [PubMed: 35236940]
25. Mun DG et al. (2019) Proteogenomic Characterization of Human Early-Onset Gastric Cancer. *Cancer Cell* 35, 111–124.e110. 10.1016/j.ccell.2018.12.003 [PubMed: 30645970]
26. Burke AP et al. (1990) Lymphoepithelial carcinoma of the stomach with Epstein-Barr virus demonstrated by polymerase chain reaction. *Mod Pathol* 3, 377–380 [PubMed: 2163534]
27. Malfertheiner P. et al. (2014) *Helicobacter pylori*: perspectives and time trends. *Nat Rev Gastroenterol Hepatol* 11, 628–638. 10.1038/nrgastro.2014.99 [PubMed: 25001975]
28. Woese CR (1987) Bacterial evolution. *Microbiological Reviews* 51, 221–271. doi:10.1128/mr.51.2.221-271.1987 [PubMed: 2439888]

29. Clarridge JE 3rd (2004) Impact of 16S rRNA gene sequence analysis for identification of bacteria on clinical microbiology and infectious diseases. *Clin Microbiol Rev* 17, 840–862, table of contents. 10.1128/cmr.17.4.840-862.2004 [PubMed: 15489351]
30. Stewart OA et al. (2020) The role of gastric microbiota in gastric cancer. *Gut Microbes* 11, 1220–1230. 10.1080/19490976.2020.1762520 [PubMed: 32449430]
31. Nash AK et al. (2017) The gut mycobiome of the Human Microbiome Project healthy cohort. *Microbiome* 5, 153. 10.1186/s40168-017-0373-4 [PubMed: 29178920]
32. Walker MA et al. (2018) GATK PathSeq: a customizable computational tool for the discovery and identification of microbial sequences in libraries from eukaryotic hosts. *Bioinformatics* 34, 4287–4289. 10.1093/bioinformatics/bty501 [PubMed: 29982281]
33. Dohlman AB et al. (2022) A pan-cancer mycobiome analysis reveals fungal involvement in gastrointestinal and lung tumors. *Cell* 185, 3807–3822.e3812. 10.1016/j.cell.2022.09.015 [PubMed: 36179671]
34. Huang KK et al. (2023) Spatiotemporal genomic profiling of intestinal metaplasia reveals clonal dynamics of gastric cancer progression. *Cancer Cell* 41, 2019–2037.e2018. 10.1016/j.ccell.2023.10.004 [PubMed: 37890493]
35. Louis P. et al. (2014) The gut microbiota, bacterial metabolites and colorectal cancer. *Nat Rev Microbiol* 12, 661–672. 10.1038/nrmicro3344 [PubMed: 25198138]
36. Alseekh S. et al. (2021) Mass spectrometry-based metabolomics: a guide for annotation, quantification and best reporting practices. *Nat Methods* 18, 747–756. 10.1038/s41592-021-01197-1 [PubMed: 34239102]
37. Dai D. et al. (2021) Interactions between gastric microbiota and metabolites in gastric cancer. *Cell Death Dis* 12, 1104. 10.1038/s41419-021-04396-y [PubMed: 34819503]
38. Cuomo ASE et al. (2023) Single-cell genomics meets human genetics. *Nat Rev Genet* 24, 535–549. 10.1038/s41576-023-00599-5 [PubMed: 37085594]
39. Baysoy A. et al. (2023) The technological landscape and applications of single-cell multi-omics. *Nat Rev Mol Cell Biol* 24, 695–713. 10.1038/s41580-023-00615-w [PubMed: 37280296]
40. (2023) A focus on single-cell omics. *Nat Rev Genet* 24, 485. 10.1038/s41576-023-00628-3 [PubMed: 37464023]
41. Wang R. et al. (2021) Single-cell dissection of intratumoral heterogeneity and lineage diversity in metastatic gastric adenocarcinoma. *Nat Med* 27, 141–151. 10.1038/s41591-020-1125-8 [PubMed: 33398161]
42. Kumar V. et al. (2022) Single-Cell Atlas of Lineage States, Tumor Microenvironment, and Subtype-Specific Expression Programs in Gastric Cancer. *Cancer Discov* 12, 670–691. 10.1158/2159-8290.Cd-21-0683 [PubMed: 34642171]
43. Sun C. et al. (2023) Spatially resolved multi-omics highlights cell-specific metabolic remodeling and interactions in gastric cancer. *Nat Commun* 14, 2692. 10.1038/s41467-023-38360-5 [PubMed: 37164975]
44. Ren L. et al. (2024) Applications of single-cell omics and spatial transcriptomics technologies in gastric cancer (Review). *Oncol Lett* 27, 152. 10.3892/ol.2024.14285 [PubMed: 38406595]
45. Ma H. et al. (2025) Spatially Resolved Tumor Ecosystems and Cell States in Gastric Adenocarcinoma Progression and Evolution. *Cancer Discov* 15, 767–792. 10.1158/2159-8290.Cd-24-0605 [PubMed: 39774838]
46. Yasuda T and Wang YA (2024) Gastric cancer immunosuppressive microenvironment heterogeneity: implications for therapy development. *Trends Cancer* 10, 627–642. 10.1016/j.trecan.2024.03.008 [PubMed: 38600020]
47. Andor N. et al. (2020) Joint single cell DNA-seq and RNA-seq of gastric cancer cell lines reveals rules of in vitro evolution. *NAR Genom Bioinform* 2, lqaa016. 10.1093/nargab/lqaa016 [PubMed: 32215369]
48. Wang R et al. (2023) Evolution of immune and stromal cell states and ecotypes during gastric adenocarcinoma progression. *Cancer Cell* 41, 1407–1426.e1409. 10.1016/j.ccell.2023.06.005 [PubMed: 37419119]

49. Simeonov KP et al. (2021) Single-cell lineage tracing of metastatic cancer reveals selection of hybrid EMT states. *Cancer Cell* 39, 1150–1162.e1159. 10.1016/j.ccell.2021.05.005 [PubMed: 34115987]
50. Yang D. et al. (2022) Lineage tracing reveals the phylodynamics, plasticity, and paths of tumor evolution. *Cell* 185, 1905–1923.e1925. 10.1016/j.cell.2022.04.015 [PubMed: 35523183]
51. Bian S. et al. (2023) Integrative single-cell multiomics analyses dissect molecular signatures of intratumoral heterogeneities and differentiation states of human gastric cancer. *Natl Sci Rev* 10, nwad094. 10.1093/nsr/nwad094 [PubMed: 37347037]
52. Yonemura Y. et al. (1992) Correlation of c-erbB-2 protein expression and lymph node status in early gastric cancer. *Oncology* 49, 363–367. 10.1159/000227074 [PubMed: 1356259]
53. Totoki Y. et al. (2023) Multiancestry genomic and transcriptomic analysis of gastric cancer. *Nat Genet* 55, 581–594. 10.1038/s41588-023-01333-x [PubMed: 36914835]
54. (2014) Comprehensive molecular characterization of gastric adenocarcinoma. *Nature* 513, 202–209. 10.1038/nature13480 [PubMed: 25079317]
55. Cristescu R. et al. (2015) Molecular analysis of gastric cancer identifies subtypes associated with distinct clinical outcomes. *Nat Med* 21, 449–456. 10.1038/nm.3850 [PubMed: 25894828]
56. Nagaraja AK et al. (2019) Genomics and Targeted Therapies in Gastroesophageal Adenocarcinoma. *Cancer Discov* 9, 1656–1672. 10.1158/2159-8290.Cd-19-0487 [PubMed: 31727671]
57. Kim ST et al. (2018) Comprehensive molecular characterization of clinical responses to PD-1 inhibition in metastatic gastric cancer. *Nat Med* 24, 1449–1458. 10.1038/s41591-018-0101-z [PubMed: 30013197]
58. Liu Y. et al. (2018) Comparative Molecular Analysis of Gastrointestinal Adenocarcinomas. *Cancer Cell* 33, 721–735.e728. 10.1016/j.ccell.2018.03.010 [PubMed: 29622466]
59. Kakiuchi M. et al. (2014) Recurrent gain-of-function mutations of RHOA in diffuse-type gastric carcinoma. *Nat Genet* 46, 583–587. 10.1038/ng.2984 [PubMed: 24816255]
60. Zhang H. et al. (2020) Gain-of-Function RHOA Mutations Promote Focal Adhesion Kinase Activation and Dependency in Diffuse Gastric Cancer. *Cancer Discov* 10, 288–305. 10.1158/2159-8290.Cd-19-0811 [PubMed: 31771969]
61. Shu Y. et al. (2018) Prognostic significance of frequent CLDN18-ARHGAP26/6 fusion in gastric signet-ring cell cancer. *Nat Commun* 9, 2447. 10.1038/s41467-018-04907-0 [PubMed: 29961079]
62. Nakayama I. et al. (2024) Claudin 18.2 as a novel therapeutic target. *Nat Rev Clin Oncol* 21, 354–369. 10.1038/s41571-024-00874-2 [PubMed: 38503878]
63. Wang Y. et al. (2024) Immunotherapies targeting the oncogenic fusion gene CLDN18-ARHGAP in gastric cancer. *EMBO Mol Med* 16, 2170–2187. 10.1038/s44321-024-00120-3 [PubMed: 39164472]
64. Zeng Y and Jin RU (2022) Molecular pathogenesis, targeted therapies, and future perspectives for gastric cancer. *Semin Cancer Biol* 86, 566–582. 10.1016/j.semcancer.2021.12.004 [PubMed: 34933124]
65. Zhang P. et al. (2019) Dissecting the Single-Cell Transcriptome Network Underlying Gastric Premalignant Lesions and Early Gastric Cancer. *Cell Rep* 27, 1934–1947.e1935. 10.1016/j.celrep.2019.04.052 [PubMed: 31067475]
66. Ge S. et al. (2018) A proteomic landscape of diffuse-type gastric cancer. *Nat Commun* 9, 1012. 10.1038/s41467-018-03121-2 [PubMed: 29520031]
67. Shi W. et al. (2023) Multilevel proteomic analyses reveal molecular diversity between diffuse-type and intestinal-type gastric cancer. *Nat Commun* 14, 835. 10.1038/s41467-023-35797-6 [PubMed: 36788224]
68. Sohn BH et al. (2017) Clinical Significance of Four Molecular Subtypes of Gastric Cancer Identified by The Cancer Genome Atlas Project. *Clin Cancer Res* 23, 4441–4449. 10.1158/1078-0432.CCR-16-2211 [PubMed: 28747339]
69. Gibbs SN et al. (2023) Comprehensive Review on the Clinical Impact of Next-Generation Sequencing Tests for the Management of Advanced Cancer. *JCO Precis Oncol* 7, e2200715. 10.1200/PO.22.00715 [PubMed: 37285561]

70. Ratti M. et al. (2018) Microsatellite instability in gastric cancer: molecular bases, clinical perspectives, and new treatment approaches. *Cell Mol Life Sci* 75, 4151–4162. 10.1007/s00018-018-2906-9 [PubMed: 30173350]
71. Wang Y and Consortium, C.P.T.A. (2024) Abstract 1851: Comprehensive multilevel proteomics characterization of gastric cancer. *Cancer Research* 84, 1851–1851. 10.1158/1538-7445.Am2024-1851
72. Zeng R. et al. (2024) Stomach microbiota in gastric cancer development and clinical implications. *Gut* 73, 2062–2073. 10.1136/gutjnl-2024-332815 [PubMed: 38886045]
73. Malfertheiner P. et al. (2023) *Helicobacter pylori* infection. *Nat Rev Dis Primers* 9, 19. 10.1038/s41572-023-00431-8 [PubMed: 37081005]
74. Liu X. et al. (2019) Alterations of gastric mucosal microbiota across different stomach microhabitats in a cohort of 276 patients with gastric cancer. *EBioMedicine* 40, 336–348. 10.1016/j.ebiom.2018.12.034 [PubMed: 30584008]
75. Vinasco K. et al. (2019) Microbial carcinogenesis: Lactic acid bacteria in gastric cancer. *Biochim Biophys Acta Rev Cancer* 1872, 188309. 10.1016/j.bbcan.2019.07.004 [PubMed: 31394110]
76. Coker OO et al. (2018) Mucosal microbiome dysbiosis in gastric carcinogenesis. *Gut* 67, 1024–1032. 10.1136/gutjnl-2017-314281 [PubMed: 28765474]
77. Chattopadhyay I. et al. (2023) Diversification and deleterious role of microbiome in gastric cancer. *Cancer Rep (Hoboken)* 6, e1878. 10.1002/cnr2.1878 [PubMed: 37530125]
78. Fu K. et al. (2024) *Streptococcus anginosus* promotes gastric inflammation, atrophy, and tumorigenesis in mice. *Cell* 187, 882–896.e817. 10.1016/j.cell.2024.01.004 [PubMed: 38295787]
79. Zhao LY et al. (2023) Role of the gut microbiota in anticancer therapy: from molecular mechanisms to clinical applications. *Signal Transduct Target Ther* 8, 201. 10.1038/s41392-023-01406-7 [PubMed: 37179402]
80. Kumar S. et al. (2020) Risk Factors and Incidence of Gastric Cancer After Detection of *Helicobacter pylori* Infection: A Large Cohort Study. *Gastroenterology* 158, 527–536 e527. 10.1053/j.gastro.2019.10.019 [PubMed: 31654635]
81. Li D. et al. (2023) Effect of *Helicobacter pylori* Eradication Therapy on the Incidence of Noncardia Gastric Adenocarcinoma in a Large Diverse Population in the United States. *Gastroenterology* 165, 391–401 e392. 10.1053/j.gastro.2023.04.026 [PubMed: 37142201]
82. Choi IJ et al. (2018) *Helicobacter pylori* Therapy for the Prevention of Metachronous Gastric Cancer. *N Engl J Med* 378, 1085–1095. 10.1056/NEJMoa1708423 [PubMed: 29562147]
83. Zhao Z. et al. (2024) Anti-*Helicobacter pylori* Treatment in Patients With Gastric Cancer After Radical Gastrectomy. *JAMA Netw Open* 7, e243812. 10.1001/jamanetworkopen.2024.3812 [PubMed: 38546641]
84. Fang X et al. (2017) Positive *Helicobacter pylori* status is associated with better overall survival for gastric cancer patients: evidence from case-cohort studies. *Oncotarget* 8, 79604–79617. 10.18632/oncotarget.18758 [PubMed: 29108340]
85. Magahis PT et al. (2023) Impact of *Helicobacter pylori* infection status on outcomes among patients with advanced gastric cancer treated with immune checkpoint inhibitors. *J Immunother Cancer* 11. 10.1136/jitc-2023-007699
86. Zhang Z et al. (2022) Liquid biopsy in gastric cancer: predictive and prognostic biomarkers. *Cell Death Dis* 13, 903. 10.1038/s41419-022-05350-2 [PubMed: 36302755]
87. Yap TA et al. (2014) Circulating tumor cells: a multifunctional biomarker. *Clin Cancer Res* 20, 2553–2568. 10.1158/1078-0432.Ccr-13-2664 [PubMed: 24831278]
88. Kang HM et al. (2017) Circulating tumor cells detected by lab-on-a-disc: Role in early diagnosis of gastric cancer. *PLoS One* 12, e0180251. 10.1371/journal.pone.0180251 [PubMed: 28662130]
89. Vaiopoulos AG et al. (2014) Detection of circulating tumor cells in colorectal and gastric cancer using a multiplex PCR assay. *Anticancer Res* 34, 3083–3092 [PubMed: 24922677]
90. Pessoa LS et al. (2020) ctDNA as a cancer biomarker: A broad overview. *Crit Rev Oncol Hematol* 155, 103109. 10.1016/j.critrevonc.2020.103109 [PubMed: 33049662]
91. Mencil J. et al. (2022) The Role of ctDNA in Gastric Cancer. *Cancers (Basel)* 14. 10.3390/cancers14205105

92. Qian C. et al. (2017) Alu-based cell-free DNA: a novel biomarker for screening of gastric cancer. *Oncotarget* 8, 54037–54045. 10.18632/oncotarget.11079 [PubMed: 28903321]
93. Varkalaite G. et al. (2021) Liquid Biopsy in Gastric Cancer: Analysis of Somatic Cancer Tissue Mutations in Plasma Cell-Free DNA for Predicting Disease State and Patient Survival. *Clin Transl Gastroenterol* 12, e00403. 10.14309/ctg.0000000000000403 [PubMed: 34644276]
94. Lan YT et al. (2017) Clinical relevance of cell-free DNA in gastrointestinal tract malignancy. *Oncotarget* 8, 3009–3017. 10.18632/oncotarget.13821 [PubMed: 27936467]
95. Yang J. et al. (2020) Deep sequencing of circulating tumor DNA detects molecular residual disease and predicts recurrence in gastric cancer. *Cell Death Dis* 11, 346. 10.1038/s41419-020-2531-z [PubMed: 32393783]
96. Hayes DF et al. (2006) Circulating tumor cells at each follow-up time point during therapy of metastatic breast cancer patients predict progression-free and overall survival. *Clin Cancer Res* 12, 4218–4224. 10.1158/1078-0432.CCR-05-2821 [PubMed: 16857794]
97. Cohen SJ et al. (2008) Relationship of circulating tumor cells to tumor response, progression-free survival, and overall survival in patients with metastatic colorectal cancer. *J Clin Oncol* 26, 3213–3221. 10.1200/JCO.2007.15.8923 [PubMed: 18591556]
98. de Bono JS et al. (2008) Circulating tumor cells predict survival benefit from treatment in metastatic castration-resistant prostate cancer. *Clin Cancer Res* 14, 6302–6309. 10.1158/1078-0432.CCR-08-0872 [PubMed: 18829513]
99. Tie J. et al. (2022) Circulating Tumor DNA Analysis Guiding Adjuvant Therapy in Stage II Colon Cancer. *N Engl J Med* 386, 2261–2272. 10.1056/NEJMoa2200075 [PubMed: 35657320]
100. Chung DC et al. (2024) A Cell-free DNA Blood-Based Test for Colorectal Cancer Screening. *N Engl J Med* 390, 973–983. 10.1056/NEJMoa2304714 [PubMed: 38477985]
101. Schrag D. et al. (2023) Blood-based tests for multicancer early detection (PATHFINDER): a prospective cohort study. *Lancet* 402, 1251–1260. 10.1016/S0140-6736(23)01700-2 [PubMed: 37805216]
102. Nicholson BD et al. (2023) Multi-cancer early detection test in symptomatic patients referred for cancer investigation in England and Wales (SYMPLOY): a large-scale, observational cohort study. *Lancet Oncol* 24, 733–743. 10.1016/S1470-2045(23)00277-2 [PubMed: 37352875]
103. Xie S. et al. (2020) Non-coding RNAs in gastric cancer. *Cancer Lett* 493, 55–70. 10.1016/j.canlet.2020.06.022 [PubMed: 32712234]
104. Tan H. et al. (2020) Long non-coding RNAs in gastric cancer: New emerging biological functions and therapeutic implications. *Theranostics* 10, 8880–8902. 10.7150/thno.47548 [PubMed: 32754285]
105. Chen B. et al. (2022) Targeting non-coding RNAs to overcome cancer therapy resistance. *Signal Transduct Target Ther* 7, 121. 10.1038/s41392-022-00975-3 [PubMed: 35418578]
106. Johnstone RM et al. (1987) Vesicle formation during reticulocyte maturation. Association of plasma membrane activities with released vesicles (exosomes). *J Biol Chem* 262, 9412–9420 [PubMed: 3597417]
107. Fu M. et al. (2019) Exosomes in gastric cancer: roles, mechanisms, and applications. *Mol Cancer* 18, 41. 10.1186/s12943-019-1001-7 [PubMed: 30876419]
108. Shojaeian A. et al. (2024) Harnessing exosomes in theranostic applications: advancements and insights in gastrointestinal cancer research. *Discov Oncol* 15, 162. 10.1007/s12672-024-01024-x [PubMed: 38743146]
109. Zhang TN et al. (2021) Non-Coding RNAs and Extracellular Vehicles: Their Role in the Pathogenesis of Gestational Diabetes Mellitus. *Front Endocrinol (Lausanne)* 12, 664287. 10.3389/fendo.2021.664287 [PubMed: 34093439]
110. Valadi H. et al. (2007) Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nat Cell Biol* 9, 654–659. 10.1038/ncb1596 [PubMed: 17486113]
111. Nakayasu ES et al. (2021) Tutorial: best practices and considerations for massspectrometry-based protein biomarker discovery and validation. *Nat Protoc* 16, 3737–3760. 10.1038/s41596-021-00566-6 [PubMed: 34244696]

112. Pinho SS et al. (2013) Gastric cancer: adding glycosylation to the equation. *Trends Mol Med* 19, 664–676. 10.1016/j.molmed.2013.07.003 [PubMed: 23932995]
113. Zhou F. et al. (2022) The role and potential mechanism of O-Glycosylation in gastrointestinal tumors. *Pharmacol Res* 184, 106420. 10.1016/j.phrs.2022.106420 [PubMed: 36049664]
114. Repetto O. et al. (2023) Circulating Proteins as Diagnostic Markers in Gastric Cancer. *Int J Mol Sci* 24. 10.3390/ijms242316931
115. Jeong S. et al. (2020) Glycosylation of serum haptoglobin as a marker of gastric cancer: an overview for clinicians. *Expert Rev Proteomics* 17, 109–117. 10.1080/14789450.2020.1740091 [PubMed: 32149536]
116. Zhang D. et al. (2016) Disease-specific IgG Fc N-glycosylation as personalized biomarkers to differentiate gastric cancer from benign gastric diseases. *Sci Rep* 6, 25957. 10.1038/srep25957 [PubMed: 27173519]
117. Shu J. et al. (2017) Salivary glycoproteins as potential biomarkers for diagnosis of gastric cancer. *Oncotarget* 8, 35718–35727. 10.18632/oncotarget.16082 [PubMed: 28415698]
118. Duarte HO et al. (2016) Mucin-Type O-Glycosylation in Gastric Carcinogenesis. *Biomolecules* 6. 10.3390/biom6030033
119. Pinho SS et al. (2009) Role of E-cadherin N-glycosylation profile in a mammary tumor model. *Biochem Biophys Res Commun* 379, 1091–1096. 10.1016/j.bbrc.2009.01.024 [PubMed: 19159617]
120. Pinho SS et al. (2013) E-cadherin and adherens-junctions stability in gastric carcinoma: functional implications of glycosyltransferases involving N-glycan branching biosynthesis, N-acetylglucosaminyltransferases III and V. *Biochim Biophys Acta* 1830, 2690–2700. 10.1016/j.bbagen.2012.10.021 [PubMed: 23671930]
121. Shah SC et al. (2020) Histologic Subtyping of Gastric Intestinal Metaplasia: Overview and Considerations for Clinical Practice. *Gastroenterology* 158, 745–750. 10.1053/j.gastro.2019.12.004 [PubMed: 31887261]
122. Chan AO et al. (2003) Deregulation of E-cadherin-catenin complex in precancerous lesions of gastric adenocarcinoma. *J Gastroenterol Hepatol* 18, 534–539. 10.1046/j.1440-1746.2003.02998.x [PubMed: 12702045]
123. Humar B. et al. (2009) E-cadherin deficiency initiates gastric signet-ring cell carcinoma in mice and man. *Cancer Res* 69, 2050–2056. 10.1158/0008-5472.Can-08-2457 [PubMed: 19223545]
124. Guan WL et al. (2023) Gastric cancer treatment: recent progress and future perspectives. *J Hematol Oncol* 16, 57. 10.1186/s13045-023-01451-3 [PubMed: 37245017]
125. Iqbal N and Iqbal N (2014) Human Epidermal Growth Factor Receptor 2 (HER2) in Cancers: Overexpression and Therapeutic Implications. *Mol Biol Int* 2014, 852748. 10.1155/2014/852748 [PubMed: 25276427]
126. Zhu Y. et al. (2021) HER2-targeted therapies in gastric cancer. *Biochim Biophys Acta Rev Cancer* 1876, 188549. 10.1016/j.bbcan.2021.188549 [PubMed: 33894300]
127. Kelly CM and Janjigian YY (2016) The genomics and therapeutics of HER2-positive gastric cancer-from trastuzumab and beyond. *J Gastrointest Oncol* 7, 750–762. 10.21037/jgo.2016.06.10 [PubMed: 27747089]
128. Hu Q. et al. (2023) Genomic characterization between HER2-positive and negative gastric cancer patients in a prospective trial. *Cancer Med* 12, 16649–16660. 10.1002/cam4.6269 [PubMed: 37325934]
129. Li Y. et al. (2022) Proteomic characterization of gastric cancer response to chemotherapy and targeted therapy reveals new therapeutic strategies. *Nat Commun* 13, 5723. 10.1038/s41467-022-33282-0 [PubMed: 36175412]
130. Zhang M. et al. (2022) Targeting HER3 or MEK overcomes acquired Trastuzumab resistance in HER2-positive gastric cancer-derived xenograft. *Cell Death Discov* 8, 478. 10.1038/s41420-022-01259-z [PubMed: 36463209]
131. Wang J. et al. (2024) Metabolic heterogeneity affects trastuzumab response and survival in HER2-positive advanced gastric cancer. *Br J Cancer* 130, 1036–1045. 10.1038/s41416-023-02559-6 [PubMed: 38267634]


132. He X and Xu C (2020) Immune checkpoint signaling and cancer immunotherapy. *Cell Res* 30, 660–669. 10.1038/s41422-020-0343-4 [PubMed: 32467592]
133. Muro K. et al. (2016) Pembrolizumab for patients with PD-L1-positive advanced gastric cancer (KEYNOTE-012): a multicentre, open-label, phase 1b trial. *Lancet Oncol* 17, 717–726. 10.1016/s1470-2045(16)00175-3 [PubMed: 27157491]
134. Janjigian YY et al. (2021) First-line nivolumab plus chemotherapy versus chemotherapy alone for advanced gastric, gastro-oesophageal junction, and oesophageal adenocarcinoma (CheckMate 649): a randomised, open-label, phase 3 trial. *Lancet* 398, 27–40. 10.1016/s0140-6736(21)00797-2 [PubMed: 34102137]
135. Jiang R. et al. (2024) Plasma circulating tumor DNA unveils the efficacy of PD-1 inhibitors and chemotherapy in advanced gastric cancer. *Sci Rep* 14, 14027. 10.1038/s41598-024-63486-x [PubMed: 38890392]
136. Longuespée R. et al. (2023) Molecular prediction of clinical response to anti-PD-1/anti-PD-L1 immune checkpoint inhibitors: New perspectives for precision medicine and mass spectrometry-based investigations. *Int J Cancer* 153, 252–264. 10.1002/ijc.34366 [PubMed: 36408912]
137. Sun J. et al. (2024) Proteomic profiling and biomarker discovery for predicting the response to PD-1 inhibitor immunotherapy in gastric cancer patients. *Front Pharmacol* 15, 1349459. 10.3389/fphar.2024.1349459 [PubMed: 38881867]
138. Sun JR et al. (2024) Integrated analysis of single-cell and bulk RNA-sequencing reveals a novel signature based on NK cell marker genes to predict prognosis and immunotherapy response in gastric cancer. *Sci Rep* 14, 7648. 10.1038/s41598-024-57714-7 [PubMed: 38561388]
139. Hou W. et al. (2023) Predictive Biomarkers for Immunotherapy in Gastric Cancer: Current Status and Emerging Prospects. *Int J Mol Sci* 24. 10.3390/ijms242015321
140. Li Z. et al. (2020) Multi-omics characterization of molecular features of gastric cancer correlated with response to neoadjuvant chemotherapy. *Sci Adv* 6, eaay4211. 10.1126/sciadv.aay4211 [PubMed: 32133402]
141. Zeng Y. et al. (2024) The preclinical discovery and development of zolbetuximab for the treatment of gastric cancer. *Expert Opin Drug Discov* 19, 873–886. 10.1080/17460441.2024.2370332 [PubMed: 38919123]
142. Hashimoto T. et al. (2024) Whole-transcriptome sequencing in advanced gastric or gastroesophageal cancer: A deep dive into its clinical potential. *Cancer Sci* 115, 1622–1633. 10.1111/cas.16109 [PubMed: 38429886]
143. Qiu S. et al. (2023) Targeting Trop-2 in cancer: Recent research progress and clinical application. *Biochim Biophys Acta Rev Cancer* 1878, 188902. 10.1016/j.bbcan.2023.188902 [PubMed: 37121444]
144. Chen H. et al. (2024) Molecular characterization and clinical relevance of metabolic signature subtypes in gastric cancer. *Cell Rep* 43, 114424. 10.1016/j.celrep.2024.114424 [PubMed: 38959111]
145. Akhoundova D and Rubin MA (2022) Clinical application of advanced multi-omics tumor profiling: Shaping precision oncology of the future. *Cancer Cell* 40, 920–938. 10.1016/j.ccell.2022.08.011 [PubMed: 36055231]
146. Jimenez-Santos MJ et al. (2022) Bioinformatics roadmap for therapy selection in cancer genomics. *Mol Oncol* 16, 3881–3908. 10.1002/1878-0261.13286 [PubMed: 35811332]
147. Ballman KV (2015) Biomarker: Predictive or Prognostic? *J Clin Oncol* 33, 3968–3971. 10.1200/JCO.2015.63.3651 [PubMed: 26392104]
148. Rocken C. (2023) Predictive biomarkers in gastric cancer. *J Cancer Res Clin Oncol* 149, 467–481. 10.1007/s00432-022-04408-0 [PubMed: 36260159]
149. Chao J. et al. (2021) Assessment of Pembrolizumab Therapy for the Treatment of Microsatellite Instability-High Gastric or Gastroesophageal Junction Cancer Among Patients in the KEYNOTE-059, KEYNOTE-061, and KEYNOTE-062 Clinical Trials. *JAMA Oncol* 7, 895–902. 10.1001/jamaoncol.2021.0275 [PubMed: 33792646]
150. Janjigian YY et al. (2024) First-Line Nivolumab Plus Chemotherapy for Advanced Gastric, Gastroesophageal Junction, and Esophageal Adenocarcinoma: 3-Year Follow-Up of the Phase III CheckMate 649 Trial. *J Clin Oncol* 42, 2012–2020. 10.1200/JCO.23.01601 [PubMed: 38382001]

151. Sahin U. et al. (2008) Claudin-18 splice variant 2 is a pan-cancer target suitable for therapeutic antibody development. *Clin Cancer Res* 14, 7624–7634. 10.1158/1078-0432.CCR-08-1547 [PubMed: 19047087]
152. Shah MA et al. (2023) Zolbetuximab plus CAPOX in CLDN18.2-positive gastric or gastroesophageal junction adenocarcinoma: the randomized, phase 3 GLOW trial. *Nat Med* 29, 2133–2141. 10.1038/s41591-023-02465-7 [PubMed: 37524953]
153. Shitara K. et al. (2023) Zolbetuximab plus mFOLFOX6 in patients with CLDN18.2-positive, HER2-negative, untreated, locally advanced unresectable or metastatic gastric or gastro-oesophageal junction adenocarcinoma (SPOTLIGHT): a multicentre, randomised, double-blind, phase 3 trial. *Lancet* 401, 1655–1668. 10.1016/S0140-6736(23)00620-7 [PubMed: 37068504]
154. (2024). FDA approves zolbetuximab-clzb with chemotherapy for gastric or gastroesophageal junction adenocarcinoma.
155. Mertins P. et al. (2018) Reproducible workflow for multiplexed deep-scale proteome and phosphoproteome analysis of tumor tissues by liquid chromatography-mass spectrometry. *Nat Protoc* 13, 1632–1661. 10.1038/s41596-018-0006-9 [PubMed: 29988108]
156. Meier F. et al. (2021) Trapped Ion Mobility Spectrometry and Parallel Accumulation-Serial Fragmentation in Proteomics. *Mol Cell Proteomics* 20, 100138. 10.1016/j.mcpro.2021.100138 [PubMed: 34416385]
157. Wang Y. et al. (2024) Multi-omic profiling of intraductal papillary neoplasms of the pancreas reveals distinct expression patterns and potential markers of progression. *bioRxiv*. 10.1101/2024.07.07.602385
158. Sato H and Noshiro H (2019) Report about the present condition of Billroth's stomach specimens. *Gastric Cancer* 22, 1310–1311. 10.1007/s10120-019-00987-z [PubMed: 31289946]
159. Catarci M. et al. (1997) History of reconstruction after total gastrectomy. *J R Coll Surg Edinb* 42, 73–81 [PubMed: 9114673]
160. López Sala P. et al. (2023) Gastric adenocarcinoma: A review of the TNM classification system and ways of spreading. *Radiologia (Engl Ed)* 65, 66–80. 10.1016/j.rxeng.2022.10.011 [PubMed: 36842787]
161. Joshi SS and Badgwell BD (2021) Current treatment and recent progress in gastric cancer. *CA Cancer J Clin* 71, 264–279. 10.3322/caac.21657 [PubMed: 33592120]
162. Marshall BJ and Warren JR (1984) Unidentified curved bacilli in the stomach of patients with gastritis and peptic ulceration. *Lancet* 1, 1311–1315. 10.1016/s0140-6736(84)91816-6 [PubMed: 6145023]
163. Guilford P. et al. (1998) E-cadherin germline mutations in familial gastric cancer. *Nature* 392, 402–405. 10.1038/32918 [PubMed: 9537325]
164. van Dijk EL et al. (2018) The Third Revolution in Sequencing Technology. *Trends Genet* 34, 666–681. 10.1016/j.tig.2018.05.008 [PubMed: 29941292]
165. Peng L. et al. (2019) Characterization and validation of somatic mutation spectrum to reveal heterogeneity in gastric cancer by single cell sequencing. *Sci Bull (Beijing)* 64, 236–244. 10.1016/j.scib.2018.12.014 [PubMed: 36659713]
166. Chen Y. et al. (2021) Dysregulated KRAS gene-signaling axis and abnormal chromatin remodeling drive therapeutic resistance in heterogeneous-sized circulating tumor cells in gastric cancer patients. *Cancer Lett* 517, 78–87. 10.1016/j.canlet.2021.06.002 [PubMed: 34126192]
167. Casado-Pelaez M. et al. (2022) Single cell cancer epigenetics. *Trends Cancer* 8, 820–838. 10.1016/j.trecan.2022.06.005 [PubMed: 35821003]
168. Singh H. et al. (2021) Hybrid Stomach-Intestinal Chromatin States Underlie Human Barrett's Metaplasia. *Gastroenterology* 161, 924–939.e911. 10.1053/j.gastro.2021.05.057 [PubMed: 34090884]
169. Kim J. et al. (2022) Single-cell analysis of gastric pre-cancerous and cancer lesions reveals cell lineage diversity and intratumoral heterogeneity. *NPJ Precis Oncol* 6, 9. 10.1038/s41698-022-00251-1 [PubMed: 35087207]
170. Sathe A. et al. (2020) Single-Cell Genomic Characterization Reveals the Cellular Reprogramming of the Gastric Tumor Microenvironment. *Clin Cancer Res* 26, 2640–2653. 10.1158/1078-0432.Ccr-19-3231 [PubMed: 32060101]

171. Li Y. et al. (2022) Single-cell landscape reveals active cell subtypes and their interaction in the tumor microenvironment of gastric cancer. *Theranostics* 12, 3818–3833. 10.7150/thno.71833 [PubMed: 35664061]
172. Sundar R. et al. (2022) Epigenetic promoter alterations in GI tumour immune-editing and resistance to immune checkpoint inhibition. *Gut* 71, 1277–1288. 10.1136/gutjnl-2021-324420 [PubMed: 34433583]
173. Li X et al. (2022) Single-cell RNA sequencing reveals a pro-invasive cancer-associated fibroblast subgroup associated with poor clinical outcomes in patients with gastric cancer. *Theranostics* 12, 620–638. 10.7150/thno.60540 [PubMed: 34976204]
174. Lo YH et al. (2021) A CRISPR/Cas9-Engineered ARID1A-Deficient Human Gastric Cancer Organoid Model Reveals Essential and Nonessential Modes of Oncogenic Transformation. *Cancer Discov* 11, 1562–1581. 10.1158/2159-8290.Cd-20-1109 [PubMed: 33451982]
175. Bennett HM et al. (2023) Single-cell proteomics enabled by next-generation sequencing or mass spectrometry. *Nat Methods* 20, 363–374. 10.1038/s41592-023-01791-5 [PubMed: 36864196]
176. Specht H. et al. (2021) Single-cell proteomic and transcriptomic analysis of macrophage heterogeneity using SCoPE2. *Genome Biol* 22, 50. 10.1186/s13059-021-02267-5 [PubMed: 33504367]
177. Siyal AA et al. (2021) Sample Size-Comparable Spectral Library Enhances Data-Independent Acquisition-Based Proteome Coverage of Low-Input Cells. *Anal Chem* 93, 17003–17011. 10.1021/acs.analchem.1c03477 [PubMed: 34904835]
178. Wang Y. et al. (2022) Optimized data-independent acquisition approach for proteomic analysis at single-cell level. *Clin Proteomics* 19, 24. 10.1186/s12014-022-09359-9 [PubMed: 35810282]
179. Rao A. et al. (2021) Exploring tissue architecture using spatial transcriptomics. *Nature* 596, 211–220. 10.1038/s41586-021-03634-9 [PubMed: 34381231]
180. Sundar R. et al. (2021) Spatial profiling of gastric cancer patient-matched primary and locoregional metastases reveals principles of tumour dissemination. *Gut* 70, 1823–1832. 10.1136/gutjnl-2020-320805 [PubMed: 33229445]
181. Jeong HY et al. (2021) Spatially Distinct Reprogramming of the Tumor Microenvironment Based On Tumor Invasion in Diffuse-Type Gastric Cancers. *Clin Cancer Res* 27, 6529–6542. 10.1158/1078-0432.Ccr-21-0792 [PubMed: 34385296]
182. Kelsen D. et al. (1992) FAMTX versus etoposide, doxorubicin, and cisplatin: a random assignment trial in gastric cancer. *J Clin Oncol* 10, 541–548. 10.1200/jco.1992.10.4.541 [PubMed: 1548519]

Box 1.**Proteomics approach towards clinical cancer studies**

Features of proteins, PTMs, and protein-protein interactions (PPIs) can exceed 10,000 in cancer studies [24], multi-dimensional separation approaches are essential when characterization with MS. Liquid chromatography (LC) combined with MS tandem mass spectrometry (MS/MS) is currently the most popular method used in bottom-up proteomic studies [24] (Figure I). Additional enrichments are necessary for PTM analyses [20,25,155].

**Figure I.**

LC-MS/MS systems can perform label-free quantification (via data-independent acquisition, DIA) and isobaric chemical tag labeling, such as tandem mass tag (TMT or TMTpro) for tumor samples. In TMTpro analysis, Orbitrap system can multiplex up to 18 samples in one injection, detecting over ~10,000 protein features [20], while in DIA mode, it typically detects a similar number of ~10,000 protein features [156,157]. TMT labeled and label-free samples are both suitable for deep coverage of protein quantification.

Box 2.**Technological innovations in the history of gastric cancer treatments.**

The earliest treatment milestones for GC were marked by the first successful gastric surgery in 1881 [158] and the first successful total gastrectomy in 1897 [159], both made possible by advancements in surgical instrumentation, aseptic techniques, and anesthesia. Since these surgical milestones, significant progress has been made in tumor classification, tumor staging, risk factor identification, and therapeutics for GC. In addition of histomorphological classification of GC, in 1974, the introduction of the tumor-node-metastasis (TNM) staging system for GC provided a critical framework for assessing the degree of tumor invasion, supported by developments in radiographic imaging technologies like computed tomography (CT) scans [160]. Guided by histological classification and TNM staging, surgery remained the primary treatment for GC until chemotherapy and chemoradiation combinations emerged as additional key therapeutic interventions in the 1980s [161]. Following the advent of improved endoscopic technology, endoscopic submucosal dissection was introduced into clinical practice, and *Helicobacter pylori* was successfully isolated, cultured, and linked to the GC development [27,73,162]. In the 1990s, cancer research entered the genomic era. The PCR and RT-PCR enable the detection of EBV and *CDH1* germline mutations in GC, confirming their roles as carcinogenic factors [26,163]. Subsequently, advances in NGS allowed for detailed genomic characterization of GC [13,164]. For example, TCGA and ACRG provided foundational molecular subtypes for GC classification [54,55,64]. Recently, advances in single-cell sequencing [41,42,48] and mass spectrometry [25,66,67,71], particularly with the development of Orbitrap and timsTOF technologies, have led to ground-breaking single-cell analyses and comprehensive multi-omics investigations of GC. A comprehensive understanding of current technologies and emerging trends is essential to further guide improvements in GC research and treatment. From early surgical practices to modern genomic and multi-omics approaches, technological innovations have driven the advancement of understanding and treatment strategies in GC (Table I).

Table I. Key events in understanding and treatment of gastric cancer

Year	Key Event/Finding	Supporting Technology
1881	First successful surgery	surgical instruments and aseptic techniques [158]
1897	First successful total gastrectomy	surgical techniques and anesthesia [159]
1965	Lauren histological classification	microscopy combined with histological staining [2]
1974	First TNM staging for gastric cancer	radiographic imaging (X-rays, CT scans) [160]
1980s	FAM and FAMTX chemotherapy for advanced gastric cancer	synthetic chemistry and toxicity management [182]
1988	Endoscopic submucosal dissection	endoscopic technology [2]
1990	Epstein–Barr virus infection linked to gastric cancer risk	viral DNA detection, PCR [26]

Year	Key Event/Finding	Supporting Technology
1991	<i>Helicobacter pylori</i> infection linked to gastric cancer risk	fiberoptic biopsy, endoscopic technology [27,162]
1997	ECF chemotherapy	development on combination therapy [161]
1998	CDH1 gene mutation linked to diffuse gastric cancer	gene sequencing technology (RT-PCR) [163]
2010	ToGA clinical trial and food and drug administration (FDA) approval of HER2-targeted drug trastuzumab	targeted therapy and biologics drug development [161]
2014-2015	TCGA and ACRG comprehensive genomic studies	gene sequencing (next-generation sequencing) [54,55]
2017	FDA approval of pembrolizumab (then nivolumab) for PD-L1 expressing gastric cancer	cancer immunotherapy [161]
2021	Enhertu (DS-8201) as FDA first approval ADC for gastric cancer	biologics and bioconjugation drug development [126]
2021-2022	single-cell analysis of gastric cancer	single-cell sequencing [41,42,48]
2024-2025	Clinical proteomic tumor analysis consortium (CPTAC) comprehensive multi-omics characterization of gastric cancer	mass spectrometry [71]

Outstanding questions

- How can multi-omics technologies comprehensively capture the molecular and cellular heterogeneity of gastric cancer, facilitating earlier detection across its diverse subtypes to improve prevention, stratification, and risk assessment for better clinical outcomes?
- Can proteogenomic analyses provide more precise insights into the identification of intracellular and cell surface targets, enabling the development of subtype-specific therapeutic strategies for gastric cancer and overcoming treatment resistance?
- Which advancements in single-cell technologies can most effectively reveal the complexity of gastric cancer at the cellular level, and how can these innovations be leveraged to transform early diagnosis, treatment personalization, and prognosis prediction?
- Are these technologies in molecular characterization, such as integrated omics approaches, that can revolutionize the development of targeted biologics and personalized therapies, particularly in addressing the unmet clinical needs for gastric cancer?
- How can AI and machine learning approaches be applied to integrate multi-omics and clinical data to predict disease characteristics, model therapeutic responses, and enhance personalized treatment plans for gastric cancer?

Highlights

- Gastric cancer is a complex and highly heterogeneous disease, presenting significant challenges in treatment due to its molecular diversity and variations in coding and non-coding mutations. The complexities make it difficult to develop universally effective therapies.
- Recent technological advancements, including proteomics, proteogenomics, microbiome profiling, and single-cell analyses with spatial information, have greatly improved our capability to understand gastric cancer biology.
- The integration of multi-omics technologies offers new opportunities for precision treatment. These approaches not only directly identify therapeutically targetable proteins and their modifications, but also pave the way for more refined and personalized cancer treatments with greater promise.




Figure 1. overview of technology driven gastric cancer study. (A) Features of emerging technologies towards GC study. Genomic analysis enables the measurement of gene expression and genomic variations, providing insights into gene regulation and the identification of distinct cell types and states through RNA-seq data. Proteomic analysis directly analyzes functional targets, tracks changes in signaling pathways, and uncovers molecular alterations associated with therapeutic responses. Single-cell analysis identifies signature markers specific to cell types and states, tracks cell differentiation and developmental trajectories, and characterizes gene variants across diverse cell populations. Microbiome and metabolomic analyses elucidate the composition of the microbiota and identifies metabolites derived from both tissues and the microbiome, offering a comprehensive view of host-microbiome interactions and metabolic processes. (B) The component for emerging technologies. Sample quality is a critical factor influencing data reliability, as degradation of RNA and protein modifications especially phosphorylation during collection and processing can hinder analysis, particularly in spatial analysis, which requires intact, well-preserved tissue samples. Additionally, cost remains a significant barrier, especially for high-throughput multi-omics and single-cell technologies, which are often high cost compared to other studies.




Figure 2.
key findings of omics based gastric cancer studies.

Table 1. Single-cell omics technologies in gastric cancer

Single-cell omics	Features	Applications in gastric cancer
Single-cell genome sequencing	This technique enables the analysis of the genomic DNA of individual cells, revealing genetic variations and alterations within tumors. It has been used to identify mutated genes, track clonal evolution and understand the genetic basis of GC.	Characterizations were made in gastric cancer cell line [47] and tumor tissue [165,166].
Single-cell epigenome sequencing	This method focuses on analyzing the epigenetic modifications, such as DNA methylation and histone modifications, in individual cells. It provides insights into the regulation of gene expression and the epigenetic changes in GC.	Several achievements were made for this technology [167] and used in the measurement of metaplasia [168].
Single-cell RNA sequencing (scRNA-seq)	This technology allows for the profiling of gene expression in individual cells, providing insights into cellular heterogeneity and molecular landscapes. It has been used to study tumor heterogeneity and the TME. When integrated with CRISPR-based lineage tracing, it provides a powerful framework to trace the trajectories of tumor progression with unprecedented resolution.	The studies were completed on metastatic GC [41], all stages and subtypes with paired normal samples [42,169], TME [48,170-172], and tumor progression [34,48,65,173]. Limited studies were published on the joint application of CRISPR-based lineage tracing and scRNA-seq on GC [174].
Single-cell proteomics (sc-proteomics)	This technology allows for the analysis of protein expression at the single-cell level. It provides information about the protein profiles and signaling pathways within individual cells, contributing to the understanding of cellular dynamics and disease mechanisms.	sc-proteomics could be achieved by either MS or NGS [175]. MS techniques are more common in sc-proteomics including DDA [176] or DIA approach [177,178]. However, no study was reported in GC as of now.
Single-cell multiomics	This technology integrates multiple layers of omics information, at the single-cell level. It allows for a holistic understanding of cellular states, signaling networks, and interactions, offering deeper insights into tumor heterogeneity and the TME.	The study contained genomic, epigenomic, and transcriptomic analysis from single cell, uncovering intratumoral heterogeneity and differentiation states in gastric cancer [51].
Spatial transcriptomics (ST)	ST technologies provide information about the spatial distribution of gene expression within tissues. ST techniques have been used to study spatial heterogeneity in tumors and understand the spatial organization of gene expression in GC.	They included imaging-based and sequencing-based techniques [179]. Currently, sequencing-based <i>in situ</i> barcoding (ISB) is the most common approach in GC studies [40,42,43,180,181].
Spatial multi-omics	This technology integrates spatially resolved omics data with tissue architecture, to map cellular and molecular features directly within their native spatial context. It enables the exploration of cell-cell interactions and the TME with unprecedented resolution.	The study integrated spatial information from MS imaging-based metabolomics and lipidomics with microarray-based transcriptomics to reveal the metabolic remodeling feature [43,45].