

Vascular thiol isomerases

Robert Flaumenhaft and Bruce Furie

Division of Hemostasis and Thrombosis, Department of Medicine, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA

Thiol isomerases are multifunctional enzymes that influence protein structure via their oxidoreductase, isomerase, and chaperone activities. These enzymes localize at high concentrations in the endoplasmic reticulum of all eukaryotic cells where they serve an essential function in folding nascent proteins. However, thiol isomerases can escape endoplasmic retention and be secreted and localized on plasma membranes. Several thiol isomerases including protein disulfide isomerase,

ERp57, and ERp5 are secreted by and localize to the membranes of platelets and endothelial cells. These vascular thiol isomerases are released following vessel injury and participate in thrombus formation. Although most of the activities of vascular thiol isomerases that contribute to thrombus formation are yet to be defined at the molecular level, allosteric disulfide bonds that are modified by thiol isomerases have been described in substrates such as α IIb β 3, α v β 3, GPIIb α , tissue

factor, and thrombospondin. Vascular thiol isomerases also act as redox sensors. They respond to the local redox environment and influence S-nitrosylation of surface proteins on platelets and endothelial cells. Despite our rudimentary understanding of the mechanisms by which thiol isomerases control vascular function, the clinical utility of targeting them in thrombotic disorders is already being explored in clinical trials. (Blood. 2016; 128(7):893-901)

Introduction to thiol isomerases

In the early 1960s, Anfinsen et al described the ability of the inactive, denatured form of ribonuclease to refold into its native form and express full enzymatic activity.¹ An enzyme from liver, called disulfide interchange enzyme, reshuffle, or rearrangease,² was identified as facilitating the refolding of ribonuclease by forming the disulfide bonds to obtain the lowest energy structure.³ This enzyme came to be known as protein disulfide isomerase (PDI; EC.5.3.4.1) and was established as a critical component of the biosynthetic pathway for proteins.⁴ PDI, present in the endoplasmic reticulum, is the prototype thiol isomerase and has been extensively studied.⁵

All thiol isomerases, of which now there are 21 in the PDI family, have evolved from an ancestral thioredoxin found in primitive unicellular organisms. PDI is the archetypal and most abundant thiol isomerase (Figure 1). These proteins contain a thioredoxin-like domain, which consists of a 4-stranded antiparallel β sheet surrounded by 3 α helices (Figure 1A).^{6,7} PDI family thiol isomerases have an active site motif Cys-X-X-Cys (CXXC) where X is any amino acid. Duplication and evolution of the thioredoxin-like fold⁸ resulted in multidomain thiol isomerases of the PDI family that contain both catalytic domains (termed a, a', or a⁰ by convention), which contain the CXXC motif, and substrate-binding domains (termed b or b' by convention). Catalytic domains contain the CXXC motif and perform oxidoreductase functions (Figure 2). Although the a and a' domains are homologous, they are not functionally redundant. For example, only the a' domain of PDI is critical for thrombus formation.⁹ The substrate binding domains have lost the catalytic motifs and developed features that facilitate protein-protein interactions such as the hydrophobic pocket on PDI (Figure 1B).¹⁰

Depending on the context, thiol isomerases function as reductases, oxidases, or isomerases (Figure 3). Thiol isomerases can act as oxidases when engaging a thiol-containing substrate with a lower redox potential, forming disulfide bonds within the substrate (Figure 3A, reaction from left to right). Acting as a reductase, thiol isomerases

cleave disulfide bonds within substrate proteins (Figure 3A, reaction from right to left). They can also act as an isomerase, changing protein structure by modified disulfide bond formation without a net change in free thiol content (Figure 3B). By facilitating protein folding in the absence of catalytic activity, they can act as chaperones (Figure 3C).^{11,12} Modifications of the catalytic cysteines enable thiol isomerases to act as nitrosylases/denitrosylases, transferring nitric oxide (NO) to or from substrates, respectively (Figure 3D).¹³⁻¹⁵ Variations in vascular NO levels may drive these S-nitrosylation reactions. The catalytic cysteines of thiol isomerases can also react with glutathione and other redox active plasma species.^{16,17} Thus, thiol isomerases are multidomain, multifunctional proteins.

Thiol isomerases as redox sensors

Effect of redox environment on thiol isomerases

The activity of cell surface thiol isomerases is controlled by the redox environment. In a reducing environment, the catalytic cysteines within the active site motif are reduced and thiol isomerases act as reductases. Under oxidizing conditions, the catalytic cysteines will form a disulfide bond and thiol isomerases act as oxidases. The main redox buffers in the vasculature are the glutathione system (GSH/GSSG) and the cysteine system (Cys/CySS) and these determine the redox potential of plasma.¹⁸ Plasma redox potentials (E_h) are -130.9 ± 22.9 mV for GSH/GSSG and -72.4 ± 12.8 mV for Cys/CySS.¹⁹ The active site cysteines in PDI have a redox potential of -190 ± 10 mV.²⁰ Based on these considerations, one would expect vascular PDI to be primarily oxidized in the extracellular environment. However, redox conditions can vary locally. In addition, protein disulfide oxidases such as ERO1 α , which undergoes activation-dependent cell surface expression in platelets,²¹ and protein disulfide reductases can enzymatically modify




Figure 1. The architecture of thiol isomerases. Thiol isomerases consist of tandem domains containing a thioredoxin-fold. (A) The structure of the thioredoxin-fold.⁷ (B) PDI is formed from 4 tandem domains containing thioredoxin-like folds that assume a U-shape. The 2 catalytic domains (a and a') contain the CGHC motifs that contain the catalytic cysteines (shown in red). The catalytic cysteines face one another and are responsible for oxidoreductive activities. The substrate binding domains (b and b') form the bottom of the U-shape. The b' domain contains a hydrophobic binding pocket that is primarily responsible for substrate binding. The b' and a' domains are connected by a flexible 19-amino-acid peptide termed the x-linker.

catalytic cysteines within active site motifs.²² Little is known about the function of enzymatic control of redox states of extracellular vascular thiol isomerases.

Thiol isomerases are sensitive to changes in redox potential, undergoing substantial conformational changes in response to modifications the redox environment. In reducing conditions, PDI assumes a compact structure (Figure 4A-B).²³ The conformational change is mediated by reduction of the catalytic cysteines in the CGHC motif, which triggers rotation of Trp-396 on the a' domain, enabling it to interact with Arg-300 on the b' domain (Figure 4C). This switch initiates a series of interactions at the b'-a' interface that positions the a' domain over the hydrophobic binding site on the b' domain.^{23,24} This constrained conformation is thought to be a means to limit substrate binding under reducing conditions.²⁴ Under oxidizing conditions, the constraint of disulfide bond formation between Cys-397 and Cys-400 prevents Trp-396 from interacting with Arg-300 and the hydrophobic binding site on b' remains exposed and facilitating interactions with substrates. PDI is thus a more effective chaperone under oxidizing conditions,²⁴ but can also engage substrates under reducing conditions. Recent studies show that on binding a substrate, PDI assumes a more compact conformation.²⁵ Substrate binding at the hydrophobic pocket activates an allosteric switch involving the x-linker, which results in enhanced reductase activity in the a and a' domains and appears to be a cooperative mechanism whereby substrate engagement activates enzymatic activity.

Generating surface free thiols

Activation of platelets results in a greater than fourfold increase in surface protein free thiols.²⁶ Although the percentage of active sites that are dithiols is only 26% in resting platelets, the number increases to

81% in activated platelets.²⁶ PDI is prominent among the sources of activation-dependent free thiols. Free thiols within catalytic domains have several potential functions in addition to the oxidoreductive functions discussed above. They could react with reactive oxygen species, thereby buffering platelets from oxidant damage. Free thiols within the catalytic domains of PDI could also act as denitrosylases, activating S-nitrosylated proteins, although this function has not been well characterized.

In endothelium, reaction of free thiols with oxidants and free radicals is thought to protect endothelium from oxidant damage. Several thiol isomerases including PDI, ERp46, and ERp5 are oxidized on incubation of endothelial cells with the neutrophil-derived oxidant HOCl.²⁷ Similarly, thiol isomerases, particularly thioredoxin, are capable of regenerating protein inactivated by oxidative stress.^{28,29} Thiol isomerases on endothelium expose a free thiol (Cys326) on the surface of β_2 GPI and β_2 GPI reduced by endothelial thiol isomerases protects endothelium from cell death induced by oxidative stress.³⁰

S-nitrosylation of thiol isomerases

Generation of NO by endothelial cells is an essential mechanism for maintaining vascular quiescence. NO produced by endothelium acts on platelets to maintain them in a resting state. Exposure to NO results in a decreased number of free thiols on megakaryocyte and platelet surfaces³¹ and S-nitrosylation of multiple platelet receptors including $\alpha_{IIb}\beta_3$ is observed following exposure to NO.³² PDI undergoes S-nitrosylation following exposure to NO donors.³³ In contrast to S-nitrosylation at regulatory cysteines within platelet receptors such as $\alpha_{IIb}\beta_3$,³⁴ S-nitrosylation of catalytic cysteines within PDI enables transfer of NO to other proteins and into cells. Several lines of evidence demonstrate this transnitrosylase activity of PDI. Delivery of NO from an S-NO donor into cytosol is inhibited following knockdown of PDI in HEL cells.¹³ Inhibition of PDI blocks the delivery of NO from the physiologic NO donor, S-nitrosoglutathione, into both megakaryocytes³¹ and into platelets.³⁵ PDI-mediated transnitrosylation is proposed to occur via the delivery of NO into cell membranes in the form of N_2O_3 .¹⁴ Intracellular substrates are S-nitrosylated as they interact at the membrane-cytosol interface.¹⁵ NO can be transferred from PDI to the heme moiety in guanylyl cyclase, as evidenced by its activation and marked increased in guanosine 3',5'-cyclic monophosphate (cGMP) levels.^{13,35} PDI could also transfer NO to surface proteins¹⁴ including platelet receptors, maintaining them in an inactivated conformation. Thiol isomerase-mediated S-nitrosylation of surface proteins could therefore function in maintaining vascular quiescence (Figure 5). However, the significance of this mechanism in vivo has not yet been tested.





Figure 2. Mechanism of oxidoreductive cleavage by thiol isomerases. The reduced thiol isomerase aligns with a substrate that contains a disulfide bond such that the free thiolate forms an 180° angle with the vector of the disulfide. A second-order nucleophilic substitution S_N2 -type reaction then occurs in which the active site sulfur ion nucleophile of the thiol isomerase attacks the adjacent sulfur atoms of the disulfide bond.⁷⁴ This nucleophilic substitution results in a transient mixed disulfide (blue). The mixed disulfide spontaneously decomposes, resulting in the formation of a disulfide bond in the thiol isomerase and reduction of the disulfide bond in the substrate.

Figure 3. Multiple functions of PDI. (A) PDI is an oxidoreductase that can both form (oxidase) and cleave (reductase) disulfide bonds. Active site cysteines are shown in red. Note that the catalytic cysteines are either reduced or oxidized during these processes. (B) Through its oxidoreductase activity, PDI acts as an isomerase changing disulfide bond patterns within a protein. In this example, PDI modifies disulfides formed between cysteines 1-2 and 3-4 into disulfides formed between 1-3 and 2-4 (numbering in red). Note that there is no net change in the redox state of the catalytic cysteines on PDI. (C) PDI can act as a chaperone, modifying the conformation of a protein independent of its oxidoreductase and chaperone activities. The redox state of the catalytic cysteines is unchanged. (D) The active site cysteines within PDI can also become *S*-nitrosylated. PDI can then transfer NO to substrates, thus acting as a nitrosylase. PDI can also act as a denitrosylase, removing NO from substrates. *S*-nitrosylation typically inactivates substrates, whereas denitrosylation activates substrates.



Vascular thiol isomerases

A subset of thiol isomerases is found in the vasculature (Table 1)^{36,37} and some of these, including PDI, ERp57, and ERp5, have known functional importance on the surfaces of platelets and endothelial cells (Figure 6). The first report of a thiol isomerase being released from platelets was in the early 1990s, when Chen et al showed that PDI is released by activated platelets.³⁸ The Essex laboratory subsequently showed that PDI localizes to the platelet membrane^{39,40} and functions in platelet activation.⁴¹ Several other groups subsequently demonstrated that thiol isomerases traffic to secretory granules/membrane systems and are released from platelets in an activation-dependent manner.^{36,42-45} Kim et al showed that platelets lacking PDI have reduced activation-dependent aggregation.⁴⁶ Platelet surface PDI has been postulated to modulate $\alpha_{IIb}\beta_3$ function,⁴⁷⁻⁴⁹

collagen binding to $\alpha_2\beta_1$,⁵⁰ GP1b α ,²⁶ thrombospondin 1,⁵¹⁻⁵³ and vitronectin.⁵⁴ ERp5⁵⁵ and ERp57^{56,57} also localize to the platelet surface and their inhibition blocks platelet activation. The precise mechanisms by which thiol isomerases function in the activation of $\alpha_{IIb}\beta_3$ or other platelet receptors, however, are not well understood.

Thiol isomerases have also been found on the surface of the endothelium. Cell-impermeable probes capable of reacting with closely spaced thiols on cell surfaces have been used to identify PDI on the surface of endothelial cells.^{15,58} Subsequent studies showed that PDI forms a complex with $\alpha_v\beta_3$ on the endothelial cell membranes, providing a mechanism by which it is retained on these membranes.⁵⁹ This endothelial cell surface PDI is capable of activating both β_1 and β_3 integrins.^{59,60} Cell surface PDI has also been shown to regulate phosphatidylserine exposure,⁶¹ as well as nitric oxide delivery on endothelial cells.^{15,62} Multiple endothelial cell agonists including




Figure 4. Redox sensitivity of PDI. (A) In a reducing environment, PDI assumes a compact structure characterized by a 15-Å distance between the a and a' domains. (B) In an oxidizing environment, the distance between the a and a' domain increases to 30 Å. The increase in distance results from loss of interactions between the b' and a' domains that allows movement of the x-linker and a' domains. Adapted from Wang et al.²³ (C) In a reducing environment, the disulfide bond between C397 and C400 is broken, enabling the rotation of W396. This rotation positions W396 so that it can interact with R300. The interaction between W396 and R300 brings the a' domain in closer proximity to the b' domain, facilitating additional interactions that lead to a more compact structure. Adapted from Wang et al.²³

thrombin, histamine, and phorbol 12-myristate 13-acetate have been shown to stimulate PDI release from endothelial cells, and PDI is released from endothelium *in vivo* following laser injury.⁶³ However, it was not until the function of thiol isomerases was evaluated in the context of thrombus formation that the *in vivo* importance was appreciated.

Thiol isomerases in thrombus formation

The discovery that thiol isomerases participate in thrombus formation was accidental and facilitated by the intersection of the development and application of high-speed multicolor digital intravital microscopy to the study of thrombus development in a live mouse,⁶⁴ the hypothesis of Hogg et al that the mechanism of tissue factor deencryption involved oxidation of 2 free sulphhydryls and formation of a disulfide bond,⁶⁵ and the suggestion that PDI might be the enzyme that catalyzes the formation of that disulfide bond.⁶⁶ Although subsequent efforts to examine the oxidation of tissue factor following tissue injury in *in vivo* systems neither proved nor disproved the Hogg hypothesis, several studies showed that platelets and endothelium secrete PDI on tissue injury in a live animal.^{63,67} That PDI, important for disulfide bond formation during protein synthesis, is released from vascular cells and plays an important extracellular role was in itself a surprising finding. However, the observation that inhibition of PDI with blocking antibodies demonstrated inhibition of thrombus formation provided direct evidence that PDI is required for thrombus formation after tissue injury.⁶⁷ The importance of PDI in thrombus formation has now been demonstrated in several models of thrombus formation including a laser injury model,⁶⁷ a ligation model,⁶⁸ and a ferric chloride model (Table 2).^{69,70}

Since the discovery of the importance of PDI in thrombus formation, similar observations have been extended to other thiol isomerases, including ERp57 and ERp5 (Table 2).^{45,56,57} ERp57, with a domain organization parallel to PDI, is also secreted by platelets and endothelial cells on cell activation. Three groups have related ERp57 to thrombus formation using (1) blocking antibodies to ERp57⁵⁶; (2) genetic alterations of megakaryocyte/platelet PDIA3,⁷¹ the gene that encodes ERp57; and (3) *in vivo* morpholinos in mice to decrease expression of ERp57.⁷⁰

ERp5 is yet another thiol isomerase whose inhibition blocks thrombus formation. Using polyclonal monospecific, immunoaffinity purified antibodies to ERp5 antibodies *in vivo* in the laser injury model in mice, Passam et al showed that inhibition of ERp5 blocked thrombus formation including both platelet aggregation and fibrin generation.⁴⁵ ERp5 is derived from both platelets and endothelial cells on cell activation with vascular injury. ERp5 binds to $\alpha_{IIb}\beta_3$ but does not bind through the active site domain.

In summary, a series of thiol isomerase has been identified in the vessel wall and in platelets that, on cell activation, stimulates activation of extracellular thiol isomerase activity⁷² and accompanies thrombus formation *in vivo*.^{45,67} Each of these thiol isomerases is either absolutely required or partially required for thrombus formation. In addition to these *in vivo* murine models, a role for thiol isomerases in platelet activation and thrombus formation has also been shown *in vitro* using human samples.⁷² Several surprises can be noted. First, these thiol isomerases reside in the endoplasmic reticulum and, in the case of PDI, also the T-granule⁴³ and yet have extracellular function; second, they have important function with regard to thrombus formation. An initiation pathway requiring thiol isomerases for thrombus formation has never been previously described.

The master switch hypothesis

As bits and pieces of this new system were discovered, it was difficult to synthesize these observations to understand their biological function. What is the pathway in which inhibition of PDI or ERp57 or ERp5 individually block both platelet activation and fibrin generation *in vivo*? Are these thiol isomerases acting in series in the pathway? That is, is there an electron transport chain in which thiol isomerases act in a linear fashion as enzyme and substrate in a redox-active manner? Or do they act in parallel, with each of the vascular thiol isomerases acting on a unique set of substrates? Are the actions of thiol isomerases on substrates only about changes in the covalent structure of disulfide bonds or are the other functions of thiol isomerases involved?

With only fragments of information available about this new pathway(s), we considered that the intravascular release of these thiol isomerases primed the triggers for initiating thrombus formation. We




Figure 5. Proposed model for the role of PDI in regulating vascular NO activity. In the quiescent vasculature, when NO levels are high, PDI may be S-nitrosylated. Endothelial SNO-PDI could transfer NO to surface proteins, thereby inhibiting their activity, or transfer NO intracellularly. Intracellular NO could bind to guanylyl cyclase, stimulating the generation of cGMP and contributing to the platelet quiescence.

posit that (1) vascular thiol isomerases are intracellular under normal conditions; (2) thiol isomerases are released on tissue damage or cellular activation; (3) only on release of thiol isomerases can thrombus formation be initiated; and (4) extracellular thiol isomerases catalyze changes in vascular surface receptors and plasma protein structure. In essence, this master switch hypothesis postulates that encoded into many vascular proteins are functional disulfide bonds that lock the proteins in an inactive state. Such allosteric disulfides have previously been described in vascular proteins and proposed to function in thrombus formation.^{73,74} These allosteric disulfides could form an embedded network of clamps that prevents the untoward initiation of thrombus formation. Thiol isomerases, released from activated or injured cells, are required to deactivate this embedded network of safeguards so as to enable blood coagulation and platelet activation. An essential component of this network is that its activation is required for coagulation and platelet activation, but it is not itself capable of eliciting either.

The specific substrates of thiol isomerases and identification of cysteines involved in clamping vascular proteins in an inactive state are largely undefined. Yet some examples are illustrative. PDI binds to platelet $\alpha_{IIb}\beta_3$ ⁴⁸ and platelets deficient in PDI demonstrate impaired $\alpha_{IIb}\beta_3$ activation.⁴⁶ Although the disulfides on $\alpha_{IIb}\beta_3$ that are targeted by PDI have not been identified, regulatory disulfides in the epidermal growth factor domains have been implicated.^{47,75} PDI interacts with $\alpha_v\beta_3$ on endothelial cells⁵⁹ and $\alpha_M\beta_2$ on neutrophils.⁷⁶ ERp5^{45,55} and ERp57^{56,71} also bind to and modify $\alpha_{IIb}\beta_3$ activity. Thrombospondin 1 is another thiol isomerase substrate. PDI has previously been found to form disulfide linked complexes with thrombospondin 1 from platelets⁵¹ and expose a cryptic RGD sequence that facilitates the association of thrombospondin 1 with endothelial cell $\alpha_v\beta_3$.⁵³ PDI has been proposed to activate tissue factor by forming and allosteric bond between Cys186 and Cys209.^{65,77} Although intriguing, this activation mechanism remains controversial.^{61,78} Overall, the importance of these thiol isomerase–substrate interactions in thrombus formation remains to be proven in an in vivo model. However, the hypothesis that distinct thiol isomerases act on an overlapping but distinct set of substrates to prime the vasculature for thrombus formation following vessel injury or activation is consistent with current observations.

Clinical implications of thiol isomerases in hemostasis and thrombosis

The modification of functional disulfide bonds by thiol isomerases represents a previously unrecognized layer of control of vascular function in general and specifically thrombosis. Modification of allosteric disulfide bonds by thiol isomerases is akin to cleavage of amino acid bonds by serine proteases of the coagulation cascade. Yet fundamental distinctions exist between these 2 mechanisms that have important clinical implications. Unlike proteases of the coagulation pathway, active thiol isomerases do not directly cause blood to coagulate. Thus, thiol isomerases could support hemostasis in coagulopathies without the thrombotic risks associated with infusion of coagulation proteins (eg, FEIBA, FVIIa). Conversely, because thiol isomerases control thrombus formation, but are not the final effectors of clotting, inhibition of thiol isomerases could achieve control of thrombosis without concurrent bleeding risks. In fact, genetic deletion of megakaryocyte (MK)/platelet PDI impairs thrombosis without prolonging tail bleeding times.⁴⁶ The extremely high abundance of intracellular thiol isomerases (200 μ M in endoplasmic reticulum)⁷⁹ renders likely the possibility that diseases characterized by increased cell proliferation and/or cell death will show elevated thiol isomerase levels. Chronic exposure to elevated thiol isomerase levels may predispose to thrombosis. These distinguishing characteristics of thiol isomerases compared with serine proteases have important implications with regard to the clinical utility of thiol isomerase-targeted therapeutics and diagnostics.

Table 1. Properties of vascular thiol isomerases

Thiol isomerase	Gene name	Chromosomal location	Amino acids	Molecular mass (kDa)	Domain structure	Catalytic motif
PDI	<i>P4HB</i>	17q25	508	57.1	a-b-b'-a'	CGHC
ERp57	<i>PDIA3</i>	15q15	505	56.8	a-b-b'-a'	CGHC
ERp5	<i>PDIA6</i>	2p25.1	440	48.1	a-a-b	CGHC
ERp46	<i>TXNDC5</i>	6p24.3	432	47.6	a ⁰ -a-a'	CGHC
ERp72	<i>PDIA4</i>	7q35	645	72.9	a ⁰ -a-b-b-a	CGHC
ERp44	<i>ERP44</i>	9q22.33	406	46.9	a-b-b'	CGHC
ERp29	<i>ERP29</i>	12q24.13	261	29.0	b-b'	NA




Figure 6. Domain structures of vascular thiol isomerases. (A) The domain structures of thiol isomerases with known activity in thrombus formation (PDI, ERp57, and ERp5) are shown. Catalytic domains (tan), binding domains (blue), and c domains (green) are indicated. The location of the CGHC motif is indicated in red. Numbers indicate the amino acid position in the mature protein.

Inhibition of thiol isomerases

Thiol isomerases function in coagulation primarily to modify substrates on endothelial cells, platelets, and in plasma to create conditions conducive for coagulation. Inhibition of thiol isomerases interferes with this priming mechanism and blocks thrombus formation in animal models. As described above, antibodies directed against PDI,⁶⁷ ERp57,⁵⁶ or ERp5⁸⁰ inhibit thrombus formation. Small molecules targeting ERp57 such as ADTM⁸¹ or compounds targeting PDI such as the flavonoid quercetins (eg, quercetin-3-rutinoside and isoquercetin) also inhibit thrombus formation.⁶⁹ When tested in mice, quercetin-3-rutinoside inhibited thrombus formation at doses as low as 0.5 mg/kg. Infusion of recombinant PDI reversed the inhibitory effect of quercetin-3-rutinoside in mice, indicating that PDI was the relevant target of quercetin-3-rutinoside in this thrombosis model. Orally administered quercetin-3-rutinoside was also antithrombotic.⁶⁹ Importantly, quercetin-3-rutinoside blocked injury-induced thrombus formation at concentrations commonly used by individuals who consume quercetin-3-rutinoside supplements. Epidemiologic studies indicate that consumption of flavonoids is associated with reduced risk of cardiovascular death and decreased incidence of nonfatal and fatal stroke.^{82,83}

Isoquercetin, a flavonoid quercetin that is more readily absorbed than quercetin-3-rutinoside, is currently being evaluated as an antithrombotic in the Cancer Associated Thrombosis and Isoquercetin (CAT-IQ) trial. The CAT-IQ study is a 600-patient, randomized, placebo-controlled, phase 2/3 clinical trial (#NCT02195232) designed to determine the efficacy and safety of isoquercetin in preventing cancer-related thrombosis.⁸⁴ The efficacy outcome measurements include thrombosis at 60 days and D-dimer levels. Bleeding will be monitored to assess safety. As with any antithrombotic, it will be valuable to identify assays capable of monitoring the anticoagulant effect of PDI-targeted therapies. Ancillary studies will

assess the PDI inhibitory activity in plasma to determine whether adequate plasma concentrations of isoquercetin or its metabolites are reached. Although inhibition of PDI does not affect standard coagulation studies, it does have some effect on platelet aggregation^{41,85} and platelet-dependent thrombin generation.⁸⁶ Successful completion of the CAT-IQ trial will provide proof-of-principle for inhibition of PDI in primary prophylaxis of thrombosis and allow us to develop methods to monitor PDI-targeted therapies.

Thiol isomerases in coagulopathy

If thiol isomerases do not initiate thrombosis themselves, but rather prime the vasculature for clot formation, then it is possible that they will be effective in promoting hemostasis in the setting of coagulopathy. Similarly, it is possible that lack of thiol isomerase release contributes to coagulopathy. The latter premise is supported by studies of Hermansky-Pudlak syndrome (HPS). HPS is characterized by oculocutaneous albinism and a deficiency in platelet dense granules.⁸⁷ Mice with HPS have prolonged bleeding times and impaired thrombus formation, including impaired fibrin formation, on vascular injury. Platelets and endothelial cells from HPS mice are deficient in the secretion of PDI.⁴⁴ Furthermore, infusion of recombinant PDI reverses the thrombus formation defect in HPS mice (S. H. Kim and B. Furie, unpublished observations, 2013). Whether recombinant PDI shortens bleeding times in mice has not yet been determined. Nor is it known whether infusion of recombinant thiol isomerases can improve bleeding in other coagulopathies.

Selecting candidate diseases for thiol isomerase-targeted therapy

Our immature understanding of the mechanisms by which thiol isomerases control thrombosis complicates identification of optimal clinical settings for thiol isomerase-targeted therapy. If our underlying premise is that thiol isomerases work on multiple substrates to create a prothrombotic state, then thiol isomerase-targeted therapies could be antithrombotic in almost any hypercoagulable condition. Such therapies could be most effective in disease states in which baseline thiol isomerase levels are elevated. Currently, there are very little data on circulating levels of vascular thiol isomerases other than thioredoxin. Normal ranges for any of the vascular thiol isomerases shown to function in thrombosis (PDI, ERp5, ERp57) have not been established. Autoantibodies directed against these enzymes, however, indicates that these thiol isomerases circulate at increased levels in certain disease states. Antibodies to PDI have been identified in cancer and SLE,⁸⁸ as well as in infection.⁸⁹ ERp57 autoantibodies have also been found in

Table 2. Thiol isomerases in thrombus formation

Thrombosis model	Mode of inhibition of TI	Effect on thrombus formation	Reference
Laser-induced injury cremaster arteriole	Anti-PDI antibody (RL90)	Minimal platelet accumulation No fibrin generation	67
Ligation Model	Anti-PDI antibody	~70% reduction in fibrin formation at 30 minutes	68
Laser-induced injury cremaster arteriole	Quercetin-3-rutinoside, 0.5 µg/kg	Near absence of platelet accumulation and fibrin generation	69
FeCl ₃ -induced injury	Quercetin-3-rutinoside	2.2-fold increase in occlusion time	69
Laser-induced injury cremaster arteriole	Platelet/MK-specific PDI null	~75% inhibition of peak platelet accumulation	46
FeCl ₃ -induced carotid injury	Platelet/MK-specific PDI null	Two-fold increase in occlusion time	9
FeCl ₃ -induced carotid injury	Mice expressing a mutant PDI (CGHC/SGHS)	~1.5-fold increase in occlusion time	9
FeCl ₃ -induced carotid injury	Anti-ERp57 antibody (Mab1)	Approximately threefold increase in occlusion time	57
Laser-induced injury cremaster arteriole	Anti-ERp57 antibody	~80% reduction in platelet accumulation	56
FeCl ₃ -induced carotid injury	Platelet/MK-specific ERp57 null	~1.9-fold increase in occlusion time	71
Laser-induced injury cremaster arteriole	Anti-ERp5 antibody	70% decrease in platelet accumulation 62% decrease in fibrin generation	45




Figure 7. Functional consequences of thiol isomerase-mediated disulfide bond modifications. Thiol isomerases are secreted from platelets and endothelium following cell activation and mediate modifications of functional disulfide bonds in multiple substrates. Several categories of modifications are observed. (1) Exposure of binding sites in adhesion proteins. Cleavage of disulfide bonds can reveal cryptic binding sites such as RGD sequences (shown in blue) enabling a nonbinding adhesion protein (red) to adhere to binding partners on the cell surface (green). Thrombospondin is an adhesion protein that is modified in this manner.⁵³ (2) Activation of enzyme function. Functional disulfide bond cleavage could also expose an encrypted active site within an enzyme or modify its conformation, converting the enzyme to a more active conformation. Conversely, formation of a disulfide bond could activate an enzyme or coenzyme (eg, tissue factor).^{65,66,93} (3) Receptor activation. Disulfide bond shuffling could contribute to the formation or stabilization of the active conformation of cell surface receptors (eg, $\alpha_{IIb}\beta_3$).^{47,75} (4) Release from binding protein. Cleavage of a disulfide bond could release a protein from its binding partner, thereby activating the protein (eg, TGF- β).⁹⁴

cancer,^{90,91} and infection.⁹² Thiol isomerases that bind to cell surfaces could be detected by the immune system and generate autoantibodies even if circulating levels are undetectable.

The possibility that thiol isomerase inhibition does not substantially enhance bleeding risk suggests additional indications. Currently, primary prophylaxis is not offered for certain thrombotic risk factors because the risk of bleeding from the antithrombotic outweighs the protective benefit (eg, heterozygous factor V Leiden with single thrombosis, antiphospholipid antibody without thrombosis). Thiol isomerase-targeted therapies could provide protection from thrombosis for these conditions. Because thiol isomerase inhibition acts by a mechanism other than direct inhibition of coagulation factor synthesis or activity, it is possible that the activity of the two approaches will be synergistic. The addition of thiol isomerase inhibition to existing anticoagulation may be useful in the case of refractory thrombotic syndromes. However, the finding that inhibition or deletion of thiol isomerases such as PDI is not associated with bleeding is not universally observed and requires additional testing in more reliable hemostasis models. In general, more basic understanding and clinical experience with thiol isomerase inhibition will be required to establish the optimal use of agents targeting these enzymes.

Summary

Vascular thiol isomerases serve an essential role in thrombus formation and there is growing evidence that they have additional functions in regulating the vascular redox environment. Nonetheless, investigators trying to unravel the mechanisms of how vascular thiol isomerases function in thrombus formation face several challenges. While investigators have demonstrated the role of PDI, ERp5, and ERp57 in thrombus formation in several thrombosis models, the full repertoire of thiol isomerases that participate in thrombosis remains to be identified and how extracellular thiol isomerases interact with one another in the context of thrombosis is currently unknown. Understanding how and under what conditions thiol isomerases are released is central to understanding their function and to developing the possibility of using plasma thiol isomerase levels diagnostically. Identifying substrates of thiol isomerases will be critical to understanding activation pathways that rely on disulfide bond rearrangements (Figure 7). How thiol isomerases are regulated within the vasculature remains to be determined. As these facets of vascular thiol

isomerase biology are elucidated, investigators need to form a compelling general model of how thiol isomerases prime the vasculature for thrombus formation. Even as mechanistic studies are underway, clinical studies evaluating how to best use thiol isomerase-targeted therapies are taking place and will help pave the way for new approaches to disorders of hemostasis and thrombosis.

Acknowledgments

This work was supported by National Institutes of Health, National Heart, Lung, and Blood Institute grants U54 HL112302, HL125275,

HL112809, and T32 HL007917, and National Institute on Drug Abuse grant DA032476.

Authorship

Contribution: R.F. and B.F. wrote the paper.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Robert Flaumenhaft, Division of Hemostasis and Thrombosis, Department of Medicine, Beth Israel Deaconess Medical Center, 330 Brookline Ave, Boston, MA 02215; e-mail: rflaumen@bidmc.harvard.edu.

References

- Sela M, White FH Jr, Anfinsen CB. Reductive cleavage of disulfide bridges in ribonuclease. *Science*. 1957;125(3250):691-692.
- Dani HM, Fielder JA, Rabin BR. Degranulation and sex-specific regranulation of reticular membranes from rat liver as studied using a spectrophotometric assay of protein disulphide isomerase. *FEBS Lett*. 1976;65(3):377-382.
- Goldberger RF, Epstein CJ, Anfinsen CB. Acceleration of reactivation of reduced bovine pancreatic ribonuclease by a microsomal system from rat liver. *J Biol Chem*. 1963;238:628-635.
- Bulleid NJ, Freedman RB. Defective co-translational formation of disulphide bonds in protein disulphide-isomerase-deficient microsomes. *Nature*. 1988;335(6191):649-651.
- Ellgaard L, Ruddock LW. The human protein disulphide isomerase family: substrate interactions and functional properties. *EMBO Rep*. 2005;6(1):28-32.
- Qin J, Clore GM, Gronenborn AM. The high-resolution three-dimensional solution structures of the oxidized and reduced states of human thioredoxin. *Structure*. 1994;2(6):503-522.
- Weichsel A, Gasdaska JR, Powis G, Montfort WR. Crystal structures of reduced, oxidized, and mutated human thioredoxins: evidence for a regulatory homodimer. *Structure*. 1996;4(6):735-751.
- McArthur AG, Knodler LA, Silberman JD, Davids BJ, Gillin FD, Sogin ML. The evolutionary origins of eukaryotic protein disulfide isomerase domains: new evidence from the Amitochondriate protist *Giardia lamblia*. *Mol Biol Evol*. 2001;18(8):1455-1463.
- Zhou J, Wu Y, Wang L, et al. The C-terminal CGHC motif of protein disulfide isomerase supports thrombosis. *J Clin Invest*. 2015;125(12):4391-4406.
- Tian G, Xiang S, Noiva R, Lennarz WJ, Schindelin H. The crystal structure of yeast protein disulfide isomerase suggests cooperativity between its active sites. *Cell*. 2006;124(1):61-73.
- Song JL, Wang CC. Chaperone-like activity of protein disulfide-isomerase in the refolding of rhodanese. *Eur J Biochem*. 1995;231(2):312-316.
- Yao Y, Zhou Y, Wang C. Both the isomerase and chaperone activities of protein disulfide isomerase are required for the reactivation of reduced and denatured acidic phospholipase A2. *EMBO J*. 1997;16(3):651-658.
- Zai A, Rudd MA, Scribner AW, Loscalzo J. Cell-surface protein disulfide isomerase catalyzes transnitrosation and regulates intracellular transfer of nitric oxide. *J Clin Invest*. 1999;103(3):393-399.
- Sliskovic I, Raturi A, Mutus B. Characterization of the S-denitrosation activity of protein disulfide isomerase. *J Biol Chem*. 2005;280(10):8733-8741.
- Ramachandran N, Root P, Jiang XM, Hogg PJ, Mutus B. Mechanism of transfer of NO from extracellular S-nitrosothiols into the cytosol by cell-surface protein disulfide isomerase. *Proc Natl Acad Sci USA*. 2001;98(17):9539-9544.
- Uys JD, Xiong Y, Townsend DM. Nitrosative stress-induced S-glutathionylation of protein disulfide isomerase. *Methods Enzymol*. 2011;490:321-332.
- Townsend DM, Manevich Y, He L, et al. Nitrosative stress-induced S-glutathionylation of protein disulfide isomerase leads to activation of the unfolded protein response. *Cancer Res*. 2009;69(19):7626-7634.
- Murphy DD, Reddy EC, Moran N, O'Neill S. Regulation of platelet activity in a changing redox environment. *Antioxid Redox Signal*. 2014;20(13):2074-2089.
- Go Y-M, Jones DP. Intracellular proatherogenic events and cell adhesion modulated by extracellular thiol/disulfide redox state. *Circulation*. 2005;111(22):2973-2980.
- Lundström J, Holmgren A. Determination of the reduction-oxidation potential of the thioredoxin-like domains of protein disulfide-isomerase from the equilibrium with glutathione and thioredoxin. *Biochemistry*. 1993;32(26):6649-6655.
- Swiatkowska M, Padula G, Michalec L, Stasiak M, Skurzynski S, Cierniewski CS. Ero1alpha is expressed on blood platelets in association with protein-disulfide isomerase and contributes to redox-controlled remodeling of alphaIIb beta3. *J Biol Chem*. 2010;285(39):29874-29883.
- Laurindo FRM, Pescatore LA, Fernandes DC. Protein disulfide isomerase in redox cell signaling and homeostasis. *Free Radic Biol Med*. 2012;52(9):1954-1969.
- Wang C, Li W, Ren J, et al. Structural insights into the redox-regulated dynamic conformations of human protein disulfide isomerase. *Antioxid Redox Signal*. 2013;19(1):36-45.
- Wang C, Yu J, Huo L, Wang L, Feng W, Wang CC. Human protein-disulfide isomerase is a redox-regulated chaperone activated by oxidation of domain a'. *J Biol Chem*. 2012;287(2):1139-1149.
- Bekendam RH, Bendapudi P, Lin L, et al. A substrate-driven allosteric switch that enhances PDI catalytic activity. *Nat Commun*. In press.
- Burgess JK, Hotchkiss KA, Suter C, et al. Physical proximity and functional association of glycoprotein 1balpha and protein-disulfide isomerase on the platelet plasma membrane. *J Biol Chem*. 2000;275(13):9758-9766.
- Summers FA, Forsman Quigley A, Hawkins CL. Identification of proteins susceptible to thiol oxidation in endothelial cells exposed to hypochlorous acid and N-chloramines. *Biochem Biophys Res Commun*. 2012;425(2):157-161.
- Fernando MR, Nanri H, Yoshitake S, Nagata-Kuno K, Minakami S. Thioredoxin regenerates proteins inactivated by oxidative stress in endothelial cells. *Eur J Biochem*. 1992;209(3):917-922.
- Ejima K, Nanri H, Araki M, Uchida K, Kashimura M, Ikeda M. 17beta-estradiol induces protein thiol/disulfide oxidoreductases and protects cultured bovine aortic endothelial cells from oxidative stress. *Eur J Endocrinol*. 1999;140(6):608-613.
- Ioannou Y, Zhang J-Y, Passam FH, et al. Naturally occurring free thiols within beta 2-glycoprotein 1 in vivo: nitrosylation, redox modification by endothelial cells, and regulation of oxidative stress-induced cell injury. *Blood*. 2010;116(11):1961-1970.
- Shah CM, Bell SE, Locke IC, Chowdrey HS, Gorge MP. Interactions between cell surface protein disulphide isomerase and S-nitrosoglutathione during nitric oxide delivery. *Nitric Oxide*. 2007;16(1):135-142.
- Irwin C, Roberts W, Naseem KM. Nitric oxide inhibits platelet adhesion to collagen through cGMP-dependent and independent mechanisms: the potential role for S-nitrosylation. *Platelets*. 2009;20(7):478-486.
- Uehara T, Nakamura T, Yao D, et al. S-nitrosylated protein-disulphide isomerase links protein misfolding to neurodegeneration. *Nature*. 2006;441(7092):513-517.
- Walsh GM, Leane D, Moran N, et al. S-Nitrosylation of platelet alphaIIb beta3 as revealed by Raman spectroscopy. *Biochemistry*. 2007;46(21):6429-6436.
- Bell SE, Shah CM, Gorge MP. Protein disulfide-isomerase mediates delivery of nitric oxide redox derivatives into platelets. *Biochem J*. 2007;403(2):283-288.
- Holbrook LM, Watkins NA, Simmonds AD, Jones CI, Ouweland WH, Gibbins JM. Platelets release novel thiol isomerase enzymes which are recruited to the cell surface following activation. *Br J Haematol*. 2010;148(4):627-637.
- Galligan JJ, Petersen DR. The human protein disulfide isomerase gene family. *Hum Genomics*. 2012;6:6.
- Chen K, Lin Y, Detwiler TC. Protein disulfide isomerase activity is released by activated platelets. *Blood*. 1992;79(9):2226-2228.
- Essex DW, Chen K, Swiatkowska M. Localization of protein disulfide isomerase to the external surface of the platelet plasma membrane. *Blood*. 1995;86(6):2168-2173.
- Chen K, Detwiler TC, Essex DW. Characterization of protein disulfide isomerase released from activated platelets. *Br J Haematol*. 1995;90(2):425-431.
- Essex DW, Li M. Protein disulphide isomerase mediates platelet aggregation and secretion. *Br J Haematol*. 1999;104(3):448-454.
- Crescente M, Pluthero FG, Li L, et al. Intracellular Trafficking, Localization, and Mobilization of

- Platelet-Borne Thiol Isomerases. *Arterioscler Thromb Vasc Biol.* 2016;36(6):1164-1173.
43. Thon JN, Peters CG, Machlus KR, et al. T granules in human platelets function in TLR9 organization and signaling. *J Cell Biol.* 2012;198(4):561-574.
 44. Sharda A, Kim SH, Jasuja R, et al. Defective PDI release from platelets and endothelial cells impairs thrombus formation in Hermansky-Pudlak syndrome. *Blood.* 2015;125(10):1633-1642.
 45. Passam FH, Lin L, Gopal S, et al. Both platelet- and endothelial cell-derived ERp5 support thrombus formation in a laser-induced mouse model of thrombosis. *Blood.* 2015;125(14):2276-2285.
 46. Kim K, Hahm E, Li J, et al. Platelet protein disulfide isomerase is required for thrombus formation but not for hemostasis in mice. *Blood.* 2013;122(6):1052-1061.
 47. Mor-Cohen R. Disulfide Bonds as Regulators of Integrin Function in Thrombosis and Hemostasis. *Antioxid Redox Signal.* 2016;24(1):16-31.
 48. Cho J, Kennedy DR, Lin L, et al. Protein disulfide isomerase capture during thrombus formation in vivo depends on the presence of $\beta 3$ integrins. *Blood.* 2012;120(3):647-655.
 49. Essex DW, Li M, Miller A, Feinman RD. Protein disulfide isomerase and sulfhydryl-dependent pathways in platelet activation. *Biochemistry.* 2001;40(20):6070-6075.
 50. Lahav J, Wijnen EM, Hess O, et al. Enzymatically catalyzed disulfide exchange is required for platelet adhesion to collagen via integrin $\alpha \text{IIb}\beta 3$. *Blood.* 2003;102(6):2085-2092.
 51. Hotchkiss KA, Chesterman CN, Hogg PJ. Catalysis of disulfide isomerization in thrombospondin 1 by protein disulfide isomerase. *Biochemistry.* 1996;35(30):9761-9767.
 52. Huang EM, Detwiler TC, Milev Y, Essex DW. Thiol-disulfide isomerization in thrombospondin: effects of conformation and protein disulfide isomerase. *Blood.* 1997;89(9):3205-3212.
 53. Hotchkiss KA, Matthias LJ, Hogg PJ. Exposure of the cryptic Arg-Gly-Asp sequence in thrombospondin-1 by protein disulfide isomerase. *Biochim Biophys Acta.* 1998;1388(2):478-488.
 54. Essex DW, Miller A, Swiatkowska M, Feinman RD. Protein disulfide isomerase catalyzes the formation of disulfide-linked complexes of vitronectin with thrombin-antithrombin. *Biochemistry.* 1999;38(32):10398-10405.
 55. Jordan PA, Stevens JM, Hubbard GP, et al. A role for the thiol isomerase protein ERP5 in platelet function. *Blood.* 2005;105(4):1500-1507.
 56. Holbrook LM, Sasikumar P, Stanley RG, Simmonds AD, Bicknell AB, Gibbins JM. The platelet-surface thiol isomerase enzyme ERp57 modulates platelet function. *J Thromb Haemost.* 2012;10(2):278-288.
 57. Wu Y, Ahmad SS, Zhou J, Wang L, Cully MP, Essex DW. The disulfide isomerase ERp57 mediates platelet aggregation, hemostasis, and thrombosis. *Blood.* 2012;119(7):1737-1746.
 58. Donoghue N, Yam PT, Jiang XM, Hogg PJ. Presence of closely spaced protein thiols on the surface of mammalian cells. *Protein Sci.* 2000;9(12):2436-2445.
 59. Swiatkowska M, Szymański J, Padula G, Cierniewski CS. Interaction and functional association of protein disulfide isomerase with $\alpha \text{V}\beta 3$ integrin on endothelial cells. *FEBS J.* 2008;275(8):1813-1823.
 60. Wan S-W, Lin C-F, Lu Y-T, Lei HY, Anderson R, Lin YS. Endothelial cell surface expression of protein disulfide isomerase activates $\beta 1$ and $\beta 3$ integrins and facilitates dengue virus infection. *J Cell Biochem.* 2012;113(5):1681-1691.
 61. Popescu NI, Lupu C, Lupu F. Extracellular protein disulfide isomerase regulates coagulation on endothelial cells through modulation of phosphatidylserine exposure. *Blood.* 2010;116(6):993-1001.
 62. Zhang LM, St Croix C, Cao R, et al. Cell-surface protein disulfide isomerase is required for transnitrosation of metallothionein by S-nitroso-albumin in intact rat pulmonary vascular endothelial cells. *Exp Biol Med (Maywood).* 2006;231(9):1507-1515.
 63. Jasuja R, Furie B, Furie BC. Endothelium-derived but not platelet-derived protein disulfide isomerase is required for thrombus formation in vivo. *Blood.* 2010;116(22):4665-4674.
 64. Falati S, Gross P, Merrill-Skoloff G, Furie BC, Furie B. Real-time in vivo imaging of platelets, tissue factor and fibrin during arterial thrombus formation in the mouse. *Nat Med.* 2002;8(10):1175-1181.
 65. Chen VM, Ahamed J, Versteeg HH, Berndt MC, Ruf W, Hogg PJ. Evidence for activation of tissue factor by an allosteric disulfide bond. *Biochemistry.* 2006;45(39):12020-12028.
 66. Ahamed J, Versteeg HH, Kerver M, et al. Disulfide isomerization switches tissue factor from coagulation to cell signaling. *Proc Natl Acad Sci USA.* 2006;103(38):13932-13937.
 67. Cho J, Furie BC, Coughlin SR, Furie B. A critical role for extracellular protein disulfide isomerase during thrombus formation in mice. *J Clin Invest.* 2008;118(3):1123-1131.
 68. Reinhardt C, von Brühl ML, Manukyan D, et al. Protein disulfide isomerase acts as an injury response signal that enhances fibrin generation via tissue factor activation. *J Clin Invest.* 2008;118(3):1110-1122.
 69. Jasuja R, Passam FH, Kennedy DR, et al. Protein disulfide isomerase inhibitors constitute a new class of antithrombotic agents. *J Clin Invest.* 2012;122(6):2104-2113.
 70. Schulman S, Bendapudi P, Sharda A, et al. Extracellular Thiol Isomerases and Their Role in Thrombus Formation. *Antioxid Redox Signal.* 2016;24(1):1-15.
 71. Wang L, Wu Y, Zhou J, et al. Platelet-derived ERp57 mediates platelet incorporation into a growing thrombus by regulation of the $\alpha \text{IIb}\beta 3$ integrin. *Blood.* 2013;122(22):3642-3650.
 72. Zhu S, Welsh JD, Brass LF, Diamond SL. Platelet-targeting thiol reduction sensor detects thiol isomerase activity on activated platelets in mouse and human blood under flow. *J Thromb Haemost.* 2016;14(5):1070-1081.
 73. Chen VM, Hogg PJ. Allosteric disulfide bonds in thrombosis and thrombolysis. *J Thromb Haemost.* 2006;4(12):2533-2541.
 74. Butera D, Cook KM, Chiu J, Wong JWH, Hogg PJ. Control of blood proteins by functional disulfide bonds. *Blood.* 2014;123(13):2000-2007.
 75. Leader A, Mor-Cohen R, Ram R, et al. The role of protein disulfide isomerase in the post-ligation phase of $\beta 3$ integrin-dependent cell adhesion. *Thromb Res.* 2015;136(6):1259-1265.
 76. Hahm E, Li J, Kim K, et al. Extracellular protein disulfide isomerase regulates ligand-binding activity of $\alpha \text{M}\beta 2$ integrin and neutrophil recruitment during vascular inflammation. *Blood.* 2013;121(19):3789-3800.
 77. Langer F, Ruf W. Synergies of phosphatidylserine and protein disulfide isomerase in tissue factor activation. *Thromb Haemost.* 2014;111(4):590-597.
 78. Popescu NI, Lupu C, Lupu F. Role of PDI in regulating tissue factor: FVIIa activity. *Thromb Res.* 2010;125(Suppl 1):S38-S41.
 79. Primm TP, Gilbert HF. Hormone binding by protein disulfide isomerase, a high capacity hormone reservoir of the endoplasmic reticulum. *J Biol Chem.* 2001;276(1):281-286.
 80. Lin L, Gopal S, Sharda A, et al. Quercetin-3-rutinoside Inhibits Protein Disulfide Isomerase by Binding to Its b'x Domain. *J Biol Chem.* 2015;290(39):23543-23552.
 81. Cui G, Shan L, Guo L, et al. Novel anti-thrombotic agent for modulation of protein disulfide isomerase family member ERp57 for prophylactic therapy. *Sci Rep.* 2015;5:10353.
 82. McCullough ML, Peterson JJ, Patel R, Jacques PF, Shah R, Dwyer JT. Flavonoid intake and cardiovascular disease mortality in a prospective cohort of US adults. *Am J Clin Nutr.* 2012;95(2):454-464.
 83. Geleijnse JM, Launer LJ, Van der Kuip DA, Hofman A, Witteman JC. Inverse association of tea and flavonoid intakes with incident myocardial infarction: the Rotterdam Study. *Am J Clin Nutr.* 2002;75(5):880-886.
 84. Flaumenhaft R, Furie B, Zwicker JJ. Therapeutic implications of protein disulfide isomerase inhibition in thrombotic disease. *Arterioscler Thromb Vasc Biol.* 2015;35(1):16-23.
 85. Manickam N, Sun X, Hakala KW, Weintraub ST, Essex DW. Thiols in the $\alpha \text{IIb}\beta 3$ integrin are necessary for platelet aggregation. *Br J Haematol.* 2008;142(3):457-465.
 86. Jurk K, Lahav J, VAN Aken H, Brodde MF, Nofer JR, Kehrel BE. Extracellular protein disulfide isomerase regulates feedback activation of platelet thrombin generation via modulation of coagulation factor binding. *J Thromb Haemost.* 2011;9(11):2278-2290.
 87. Huizing M, Boissy RE, Gahl WA. Hermansky-Pudlak syndrome: vesicle formation from yeast to man. *Pigment Cell Res.* 2002;15(6):405-419.
 88. Nagayama S, Yokoi T, Tanaka H, Kawaguchi Y, Shirasaka T, Kamataki T. Occurrence of autoantibody to protein disulfide isomerase in patients with hepatic disorder. *J Toxicol Sci.* 1994;19(3):163-169.
 89. Aran A, Weiner K, Lin L, et al. Post-streptococcal auto-antibodies inhibit protein disulfide isomerase and are associated with insulin resistance. *PLoS One.* 2010;5(9):e12875.
 90. Caorsi C, Niccolai E, Capello M, et al. Protein disulfide isomerase A3-specific Th1 effector cells infiltrate colon cancer tissue of patients with circulating anti-protein disulfide isomerase A3 autoantibodies. *Transl Res.* 2016;171:17-28.e2.
 91. Shen J, Wang W, Wu J, et al. Comparative proteomic profiling of human bile reveals SSP411 as a novel biomarker of cholangiocarcinoma. *PLoS One.* 2012;7(10):e47476.
 92. Komurasaki R, Imaoka S, Tada N, Okada K, Nishiguchi S, Funae Y. LKM-1 sera from autoimmune hepatitis patients that recognize ERp57, carboxylesterase 1 and CYP2D6. *Drug Metab Pharmacokinet.* 2010;25(1):84-92.
 93. Versteeg HH, Ruf W. Tissue factor coagulant function is enhanced by protein-disulfide isomerase independent of oxidoreductase activity. *J Biol Chem.* 2007;282(35):25416-25424.
 94. Brophy TM, Coller BS, Ahamed J. Identification of the thiol isomerase-binding peptide, mastoparan, as a novel inhibitor of shear-induced transforming growth factor $\beta 1$ (TGF- $\beta 1$) activation. *J Biol Chem.* 2013;288(15):10628-10639.