



OPEN ACCESS

EDITED BY

Chiranjeevi Pasala,
Sloan Kettering Institute, United States

REVIEWED BY

Shoyab Ansari,
University of Texas Southwestern Medical
Center, United States

*CORRESPONDENCE

Rita Grandori,
✉ rita.grandori@unimib.it

RECEIVED 16 July 2026
REVISED 08 August 2026
ACCEPTED 11 August 2026
PUBLISHED 03 September 2026

CITATION

Santambrogio C, Ponzini E, Sobott F and
Grandori R (2026) Recent advances in
mass spectrometry methods for the
investigation of oligomeric intermediates
of amyloid protein aggregation.
Front. Biophys. 4:1939530.
doi: 10.3389/frbis.2026.1939530

COPYRIGHT

© 2026 Santambrogio, Ponzini, Sobott
and Grandori. This is an open-access
article distributed under the terms of the
[Creative Commons Attribution License
\(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or
reproduction in other forums is permitted,
provided the original author(s) and the
copyright owner(s) are credited and that
the original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution or
reproduction is permitted which does not
comply with these terms.

Recent advances in mass spectrometry methods for the investigation of oligomeric intermediates of amyloid protein aggregation

Carlo Santambrogio¹, Erika Ponzini^{2,3}, Frank Sobott⁴ and
Rita Grandori^{1,5*}

¹Department of Biotechnology and Biosciences, University of Milano-Bicocca, Milan, Milano, Italy,

²Department of Materials Science, University of Milano-Bicocca, Milan, Milano, Italy, ³Optics and
Optometry Research Center (COMiB), University of Milano-Bicocca, Milan, Milano, Italy, ⁴Asbury Centre
for Structural Molecular Biology, School of Molecular and Cellular Biology, Faculty of Biological Sciences,
University of Leeds, Leeds, United Kingdom, ⁵Forschungszentrum Jülich, Jülich, Germany

Amyloid proteins are the etiological factors of many untreatable diseases. While structural characterization of amyloid fibrils has progressed fast, knowledge about oligomeric intermediates has lagged behind, due to their transient and dynamic nature. Mass spectrometry methods for structural biology are rapidly developing. The most relevant applications of the last 5 years are summarized and discussed here, focusing on the specific potential of the different approaches capturing oligomeric states of amyloid proteins.

KEYWORDS

amyloid aggregation, amyloid β peptide, neurodegenerative diseases, oligomer structure, protein dynamics, structural mass spectrometry, transthyretin, α -synuclein

1 Introduction

Protein misfolding and aggregation into amyloid structures is a competing pathway to the normal folding process of virtually any protein. This process leads to the accumulation of fibrils where monomers are stacked into β -rich structures (Sawaya et al., 2021) with variable degrees of structural disorder (Utami et al., 2025; Zhang et al., 2026). Amyloid structures can be physiologically relevant (Foust et al., 2026), but also correlate with degenerative pathologies, like Alzheimer's disease (AD) and Parkinson's disease (PD) (Siebner and Uversky, 2026). The aggregation process involves oligomeric structures, which can form by primary or secondary nucleation steps and represent either on-pathway or off-pathway intermediates (Wei et al., 2025). The structure of oligomeric intermediates remains elusive, due to their dynamic and heterogeneous nature. Nonetheless, their characterization could shed light on the aggregation mechanism and reveal potential targets of therapeutic strategies (Goncerz et al., 2026; Wei et al., 2025). Besides being recognized as toxic species (Gasca-Martínez et al., 2026; Ivanova et al., 2026a), they could carry structural determinants of "strains" involving the same protein in different pathologies (Lo Cascio et al., 2025; Saleem et al., 2026) and can also act as seeds in the so-called "prion-like" spreading, in which normally folded molecules of the aggregating protein are recruited in the pathogenic mechanism (Zampar et al., 2024). Thus, oligomeric states can reveal different aggregation pathways responsible of the diverse pathologies involving a given amyloid protein (Mahul-Mellier et al., 2026).




FIGURE 1
Schematic representation of the amyloid aggregation process and the structural-MS methods discussed in the present work.

Mass spectrometry (MS) has become a central tool of structural biology (Britt and Robinson, 2025; Kaltashov et al., 2024). Its specific strength is dissecting heterogeneous samples into sub-populations, thanks to the ion-sorting mechanism inherent to MS analysis, thus delivering measurables for each detected ion (Reid et al., 2023). A whole class of methods is available for protein structural investigation and designated as *structural MS* (Wagner and Gross, 2024). One approach is the so-called *native MS*, which analyses intact proteins and supramolecular complexes preserving the non-covalent interactions responsible for folding and binding (Lawrence et al., 2025). The charge-state distribution (CSD) obtained by electrospray ionization (ESI) reports on protein compactness, while mass reports on binding and modifications. The shape of gas-phase ions can be explored by ion mobility (IM), which measures the rotationally averaged collisional cross section (CCS) (Panczyk et al., 2026). Another class of methods is based on conformation-sensitive, covalent modification of proteins in solution (footprinting), followed by bottom-up or top-down MS analysis of the reaction products (Castel et al., 2023). Different modification strategies have been used, including cross-linking (XL) (Balakrishnan et al., 2026a), hydrogen-deuterium exchange (HDX) (Nyvall et al., 2026), fast-photochemical oxidation (FPOP) (Weckslar et al., 2026), covalent protein painting (CPP) (Bamberger et al., 2026) and limited proteolysis (LIP) (Figure 1) (Elsässer et al., 2026).

Such an arsenal of methods help dissect dynamic structural ensembles, either by equilibrium or time-resolved measurements (Lento and Wilson, 2022; Wagner and Gross, 2024), overcoming ensemble averaging. Furthermore, these methods can be combined with conventional MS protocols to characterize post-translational modifications and mutations on each dissected structural species. Structural MS, however, does not achieve atomic resolution and suffers of possible structural rearrangements and limitations on buffer composition and protein concentration. Nonetheless, the applicability of these methods is rapidly evolving. They are typically combined with thioflavin-T fluorescence for aggregation kinetics, small-angle x-ray scattering (SAXS) or nuclear magnetic resonance (NMR) for size distributions, NMR and computational simulations for modelling at atomic resolution (Happonen and Varjosalo, 2025). Native-MS methods also complement imaging by atomic-force microscopy (AFM), transmission and cryogenic electron microscopy (cryo-EM) and tomography (cryo-ET) (Hale and Cooper, 2025; Liang et al., 2025). This work highlights the specific contributions of structural MS dissecting heterogeneous and dynamic systems into structurally annotated sub-populations and yielding constraints for computational modelling (Cehlar et al., 2024; Ivanova et al., 2026b). The focus is kept on the oligomeric intermediates of amyloid proteins in the last 5 years (Wagner and Gross, 2024).

2 Amyloid β

A hallmark of AD is the aggregation of amyloid β (A β) peptides, derived from the cleavage of the amyloid precursor protein (APP) by β - and γ -secretases (Zetterberg and Bendlin, 2026). A β 40 and A β 42 are the most studied variants, but A β 43 is also involved in the pathology and all these fragments accumulate in the AD brain (Llorente-Saguer et al., 2026; Tambini et al., 2023). A β 42 and A β 43 have a stronger propensity to aggregate and higher cytotoxicity than A β 40 (Habeck et al., 2024). Native IM-MS has been used to compare homo- and hetero-oligomers (Habeck et al., 2024). This study shows that A β 40 dimers and trimers have bimodal CCS distributions, revealing a considerable amount of extended oligomers coexisting with a more compact component. A β 42 and A β 43, instead, lack the extended populations of dimers and trimers. Interestingly, the hybrid dimers and trimers of A β 40 with either A β 42 or A β 43 have similar CCS profiles as pure A β 42 and A β 43, where the extended population is depleted. Thus, A β 42 and A β 43 are capable of remodeling the oligomeric structure toward higher compaction. Considering their stimulating effect on A β 40 aggregation kinetics, it is suggested that the more compact oligomers have higher aggregation propensity, offering a mechanistic interpretation of the reciprocal influence of the A β variants (Habeck et al., 2024). Native IM-MS has also detected the hexameric *cylindrin* state of the A β 25–35 peptide, which represents the putative toxic form capable of inserting into membranes (Laos et al., 2021).

Further improvements are offered by cyclic IM (cIM) technology, which offers enhanced sensitivity and resolution through multiple ion cycling, amplifying differences in arrival times (Naylor and Nagy, 2025). An accurate methodological work on A β 42 oligomers (Vlk et al., 2026) sets the background for applications of cIM to amyloid oligomers, analyzing the effects of instrumental parameters. It is shown that upon optimization of the experimental settings, A β 42 oligomers up to the dodecamer can be detected, minimizing dissociation and denaturation. Gas-phase unfolding by collision-induced unfolding (CIU) highlights higher stability of the trimer relative to the dimer and distinct unfolding pathways (Vlk et al., 2026).

cIM has also been applied to the investigation of synthetic seeds consisting of covalent homo-trimers obtained by crosslinking the A β fragments corresponding to residues 17–36 or 16–36 (Guaglianone et al., 2022; Kreutzer et al., 2023). This portion spans the central and C-terminal regions, which are the most involved in aggregation (Chen et al., 2026). Each trimer is constrained in a β -hairpin conformation, as observed in the outer layers of mature fibrils (Ghosh et al., 2021), by an additional covalent bond, stapling together the N- and C-termini. These triangular seeds can associate into higher-order structures, offering a potentially useful tool to mimic early assemblies of A β 42 (Guaglianone et al., 2022; Kreutzer et al., 2023).

Alkylation of carboxylic residues has been exploited by glycine ethyl ester (GEE) footprinting to investigate the early stages of A β 42 oligomerization (Kuang et al., 2025). The results are consistent with involvement of the central and C-terminal regions in the aggregation process, while the N-terminus remains mostly exposed. Kinetic modelling identifies four populations that sequentially act as seeds of the following aggregation step: small

oligomers assemble first (M1) and convert into secondary seeds (M2) by an irreversible structural rearrangement. M2 seeds grow into protofibrils (M3) and eventually mature fibrils (M4) (Kuang et al., 2025).

3 α -Synuclein

The aggregation of the presynaptic protein α -synuclein (α -syn) plays a central role in the etiology of PD and other synucleopathies (Arino et al., 2026). The wild-type protein is involved in sporadic forms of the disease, while mutants are implicated with familial PD (Ranjan and Kumar, 2026). Three distinct protein regions have been recognized: an N-terminal region capable of forming α -helices, interacting with membranes and hosting most of the pathogenic mutations, a central region that drives aggregation (non-amyloid component, NAC, region) and an acidic C-terminal region mostly involved in interactions with other proteins (Singh et al., 2025). Native-MS spectra of wt α -syn without preincubation generally reveal predominance of the monomeric form and very small amounts of oligomers (Wagner and Gross, 2024). It has been shown, however, that an N-terminally truncated form accumulating upon expression in *Escherichia coli*, shows predominant signals of pentamers, hexamers, and heptamers, suggesting a faster aggregation kinetics (Báez Bolívar et al., 2026; McGlinchey et al., 2021). These studies add to a growing body of evidence hinting to a role of C- and N-terminal α -syn truncation in PD pathogenesis (Mahul-Mellier et al., 2025; McGlinchey et al., 2025). This study uses Tris- and NaCl-containing solutions, normally not compatible with ESI, thanks to sub-micron emitters, which expand the range of explorable conditions (Báez Bolívar and McLuckey, 2025; Österlund et al., 2025; Susa et al., 2018).

α -Syn oligomers have also been investigated by IM-MS in response to protein oxidation by reactive oxygen species (ROS), which has been implicated in pathogenesis (de Araújo et al., 2026). Controlled oxidation was achieved by low-temperature plasma (LTP) treatment, exploiting ultrafast protein modification (microsecond to millisecond timescale) (Wang et al., 2025a). Bottom-up LC-MS/MS analysis following the modification shows that oxidation upon 2–8-min incubation modifies preferentially the C-terminal region of monomeric α -syn (specifically M116, M127 and Y133). The results are consistent with this being the most disordered part of the protein at neutral pH and points to LTP treatment as an effective conformational probe, in addition to the more frequently employed FPOP (Cheng and Gross, 2025). Native IM-MS of untreated α -syn shows well-detectable oligomers including dimers, trimers and tetramers. Increasing treatment times (2–8 min) lead to progressive loss of α -syn oligomers, particularly trimers and tetramers, indicating a lower propensity to oligomerize of the modified protein (Wang et al., 2025a). Circular-dichroism analysis suggests that C-terminal α -syn oxidation hinders β -sheet formation. LTP-oxidized α -syn also shows lower propensity to form fibrils and reduced cytotoxicity while oxidation of preformed fibrils does not mitigate toxicity. This result suggests a complex relation between α -syn oxidation and pathogenic mechanisms and will have to be reconciled with the available evidence concerning the role of oxidative stress in neurodegeneration (Wang et al., 2025a).

HDX is a widely used method in structural MS (Nyvall et al., 2026). It is based on the exchange of amide hydrogen with deuterium from surrounding D₂O (or vice versa). Hydrogen bonding or solvent shielding protects from exchange, resulting in lower isotope incorporation. Since the exchange affects mass, methods and software have been developed for the analysis of reaction products by MS, besides NMR. Every residue is informative, regardless of its chemical structure. HDX-MS has been described the effects of epigallocatechin-3-gallate (EGCG) on large α -syn oligomers (~23 subunits and multiples of that) (Grønnemose et al., 2022). The results show selective protection of the N-terminus by EGCG, suggesting stabilization of a closed conformation. At the same time, EGCG induces oxidation of methionine residues in the same region, which appears highly disordered upon oxidation. Since a disordered and not modified N-terminus seems to mediate interaction with membranes and toxicity, these effects could be at the basis of EGCG protective action (Grønnemose et al., 2022).

HDX has also been used to investigate resveratrol interference with α -syn oligomerization (Illes-Toth et al., 2024). Pulsed labelling along the aggregation process shows the inhibitory role of resveratrol, particularly on peptides spanning the NAC region. Aggregation still proceeds in the presence of resveratrol, but the exchange patterns suggest remodeling towards a more open oligomeric structure (Illes-Toth et al., 2024).

Quite a unique role in structural MS is played by XL-MS. Reagents containing two reactive groups at defined distance are used to crosslink specific protein functional groups (Cupp-Sutton and Wu, 2026). Product analysis by bottom-up or top-down MS yields an upper limit to the distance between the crosslinked sites. Thus, crosslinkers act as chemical rulers to investigate protein structure, dynamics and interactions (Piersimoni et al., 2022). In this sense, XL-MS is analogous to FRET or NOE-NMR experiments, providing distance constraints that can be used for computational modelling. XL-MS has been applied to investigate α -syn oligomers bound to membrane mimics, revealing two structures (linear and circular) developing from α -syn dimers, depending on the relative orientation of the two subunits (Schwarz et al., 2023). Even low concentrations of the phase-2 drug UCB0599 lead to a rearrangement of the membrane-bound oligomers, changing crosslink pattern at the dimer interface. These results provide structural insights into the specific targeting of oligomeric α -syn by the drug (Schwarz et al., 2023). Analogously, XL-MS has been used to map the interaction between α -syn large oligomers (~30 subunits) and PSMA3, a peptide known to abrogate oligomer toxicity and to inhibit conversion to fibrils (Santos et al., 2024). Four crosslinks were detected, all located in α -syn N-terminal region (residues 1–61). HDX measurements provide consistent protection patterns and indicate that PSMA3 binding site remains accessible in the oligomeric state (Santos et al., 2024). This region could represent an interesting structural target for therapeutic strategies (Santos et al., 2024).

The aggregation behavior of the NAC region has also been investigated using synthetic peptides corresponding to α -syn residues 68–78, in the NAC region (PD1 peptide), and residues 47–56, in the N-terminal region (PD2 peptide) (Depland et al., 2026; Depraz Depland et al., 2025). These studies have been performed on recently developed hybrid instruments that combine IM-MS with

infrared multiple-photon dissociation (IRMPD) spectroscopy. Since MS does not provide direct information on protein secondary structure, the possibility to perform spectroscopic conformational analysis on the sub-populations resolved by IM-MS is extremely attractive. Despite very similar CCS values, the amide-A band shows that PD1 dimers form antiparallel β structure, while PD2 dimers contain parallel β -sheets. Dimers of the G51D pathogenic variant maintain a helical conformation similar to the monomer. The latter dimeric structures are also more compact and stable, based on resistance to ion heating by IRMPD (Depland et al., 2026; Depraz Depland et al., 2025).

4 Transthyretin

The recent technological development of direct charge detection MS (CDMS) overcomes the need for deconvolution procedures or isotopic resolution for charge assessment and mass calculation, enabling the analysis of heterogeneous and high-mass assemblies, such as higher-order protein oligomers (Kaltashov et al., 2024). IM-MS combined with CDMS has been used to study misfolding and aggregation of transthyretin (TTR) (Hampton et al., 2025). This protein is responsible for the physiological transport of thyroxine and retinol in plasma and cerebrospinal fluid. Its aggregation has been involved in senile systemic amyloidosis. TTR is natively tetrameric and aggregation is triggered by tetramer dissociation and misfolding of the monomers, which then re-associate into higher-order oligomers and eventually fibrils (Sun et al., 2025). Oligomerization has been explored as a function of pH, pre-incubation temperature and sequence variants (Hampton et al., 2025). IM-MS shows accumulation of extended monomers under acidic pH, enhanced by the pathogenic mutation L55 P. At the same time, CDMS shows that acidic pH, low temperature and the L55 P mutation induce heterogeneous and higher-order oligomers (up to 19 subunits detected). Instead, the octamer dominates in wt TTR, consistent with association of native-like tetramers, under pH and temperature conditions less favorable to aggregation (Hampton et al., 2025). The temperature-dependence of aggregation suggests a role of hydration as supported by further analyses at variable-temperature ESI and effects of D₂O. Furthermore, HDX-MS could map “cold” water molecules, whose exchange with the bulk solvent is inhibited by the interactions with the protein structure, at the dimer/dimer interface and shows that they are responsible for stabilization of the native tetramer (Lantz et al., 2025, 2024).

5 TDP-43

TAR DNA-binding protein 43 (TDP-43) is the target of emerging therapeutic strategies for amyotrophic lateral sclerosis (ALS) (Estevez-Fraga et al., 2026). Its oligomerization has been investigated by IM-MS (Depraz Depland et al., 2023; Laos et al., 2021), also in interplay with an A β peptide, revealing reciprocal influence (Laos et al., 2021). Furthermore, formation of the toxic hexameric cylindrin state of TDP-43_{307–319} could be monitored by IM-MS, testing the effects of drugs and identifying promising inhibitors (Liu X. et al., 2023). The effects of mutations and protein concentration has also been explored, identifying a dual

kinetic aggregation pathway leading to either fibrils or cylindrin and offering an interpretation of toxicity phenotypes (Jin et al., 2025).

6 Other amyloid systems

Native IM-MS has also been employed to study oligomerization of peptides (either designed or representing distinct regions of amyloid proteins), evaluating the effects of mutations and conditions on size distribution and stability (Hirschbeck et al., 2025; Jin et al., 2025; Wang et al., 2025b). Furthermore, it has been employed to explore the effect of disulfide shuffling on insulin aggregation pathway (Qin et al., 2025). Higher-order oligomers of the therapeutic peptide liraglutide have been detected by DCMS, revealing a dual aggregation pathway (Kuo et al., 2025).

XL-MS has been applied to the investigation of the early stages of aggregation of the protein $\beta 2$ microglobulin, responsible for the dialysis-related amyloidosis. An off-pathway covalent dimer has been produced by photo-induced crosslinking and shown to inhibit the aggregation of unmodified protein (Maya-Martinez et al., 2022).

HDX-MS has been applied to structural characterization of oligomeric intermediates in the aggregation pathway of insulin β -chain (Renawala and Topp, 2022), glucagon-analog peptides (Renawala et al., 2024), and tau (James et al., 2025). In the latter case, by relating the oligomer structure to fibril morphology, it was possible to show that the polymorphism due to aggregation conditions is imprinted in the early stages of aggregation (James et al., 2025).

Islet amyloid polypeptide (IAPP) is a polypeptide hormone that aggregates in the islets of Langerhans, contributing to β -cell dysfunction in type-2 diabetes and showing pathophysiological connections to insulin and $A\beta$ (Bousch et al., 2026; Dai et al., 2024; Fu et al., 2026). Structural MS has long been applied to the study of IAPP oligomeric intermediates (Abedini et al., 2016; Eldrid et al., 2021; Lim et al., 2024; Ridgway et al., 2020; Xu et al., 2022; Young et al., 2015). Among the postulated toxicity mechanisms, interactions of IAPP with membranes play a key role (Bousch et al., 2026; Dai et al., 2024; Mithu et al., 2025). These interactions have been recently investigated by native MS of IAPP bound to nanodiscs (Sanders et al., 2023), membrane mimetics consisting of lipid bilayers surrounded by a scaffold protein belt (Townsend and Marty, 2023). Oligomerization of a host protein bound to nanodiscs can be assessed by mass defect analysis of the raw nMS spectra (Townsend and Marty, 2023). This approach has shown that human IAPP forms tetramers in a nanodisc environment, while the non-amyloidogenic rat variant binds sequentially and only up to a 2:1 stoichiometry. Still another behavior is shown by the non-amyloidogenic pig protein, lacking binding at all (Sanders et al., 2023). These experiments set the basis for further mechanistic studies and screening of inhibitors.

7 Future perspectives

Two powerful tools of structural MS have not been applied yet to the investigation of amyloid oligomers, limited proteolysis coupled to MS (LiP-MS) (Elsässer et al., 2026) and covalent protein painting (CPP) (Son et al., 2023). These approaches probe local solvent

accessibility and conformational dynamics, respectively, by enzymatic proteolysis and isotope-coded chemical methylation. Downstream, quantitative, bottom-up MS generates fingerprints of the original structures, with site-specific information on the degree of solvent exposure. These methods have been applied to structural investigation of amyloid fibrils, e.g., α -syn (Serdiuk et al., 2026) and $A\beta 40/A\beta 42$ (Bamberger et al., 2026), characterizing alternative polymorphs, as well as to monomeric and fibrillar α -syn, identifying mitochondrial interactors (Serdiuk et al., 2025). Future extension of these approaches to oligomeric structures is highly promising. It has been shown that sensitivity to proteolysis discriminates stability and morphology of α -syn and tau oligomers (Lo Cascio et al., 2025).

Protein amyloid aggregation inside the cell is promoted by liquid-liquid phase separation (LLPS), a process with complex physiological and pathological implications (Lucas et al., 2026; Mukherjee et al., 2024; Tang et al., 2025). XL-MS has been successfully applied to α -syn characterization inside LLPS “droplets” showing accumulation of an elongated conformer (Ubbiali et al., 2022). It would be important to also catch the oligomeric species of amyloid proteins that form under LLPS conditions by this and other MS-based techniques (Sahin et al., 2023).

A major Frontier lies in the integration of MS with artificial intelligence (AI), for instance, facilitating automated feature extraction, spectra deconvolution, and quantitative analysis (Yu et al., 2023). AI-based tools, such as AlphaFold, can be combined with structural MS, e.g., with distance constraints derived from XL-MS, to model dynamic macromolecular assemblies and non-native or self-associating amyloid architectures (Liu, 2022). Furthermore, hybrid workflows combining structural MS with cryo-EM and cryo-ET offer exceptional synergy for integrative structural characterization (Greenblatt et al., 2024; Quintana-Gallardo et al., 2022). The combination of single particle cryo-EM and structural MS can capture transient or low-abundance species that resist conventional structure determination.

The different approaches of structural MS are increasingly implemented *ex-vivo* (El-Baba et al., 2026) and XL-MS (Balakrishnan et al., 2026b) also sees applicability *in-vivo* (Balakrishnan et al., 2026a). Methods for native ambient MS (NAMS) are rapidly growing, paving the way to tissue-derived and patient-derived oligomer analysis (Du and Cooper, 2025; Hughes et al., 2024; Liu R. et al., 2023; Venter, 2025; Villacob et al., 2024; Wong et al., 2023). Future developments in this direction will give important contributions to the understanding of amyloid biology and amyloid diseases. These methods also have the potential to reach the *omic* scale (Khanppnnavar et al., 2024) and time-resolved implementation (Nkyaagye et al., 2026; Overby et al., 2026), opening exciting new perspectives in amyloid structural biology, not only by assisting drug discovery and elucidating aggregation mechanisms, but also helping discovery of novel biomarkers and understand the biological response to protein aggregation.

Author contributions

CS: Writing – original draft, Visualization. EP: Visualization, Writing – original draft. FS: Conceptualization,

Writing – original draft. RG: Conceptualization, Supervision, Writing – original draft.

Funding

The author(s) declared that financial support was not received for this work and/or its publication.

Conflict of interest

The author(s) declared that this work was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The author RG declared that they were an editorial board member of Frontiers at the time of submission. This had no impact on the peer review process and the final decision.

References

- Abbedini, A., Plesner, A., Cao, P., Ridgway, Z., Zhang, J., Tu, L.-H., et al. (2016). Time-resolved studies define the nature of toxic IAPP intermediates, providing insight for anti-amyloidosis therapeutics. *Elife* 5, e12977. doi: 10.7554/eLife.12977
- Arino, S., Fusco, G., and De Simone, A. (2026). α -Synuclein aggregation landscape from phase separation to neurotoxic intermediates. *FEBS Lett.* 600, 1955–1965. doi: 10.1002/1873-3468.70393
- Báez Bolívar, E. G., and McLuckey, S. A. (2025). Improved mass analysis of proteins and protein complexes present in common biological buffers and physiological concentrations of NaCl using solution additives delivered by theta emitters. *Anal. Chem.* 97, 20859–20868. doi: 10.1021/acs.analchem.5c03032
- Báez Bolívar, E. G., Fortin, J. S., Ademoye, T. A., and McLuckey, S. A. (2026). Differences in α -synuclein conformational states in physiologically relevant pH/Na⁺ concentrations and ammonium acetate solutions unveiled by native mass spectrometry. *Analyst* 151, 1748–1762. doi: 10.1039/d5an01336d
- Balakrishnan, S. M., Ansari, H. R., and Nedjadi, T. (2026a). Proteome-wide structural and interaction analysis using cross-linking mass spectrometry and its applications. *Biophys. J.* S0006-3495 (26), 00265–1. doi: 10.1016/j.bpj.2026.03.062
- Balakrishnan, S. M., Nedjadi, T., Alsiary, R., and Ansari, H. R. (2026b). Cross-linking mass spectrometry: workflow enhancements for mapping large-scale interactomes. *Biophys. J.* S0006-3495 (26), 00392–9. doi: 10.1016/j.bpj.2026.05.033
- Bamberger, T. C., Diedrich, J. K., Jung, T., Lane, J. B., Schaefer, J.-H., Lander, G. C., et al. (2026). A covalent protein painting assay differentiates Amyloid- β polymorphs in Alzheimer disease. *J. Proteome Res.* 25, 788–799. doi: 10.1021/acs.jproteome.5c00735
- Bousch, C., Bérubé, F., Baby, M., Ongeri, S., and Bourgault, S. (2026). Molecular mechanisms of islet amyloid polypeptide aggregation: towards chemical strategies to prevent amyloid formation and to design non-aggregating peptide therapeutics. *Int. J. Mol. Sci.* 27, 2598. doi: 10.3390/ijms27062598
- Britt, H. M., and Robinson, C. V. (2025). Traversing the drug discovery landscape using native mass spectrometry. *Curr. Opin. Struct. Biol.* 91, 102993. doi: 10.1016/j.sbi.2025.102993
- Castel, J., Delaux, S., Hernandez-Alba, O., and Cianférani, S. (2023). Recent advances in structural mass spectrometry methods in the context of biosimilarity assessment: from sequence heterogeneities to higher order structures. *J. Pharm. Biomed. Anal.* 236, 115696. doi: 10.1016/j.jpba.2023.115696
- Cehlar, O., Njemoga, S., Horvath, M., Cizmazia, E., Bednarikova, Z., and Barrera, E. E. (2024). Structures of oligomeric states of tau protein, Amyloid- β , α -Synuclein and prion protein implicated in Alzheimer's disease, Parkinson's disease and prionopathies. *Int. J. Mol. Sci.* 25, 13049. doi: 10.3390/ijms252313049
- Chen, B., Chen, T., Sun, Z., Liu, L., Jia, Y., Ji, Y., et al. (2026). The aggregation of amyloid- β : from condensation, nucleation, and conformation to targeting therapy. *Life Sci.* 396, 124397. doi: 10.1016/j.lfs.2026.124397
- Cheng, M., and Gross, M. L. (2025). Mass spectrometry-based protein footprinting for protein structure characterization. *Acc. Chem. Res.* 58, 165–176. doi: 10.1021/acs.accounts.4c00545
- Cupp-Sutton, K. A., and Wu, S. (2026). Top-down mass spectrometry-based structural and functional proteomics: a perspective. *J. Mass Spectrom.* 61, e70044. doi: 10.1002/jms.70044
- Dai, Z., Ben-Younis, A., Vlachaki, A., Raleigh, D., and Thalassinou, K. (2024). Understanding the structural dynamics of human islet amyloid polypeptide: advancements in and applications of ion-mobility mass spectrometry. *Biophys. Chem.* 312, 107285. doi: 10.1016/j.bpc.2024.107285
- de Araújo, A. B., S. Azul, F. V. C., Carneiro, Y. C., de Sousa, C. N. S., de Vasconcelos, S. M. M., Rios, F. J., et al. (2026). Neuroinflammation and oxidative stress in Parkinson's disease, Alzheimer's disease, and COVID-19: microglia-neutrophil interaction. *ACS Omega* 11, 6922–6938. doi: 10.1021/acsomega.5c10397
- Depland, A. D., Mikromanolis, S., Stroganova, I., and Rijs, A. M. (2026). Probing early α -Synuclein oligomers: insights into aggregation pathways of NACore and preNAC segments probed by trapped ion-mobility mass spectrometry and fluorescence spectroscopy. *Proteomics* 26, 49–59. doi: 10.1002/pmic.70087
- Depraz Depland, A., Stroganova, I., Wootton, C. A., and Rijs, A. M. (2023). Developments in trapped ion mobility mass spectrometry to probe the early stages of peptide aggregation. *J. Am. Soc. Mass Spectrom.* 34, 193–204. doi: 10.1021/jasms.2c00253
- Depraz Depland, A., Verhoeve, O., Daly, S., and Rijs, A. M. (2025). IR action spectroscopy of mass- and mobility-selected α -Synuclein segments: structure identification in early-stage oligomers via the amide A region. *J. Phys. Chem. A* 129, 8064–8074. doi: 10.1021/acs.jpca.5c02819
- Du, Y., and Cooper, H. J. (2025). Native ambient mass spectrometry of membrane proteins directly from bacterial colonies. *Chem. Commun. (Camb)* 61, 4168–4171. doi: 10.1039/d4cc03881a
- El-Baba, T. J., Lutowski, C. A., Bennett, J. L., Lawrence, S. A. S., Burnap, S. A., Butroid, F. I., et al. (2026). *Molecular Dissection of Protein Complexes Isolated from Sections of Human Brain*. doi: 10.64898/2026.04.11.717886
- Eldrid, C., Ben-Younis, A., Ujma, J., Britt, H., Cragnolini, T., Kalfas, S., et al. (2021). Cyclic ion mobility-collision activation experiments elucidate protein behavior in the gas phase. *J. Am. Soc. Mass Spectrom.* 32, 1545–1552. doi: 10.1021/jasms.1c00018
- Elsässer, F., de Souza, N., and Picotti, P. (2026). 3D proteomics: structural, functional, chemical and biomarker discovery proteomics with LiP-MS. *Mol. Cell Proteomics* 25, 101552. doi: 10.1016/j.mcp.2026.101552
- Estevez-Fraga, C., Alvarez-Velasco, R., Afroz, T., Costa, M. R., Jovičić, A., Yeh, F. L., et al. (2026). Targeting TDP-43 in sporadic amyotrophic lateral sclerosis. *J. Neurol.* 273, 498. doi: 10.1007/s00415-026-14023-5
- Foust, D. J., Kolli, D., Jessel, K., Hu, Z., Chapman, M. R., and Biteen, J. S. (2026). Transient binding facilitates super-resolution imaging of functional amyloid fibrils on living bacteria. *Protein Sci.* 35, e70398. doi: 10.1002/pro.70398
- Fu, W., Lu, R., Wang, D., and Sang, Z. (2026). Shared pathophysiology and therapeutic repurposing in Alzheimer's disease and type 2 diabetes: a critical review of convergent mechanisms and clinical challenges. *Bioorg. Chem.* 180, 110044. doi: 10.1016/j.bioorg.2026.110044
- Gasca-Martínez, Y., Ontiveros-Torres, M. A., López-Gallegos, I., and Jarero-Basulto, J. J. (2026). Amyloid beta oligomers as early triggers of neuronal cytoskeleton dysfunction in Alzheimer's disease. *Pathophysiology* 33, 14. doi: 10.3390/pathophysiology33010014
- Ghosh, U., Thurber, K. R., Yau, W.-M., and Tycko, R. (2021). Molecular structure of a prevalent amyloid- β fibril polymorph from Alzheimer's disease brain tissue. *Proc. Natl. Acad. Sci. U. S. A.* 118, e2023089118. doi: 10.1073/pnas.2023089118

Generative AI statement

The author(s) declared that generative AI was not used in the creation of this manuscript.

Any alternative text (alt text) provided alongside figures in this article has been generated by Frontiers with the support of artificial intelligence and reasonable efforts have been made to ensure accuracy, including review by the authors wherever possible. If you identify any issues, please contact us.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

- Goncerz, M., Rubiś, B., Szukalska, M., Totoń, E., Nehra, M., Malesza, I., et al. (2026). Nanoscale interrogation of α -Synuclein oligomers: bridging biophysics, interface science, and neurobiology. *Adv. Colloid Interface Sci.* 356, 103965. doi: 10.1016/j.cis.2026.103965
- Greenblatt, J. F., Alberts, B. M., and Krogan, N. J. (2024). Discovery and significance of protein-protein interactions in health and disease. *Cell* 187, 6501–6517. doi: 10.1016/j.cell.2024.10.038
- Grønneimose, A. L., Österlund, E. C., Otzen, D. E., and Jørgensen, T. J. D. (2022). EGCG has dual and opposing effects on the N-terminal region of self-associating α -synuclein oligomers. *J. Mol. Biol.* 434, 167855. doi: 10.1016/j.jmb.2022.167855
- Guaglianone, G., Torrado, B., Lin, Y.-F., Watkins, M. C., Wysocki, V. H., Gratton, E., et al. (2022). Elucidating the oligomerization and cellular interactions of a trimer derived from A β through fluorescence and mass spectrometric studies. *ACS Chem. Neurosci.* 13, 2473–2482. doi: 10.1021/acscchemneuro.2c00313
- Habeck, T., Zurmühl, S. S., Figueira, A. J., Maciel, E. V. S., Gomes, C. M., and Lermyte, F. (2024). Cross-interactions of A β peptides implicated in alzheimer's disease shape amyloid oligomer structures and aggregation. *ACS Chem. Neurosci.* 15, 4295–4304. doi: 10.1021/acscchemneuro.4c00492
- Hale, O. J., and Cooper, H. J. (2025). Imaging of protein assemblies up to 231 kDa in tissues with Nano-DESI mass spectrometry. *Anal. Chem.* 97, 28343–28352. doi: 10.1021/acs.analchem.5c05767
- Hampton, J., Lantz, C., Rider, R. L., Yun, S. D., Laganowsky, A., and Russell, D. H. (2025). Effects of hydration on transthyretin conformational dynamics and oligomerization. *Biochemistry* 64, 4571–4580. doi: 10.1021/acs.biochem.5c00589
- Happonen, L. J., and Varjosalo, M. (2025). State-of-the-Art and future directions in structural proteomics. *Mol. Cell Proteomics* 24, 101065. doi: 10.1016/j.mcpro.2025.101065
- Hirschbeck, S. S., Sawaya, M. R., Lindberg, E. T., Limbach, M. N., Jang, J. H., Lazar Cantrell, K. L., et al. (2025). Amyloid oligomers: expediting crystal growth and revisiting the corkscrew structures. *J. Am. Chem. Soc.* 147, 18594–18605. doi: 10.1021/jacs.5c00656
- Hughes, J. W., Sisley, E. K., Hale, O. J., and Cooper, H. J. (2024). Laser capture microdissection and native mass spectrometry for spatially-resolved analysis of intact protein assemblies in tissue. *Chem. Sci.* 15, 5723–5729. doi: 10.1039/d3sc04933g
- Illes-Toth, E., Rempel, D. L., and Gross, M. L. (2024). Exploration of resveratrol as a potent modulator of α -Synuclein fibril formation. *ACS Chem. Neurosci.* 15, 503–516. doi: 10.1021/acscchemneuro.3c00571
- Ivanova, M. I., La Rosa, C., and Ramamoorthy, A. (2026a). Toxic mechanisms of amyloid oligomers and therapeutic strategies. *Protein Sci.* 35, e70519. doi: 10.1002/pro.70519
- Ivanova, M. I., La Rosa, C., and Ramamoorthy, A. (2026b). *In vitro*, cellular and *in vivo* studies of amyloid oligomers structure and toxicity: challenges and advances. *Protein Sci.* 35, e70525. doi: 10.1002/pro.70525
- James, E. I., Saunders, M., Lee, K. K., Guttman, M., and Nath, A. (2025). *Tau4RD Fibril Polymorphism Is Imprinted During Early Aggregation*. doi: 10.1101/2025.10.12.681486
- Jin, Y., Ganguly, P., Shea, J.-E., Buratto, S. K., and Bowers, M. T. (2025). Aggregation of TDP-43307-319: a dual pathway to forming cylindrins and fibrils and a contribution to the etiology of amyotrophic lateral sclerosis. *J. Phys. Chem. B* 129, 8099–8114. doi: 10.1021/acs.jpcc.5c02640
- Kaltashov, I. A., Ivanov, D. G., and Yang, Y. (2024). Mass spectrometry-based methods to characterize highly heterogeneous biopharmaceuticals, vaccines, and nonbiological complex drugs at the intact-mass level. *Mass Spectrom. Rev.* 43, 139–165. doi: 10.1002/mas.21829
- Khanppnavar, B., Schuster, D., Lavriha, P., Uliana, F., Özel, M., Mehta, V., et al. (2024). Regulatory sites of CaM-sensitive adenylyl cyclase AC8 revealed by cryo-EM and structural proteomics. *EMBO Rep.* 25, 1513–1540. doi: 10.1038/s44319-024-00076-y
- Kreutzer, A. G., Guaglianone, G., Yoo, S., Parrocha, C. M. T., Ruttenberg, S. M., Malonis, R. J., et al. (2023). Probing differences among A β oligomers with two triangular trimers derived from A β . *Proc. Natl. Acad. Sci. U. S. A.* 120, e2219216120. doi: 10.1073/pnas.2219216120
- Kuang, X. C., Rempel, D. L., Lin, Y., and Gross, M. L. (2025). Mass-spectrometry-based GEE footprinting characterizes kinetic mechanisms and sites of conformational change in amyloid β 1-42 aggregation. *ACS Chem. Neurosci.* 16, 2048–2059. doi: 10.1021/acscchemneuro.5c00033
- Kuo, S.-T., Xi, Z., Cong, X., Yan, X., and Russell, D. H. (2025). Dissecting hidden ligand oligomerization pathways via direct mass technology, electron-capture dissociation, and molecular dynamics. *Anal. Chem.* 97, 13465–13473. doi: 10.1021/acs.analchem.5c01851
- Lantz, C., Rider, R. L., Yun, S. D., Laganowsky, A., and Russell, D. H. (2024). Water plays key roles in stabilities of wild type and mutant transthyretin complexes. *J. Am. Soc. Mass Spectrom.* 35, 1854–1864. doi: 10.1021/jasms.4c00170
- Lantz, C., Rider, R. L., Kuo, S.-T., Xi, Z., Burningham, E., Yun, S. D., et al. (2025). Hydration (H₂O and D₂O) dictates the stabilities and conformational entropy of transthyretin tetramers. *J. Am. Chem. Soc.* 147, 35616–35626. doi: 10.1021/jacs.5c11055
- Laos, V., Bishop, D., Ganguly, P., Schonfeld, G., Trapp, E., Cantrell, K. L., et al. (2021). Catalytic cross talk between key peptide fragments that couple alzheimer's disease with amyotrophic lateral sclerosis. *J. Am. Chem. Soc.* 143, 3494–3502. doi: 10.1021/jacs.0c12729
- Lawrence, S. A. S., Dolan, A., Miller, M. M., and Robinson, C. V. (2025). Membrane protein complexity revealed through native mass spectrometry. *Annu. Rev. Biochem.* 94, 111–135. doi: 10.1146/annurev-biochem-081424-044501
- Lento, C., and Wilson, D. J. (2022). Subsecond time-resolved mass spectrometry in dynamic structural biology. *Chem. Rev.* 122, 7624–7646. doi: 10.1021/acs.chemrev.1c00222
- Liang, R., Khurshed, A., Tahirbegi, B., Torres-Flores, A. P., Qi, S., Tian, Y., et al. (2025). Amyloid- β oligomers, curvilinear and annular assemblies, imaged by cryo-ET, cryo-EM, and AFM. *Sci. Adv.* 11, ead9030. doi: 10.1126/sciadv.ad9030
- Lim, J. Y., Kim, S. Y., and Kim, H.-T. (2024). Analysis of dimer and trimer complexes of the non-amyloidogenic rat islet amyloid polypeptide 21–37 by electrospray ionization-tandem mass spectrometry. *Eur. J. Mass Spectrom. (Chichester)* 30, 254–260. doi: 10.1177/14690667241278462
- Liu, H. (2022). AlphaFold and structural mass spectrometry enable interrogations on the intrinsically disordered regions in cyanobacterial light-harvesting complex phycobilisome. *J. Mol. Biol.* 434, 167831. doi: 10.1016/j.jmb.2022.167831
- Liu, R., Xia, S., and Li, H. (2023). Native top-down mass spectrometry for higher-order structural characterization of proteins and complexes. *Mass Spectrom. Rev.* 42, 1876–1926. doi: 10.1002/mas.21793
- Liu, X., Duan, S., Jin, Y., Walker, E., Tsao, M., Jang, J. H., et al. (2023). Computationally designed molecules modulate ALS-related amyloidogenic TDP-43307-319 aggregation. *ACS Chem. Neurosci.* 14, 4395–4408. doi: 10.1021/acscchemneuro.3c00582
- Llorente-Saguer, I., Gabriele, R., Bradshaw, T. Y., Leckey, C. A., Belder, C. R. S., Chávez-Gutiérrez, L., et al. (2026). Unbiased data-driven analysis of five amyloid-beta peptides for biomarker investigations in familial Alzheimer's disease. *Brain Commun.* 8, fcag105. doi: 10.1093/braincomms/fcag105
- Lo Cascio, F., Park, S., Sengupta, U., Puangmalai, N., Bhatt, N., Shchankin, N., et al. (2025). Brain-derived tau oligomer polymorphs: distinct aggregations, stability profiles, and biological activities. *Commun. Biol.* 8, 53. doi: 10.1038/s42003-025-07499-w
- Lucas, L., Ferreón, J. C., and Ferreón, A. C. M. (2026). Condensate state as determinant of amyloid pathology in neurodegeneration. *Biomolecules* 16, 560. doi: 10.3390/biom16040560
- Mahul-Mellier, A.-L., Altay, M. F., Maharjan, N., Ait-Bouziad, N., Chiki, A., Jagannath, S., et al. (2025). Differential role of C-terminal truncations on alpha-synuclein pathology and lewy body formation. *NPJ Park. Dis.* 11, 261. doi: 10.1038/s41531-025-01084-y
- Mahul-Mellier, A.-L., van den Heuvel, L., Teixeira, M., Boussouf, M. L. N., Oudinot, G., Thonet, A., et al. (2026). Modeling Parkinson's pathology in human iPSC dopaminergic neurons uncovers key mechanisms of lewy body formation and heterogeneity. *Sci. Adv.* 12, ead8851. doi: 10.1126/sciadv.aed8851
- Maya-Martinez, R., Xu, Y., Guthertz, N., Walko, M., Karamanos, T. K., Sobott, F., et al. (2022). Dimers of D76N- β 2-microglobulin display potent anti-amyloid aggregation activity. *J. Biol. Chem.* 298, 102659. doi: 10.1016/j.jbc.2022.102659
- McGlinchey, R. P., Ni, X., Shadish, J. A., Jiang, J., and Lee, J. C. (2021). The N terminus of α -synuclein dictates fibril formation. *Proc. Natl. Acad. Sci. U. S. A.* 118, e2023487118. doi: 10.1073/pnas.2023487118
- McGlinchey, R. P., Ramos, S., Dimitriadis, E. K., Wilson, C. B., and Lee, J. C. (2025). Defining essential charged residues in fibril formation of a lysosomal derived N-terminal α -synuclein truncation. *Nat. Commun.* 16, 3825. doi: 10.1038/s41467-025-58899-9
- Mithu, V. S., Giller, K., Nimerovsky, E., Overkamp, K., Andreas, L. B., Becker, S., et al. (2025). The structure of human IAPP fibrils reflects membrane and pH conditions. *J. Am. Chem. Soc.* 147, 28943–28954. doi: 10.1021/jacs.5c06971
- Mukherjee, S., Poudyal, M., Dave, K., Kadu, P., and Maji, S. K. (2024). Protein misfolding and amyloid nucleation through liquid-liquid phase separation. *Chem. Soc. Rev.* 53, 4976–5013. doi: 10.1039/d3cs01065a
- Naylor, C. N., and Nagy, G. (2025). Recent advances in high-resolution traveling wave-based ion mobility separations coupled to mass spectrometry. *Mass Spectrom. Rev.* 44, 581–598. doi: 10.1002/mas.21902
- Nkyaagye, E., Safaoh, H. A., and Do, T. D. (2026). Cyclic peptide conformational landscapes through the lens of mass spectrometry and orthogonal spectroscopy. *J. Mass Spectrom.* 61, e70058. doi: 10.1002/jms.70058
- Nyval, H. G., Shaw, A. L., Walsh, E. E., Bhatti, H., and Burke, J. E. (2026). Use of hydrogen deuterium exchange mass spectrometry in tandem with modern structural biology. *Biochem. J.* 483, 585–602. doi: 10.1042/BCJ20250131
- Österlund, N., Jordan, J. S., Renzi, E., Szekeres, G. P., and Pagel, K. (2025). Mass spectrometry detects folding intermediates populated during urea-induced protein denaturation. *Chem. Sci.* 16, 22407–22416. doi: 10.1039/d5sc05773f
- Overby, A., Godary, T., Anderson, S., Awoyemi, O., Yuliantoro, H., Valentine, S. J., et al. (2026). Integrating simple microfluidics design/fabrication with a novel ionization source for time-resolved chemical reaction studies by mass spectrometry. *ACS Omega* 11, 26512–26522. doi: 10.1021/acsomega.5c12681
- Panczyk, E. M., Martelli, C., Ridgeway, M. E., Ekeloef, M., Hornburg, D., Jones, B. J., et al. (2026). The rise of trapped ion mobility spectrometry: principles, applications, and recent developments. *Mass Spectrom. Rev.* e70034. doi: 10.1002/mas.70034

- Piersimoni, L., Kastiris, P. L., Arlt, C., and Sinz, A. (2022). Cross-linking mass spectrometry for investigating protein conformations and protein-protein interactions—a method for all seasons. *Chem. Rev.* 122, 7500–7531. doi: 10.1021/acs.chemrev.1c00786
- Qin, W., Li, R., Liu, J., Liu, J., Wang, X., Hu, B., et al. (2025). Spatially constrained disulfide bond shuffling delays insulin aggregation and enhances neurotoxicity. *Nat. Commun.* 16, 6841. doi: 10.1038/s41467-025-62257-0
- Quintana-Gallardo, L., Maestro-López, M., Martín-Benito, J., Marcilla, M., Rutz, D., Buchner, J., et al. (2022). “Combining electron microscopy (EM) and cross-linking mass spectrometry (XL-MS) for structural characterization of protein complexes,” in *Clinical Proteomics: Methods and Protocols*. Editors F. J. Corrales, A. Paradelo, and M. Marcilla (New York, NY: Springer US), 217–232. doi: 10.1007/978-1-0716-1936-0_17
- Ranjan, P., and Kumar, A. (2026). Unfolding sequence-specific enigma in monomeric α -Synuclein: implications for parkinson's disease. *ACS Chem. Neurosci.* 17, 21–37. doi: 10.1021/acscchemneuro.5c00423
- Reid, D. J., Thibert, S., and Zhou, M. (2023). Dissecting the structural heterogeneity of proteins by native mass spectrometry. *Protein Sci.* 32, e4612. doi: 10.1002/pro.4612
- Renawala, H. K., and Topp, E. M. (2022). Fibrillation of human insulin B-chain by pulsed hydrogen-deuterium exchange mass spectrometry. *Biophys. J.* 121, 4505–4516. doi: 10.1016/j.bpj.2022.10.042
- Renawala, H. K., Chandrababu, K. B., Smith, K. J., D'Addio, S. M., and Topp, E. M. (2024). A model study to assess fibrillation and product stability to support peptide drug design. *Mol. Pharm.* 21, 2223–2237. doi: 10.1021/acs.molpharmaceut.3c00996
- Ridgway, Z., Lee, K.-H., Zhyvoloup, A., Wong, A., Eldrid, C., Hannaberry, E., et al. (2020). Analysis of baboon IAPP provides insight into amyloidogenicity and cytotoxicity of human IAPP. *Biophys. J.* 118, 1142–1151. doi: 10.1016/j.bpj.2019.12.027
- Sahin, C., Leppert, A., and Landreh, M. (2023). Advances in mass spectrometry to unravel the structure and function of protein condensates. *Nat. Protoc.* 18, 3653–3661. doi: 10.1038/s41596-023-00900-0
- Saleem, T., Möbius, W., Schmitz, M., da Silva Correia, A., Thomas, C., Canaslan, S., et al. (2026). A distinct tau oligomer strain defines the molecular and proteomic landscape of rapidly progressive Alzheimer's disease. *Acta Neuropathol.* 151, 27. doi: 10.1007/s00401-026-02998-4
- Sanders, H. M., Chalyavi, F., Fields, C. R., Kostelic, M. M., Li, M.-H., Raleigh, D. P., et al. (2023). Interspecies variation affects islet amyloid polypeptide membrane binding. *J. Am. Soc. Mass Spectrom.* 34, 986–990. doi: 10.1021/jasms.3c00005
- Santos, J., Cuellar, J., Pallarès, I., Byrd, E. J., Lends, A., Moro, F., et al. (2024). A targetable N-Terminal motif orchestrates α -Synuclein oligomer-to-fibril conversion. *J. Am. Chem. Soc.* 146, 12702–12711. doi: 10.1021/jacs.4c02262
- Sawaya, M. R., Hughes, M. P., Rodriguez, J. A., Riek, R., and Eisenberg, D. S. (2021). The expanding amyloid family: structure, stability, function, and pathogenesis. *Cell* 184, 4857–4873. doi: 10.1016/j.cell.2021.08.013
- Schwarz, T. C., Beier, A., Ledolter, K., Gossenreiter, T., Höfurther, T., Hartl, M., et al. (2023). High-resolution structural information of membrane-bound α -synuclein provides insight into the MoA of the anti-parkinson drug UCB0599. *Proc. Natl. Acad. Sci. U. S. A.* 120, e2201910120. doi: 10.1073/pnas.2201910120
- Serdiuk, T., Fleischmann, Y., Ghosh, D., Delparente, A., Reber, V., Frey, L., et al. (2025). Alpha-synuclein interacts with regulators of ATP homeostasis in mitochondria. *Nat. Commun.* 16, 7651. doi: 10.1038/s41467-025-62895-4
- Serdiuk, T., Redeker, V., Savitschenko, J., Neupane, S., Haenseler, W., Fleischmann, Y., et al. (2026). Structure-function relationship of alpha-synuclein fibrillar polymorphs derived from distinct synucleinopathies. *Mol. Syst. Biol.* 22, 868–901. doi: 10.1038/s44320-026-00199-5
- Siebnner, A. S., and Uversky, V. N. (2026). Amyloid- β , tau protein, α -Synuclein, TDP-43, and FUS in mixed pathology: and intrinsic disorder to rule them all. *Int. J. Mol. Sci.* 27, 3669. doi: 10.3390/ijms27083669
- Singh, K. S., Verma, R., Singh, N., Singh, L. R., and Gupta, A. (2025). Factors responsible for alpha-Synuclein aggregation. *Prog. Mol. Biol. Transl. Sci.* 211, 271–292. doi: 10.1016/b.smbts.2024.11.004
- Son, A., Pankow, S., Bamberger, T. C., and Yates, J. R. (2023). Quantitative structural proteomics in living cells by covalent protein painting. *Methods Enzymol.* 679, 33–63. doi: 10.1016/b.s.mie.2022.08.046
- Sun, X., Pedretti, R., Dyson, H. J., Saelices, L., and Wright, P. E. (2025). Mechanism of inhibition of acid-mediated transthyretin aggregation by designed peptides. *J. Biol. Chem.* 301, 110594. doi: 10.1016/j.jbc.2025.110594
- Susa, A. C., Lippens, J. L., Xia, Z., Loo, J. A., Campuzano, I. D. G., and Williams, E. R. (2018). Submicrometer emitter ESI tips for native mass spectrometry of membrane proteins in ionic and nonionic detergents. *J. Am. Soc. Mass Spectrom.* 29, 203–206. doi: 10.1007/s13361-017-1793-x
- Tambini, M. D., Yin, T., Yesiltepe, M., Breuillaud, L., Zehntner, S. P., d'Abramo, C., et al. (2023). A β 43 levels determine the onset of pathological amyloid deposition. *J. Biol. Chem.* 299, 104868. doi: 10.1016/j.jbc.2023.104868
- Tang, H., Andrikopoulos, N., Li, Y., Ke, S., Sun, Y., Ding, F., et al. (2025). Emerging biophysical origins and pathogenic implications of amyloid oligomers. *Nat. Commun.* 16, 2937. doi: 10.1038/s41467-025-58335-y
- Townsend, J. A., and Marty, M. T. (2023). What's the defect? Using mass defects to study oligomerization of membrane proteins and peptides in nanodiscs with native mass spectrometry. *Methods* 218, 1–13. doi: 10.1016/j.ymeth.2023.07.004
- Ubbiali, D., Fratini, M., Piersimoni, L., Ihling, C. H., Kipping, M., Heilmann, I., et al. (2022). Direct observation of “elongated” conformational states in α -Synuclein upon liquid-liquid phase separation. *Angew. Chem. Int. Ed. Engl.* 61, e202205726. doi: 10.1002/anie.202205726
- Utami, K. H., Morimoto, S., Mitsukura, Y., and Okano, H. (2025). The roles of intrinsically disordered proteins in neurodegeneration. *Biochim. Biophys. Acta Gen. Subj.* 1869, 130772. doi: 10.1016/j.bbagen.2025.130772
- Venter, A. R. (2025). Protein analysis by desorption electrospray ionization mass spectrometry. *Mass Spectrom. Rev.* 44, 757–772. doi: 10.1002/mas.21900
- Villacob, R. A., Feizi, N., Beno, S. C., and Solouki, T. (2024). Collision-induced unfolding, tandem MS, Bottom-up proteomics, and interactomics for identification of protein complexes in native surface mass spectrometry. *J. Am. Soc. Mass Spectrom.* 35, 13–30. doi: 10.1021/jasms.3c00261
- Vlk, M., Muck, A., Hey, J. A., Schaefer, J. F., Hubálek, M., and Cvačka, J. (2026). Analysis of amyloid beta oligomers by cyclic ion mobility-mass spectrometry. *Anal. Bioanal. Chem.* 418, 2273–2286. doi: 10.1007/s00216-026-06349-w
- Wagner, W. J., and Gross, M. L. (2024). Using mass spectrometry-based methods to understand amyloid formation and inhibition of alpha-synuclein and amyloid beta. *Mass Spectrom. Rev.* 43, 782–825. doi: 10.1002/mas.21814
- Wang, X., Liang, T., Jin, A., Zhang, C., Zhou, J., Li, M., et al. (2025a). C-Terminal radical oxidation inhibits α -Synuclein aggregation and cytotoxicity via an oxidative oligomer-disrupting pathway. *J. Am. Chem. Soc.* 147, 23136–23144. doi: 10.1021/jacs.5c06792
- Wang, X., Wang, N., Zhou, P., Guo, R., Song, N., Zhang, Z., et al. (2025b). Oligomerization propensity governs self-assembly morphology: insights from steady-state oligomer analysis. *Anal. Chem.* 97, 21863–21872. doi: 10.1021/acs.analchem.5c02258
- Weckslar, A. T., Wang, L., Bernstein, L. J., Huang, R. Y.-C., Gupta, S., Kristensen, L. G., et al. (2026). Recommendations and considerations for hydroxyl radical protein footprinting-mass spectrometry. *Nat. Methods* 23, 1089–1099. doi: 10.1038/s41592-026-03083-0
- Wei, J., Meisl, G., Dear, A. J., Michaels, T. C. T., and Knowles, T. P. J. (2025). Kinetics of amyloid oligomer formation. *Annu. Rev. Biophys.* 54, 185–207. doi: 10.1146/annurev-biophys-080124-122953
- Wong, K. F. C., Greatorex, R. E., Gidman, C. E., Rahman, S., and Griffiths, R. L. (2023). Surface-sampling mass spectrometry to study proteins and protein complexes. *Essays Biochem.* 67, 229–241. doi: 10.1042/EBC20220191
- Xu, Y., Maya-Martinez, R., Guthertz, N., Heath, G. R., Manfield, I. W., Breeze, A. L., et al. (2022). Tuning the rate of aggregation of hIAPP into amyloid using small-molecule modulators of assembly. *Nat. Commun.* 13, 1040. doi: 10.1038/s41467-022-28660-7
- Young, L. M., Mahood, R. A., Saunders, J. C., Tu, L.-H., Raleigh, D. P., Radford, S. E., et al. (2015). Insights into the consequences of co-polymerisation in the early stages of IAPP and A β peptide assembly from mass spectrometry. *Analyst* 140, 6990–6999. doi: 10.1039/c5an00865d
- Yu, J., Uzuner, U., Long, B., Wang, Z., Yuan, J. S., and Dai, S. Y. (2023). Artificial intelligence-based HDX (AI-HDX) prediction reveals fundamental characteristics to protein dynamics: mechanisms on SARS-CoV-2 immune escape. *iScience* 26, 106282. doi: 10.1016/j.isci.2023.106282
- Zampar, S., Di Gregorio, S. E., Grimmer, G., Watts, J. C., and Ingelsson, M. (2024). “Prion-like” seeding and propagation of oligomeric protein assemblies in neurodegenerative disorders. *Front. Neurosci.* 18, 1436262. doi: 10.3389/fnins.2024.1436262
- Zetterberg, H., and Bendlin, B. B. (2026). Biofluid biomarkers in Alzheimer's disease and other neurodegenerative dementias. *Nature* 650, 49–59. doi: 10.1038/s41586-025-10018-w
- Zhang, J. Y., Dregni, A. J., and Hong, M. (2026). Heterogeneous dynamics of the fuzzy coat of full-length phospho-mimetic tau fibrils. *J. Am. Chem. Soc.* 148, 1623–1637. doi: 10.1021/jacs.5c18540