

Cytotoxic Evaluation of Trastuzumab in HER2-Positive MDA-MB-453 Breast Cancer Cells

Saranraj S. and Priya R.*

Department of Bioinformatics, School of Life Sciences, Vels Institute of Science Technology and Advanced Studies, Chennai 600117, Tamil Nadu, INDIA

*priya.microbio@gmail.com

Abstract

This study aims to provide an integrated computational and experimental approach in the development of recombinant trastuzumab with specificity towards the HER2-positive breast cancer cell surface antigen. Using the molecular docking approach with the HADDOCK protocol, the interaction between the antibody and the antigen is shown to be stable through the computation of the docking scores and the low intermolecular RMSD. In addition, the electrostatic and van der Waals components of the interaction are significant. Using the information obtained in the molecular docking studies, a mammalian expression construct of the antibody is generated and is shown to be successfully propagated in Escherichia coli. Using the transformed cells, the recombinant antibody is shown to be successfully produced in ExpiCHO-S™ cells with sustained cell viability.


Using affinity chromatography and size exclusion chromatography-high performance liquid chromatography (SEC-HPLC) analysis, the antibody is shown to be highly homogeneous with minimal aggregation and is further confirmed through the analysis of the structure of the antibody using SDS-PAGE in reducing and non-reducing conditions. The functional evaluation by MTT assay showed a dose-dependent decrease in cell viability in MDA-MB-453 cells, with an IC₅₀ value of ~700 nM. Scratch wound

healing assays showed significant inhibition in cell migration whereas AO/EtBr staining and DNA fragmentation assays confirmed apoptosis-mediated cell death. Overall, these results validate the synthesis of high-purity biologically active rTrastuzumab and its potent cytotoxic, anti-proliferative and anti-migratory activities, making it relevant in targeted breast cancer therapy.

Keywords: Recombinant Trastuzumab, HER2-positive Breast Cancer, Molecular Docking (HADDOCK), Antibody Expression and Purification, Apoptosis and Cytotoxicity, SEC-HPLC Characterization.

Introduction

Breast Cancer Overview: The most common cancers in women all over the world is breast cancer. However, it still represents a significant health problem in terms of cancer-related morbidity and mortality¹. Despite significant progress in early detection methods, diagnostic tools and treatment regimens for breast cancer patients, this disease represents a significant problem due to its heterogeneity and the potential to develop drug-resistant strains of cancer². New therapeutic methods using targeted therapy methods are essential to improve the accuracy of diagnosis and increase the survival rate of patients suffering from breast cancer due to its heterogeneous nature³. Therefore, one of the major aims of modern cancer therapy is to develop targeted therapy methods to treat breast cancer in an accurate manner by focusing on the molecular alterations in cancer⁴.



Graphical Abstract

Role of HER2 Receptor in Breast Cancer: One of the most important biological markers related to aggressive forms of breast cancer is Human Epidermal Growth Factor Receptor 2 (HER2)^{5,6}. This receptor is a transmembrane tyrosine kinase receptor and is part of the receptor family for epidermal growth factor⁷. Cell proliferation, differentiation and survival are heavily dependent on this receptor⁸. Overexpression of the HER2 gene is related to a poor clinical prognosis in 15–20% of breast cancer patients^{9,10}. This is also associated with a higher probability of metastasis and rapid progression of cancer. Overexpression of HER2 leads to uncontrolled cell division and proliferation¹¹. This occurs due to the activation of this receptor, which initiates a series of cell signal transduction pathways such as the MAPK and PI3K/AKT pathways^{12,13}.

Mechanism of Action of Trastuzumab: Targeted therapy against the HER2 protein has significantly improved the outcomes in patients with HER2 positive breast cancer¹⁴. The most powerful drug that has been used in the treatment of HER2 positive breast cancer, is trastuzumab which is a monoclonal antibody that has been humanized¹⁵. Trastuzumab has the ability to target and bind to the HER2 protein, specifically the extracellular domain of the HER2 receptor^{16,17}. Binding of the trastuzumab antibody to the HER2 receptor prevents the dimerization of the receptors leading to the signaling cascade that leads to the proliferation of cancerous cells¹⁸. ADCC also leads to the killing of cancerous cells by the trastuzumab antibody¹⁹. These mechanisms have been proven to control the development of cancerous cells in patients with HER2 positive breast cancer and increase the survival rate of the patients^{20,21}.

Importance of Cytotoxicity Evaluation: It is essential to measure the potency of anticancer drugs in cell cultures in order to understand the biological activities of therapeutic antibodies²². It is also possible to determine the effective dosage of an anticancer drug and obtain critical information regarding anticancer activities by utilizing *in vitro* cytotoxicity studies^{23,24}. The cytotoxic activities of anticancer drugs can be examined in breast cancer cells in order to understand the response mechanisms and therapeutic potential of an anticancer drug, such as trastuzumab, particularly in understanding the survival/growth inhibition effects on HER2 positive breast cancer cells in order to determine the sensitivity/resistance profile of the medication^{25,26}.

In light of the therapeutic promise of HER2-targeted therapy, the current study aims to examine the cytotoxic and anti-proliferative effects of trastuzumab on HER2-positive breast cancer cells²⁷. The purpose of this MDA MB 453 breast cancer cell line study is to look into how trastuzumab treatment affects cell viability and proliferation^{28,29}. In addition to highlighting trastuzumab's therapeutic promise in the treatment of targeted breast cancer, this study attempts to ascertain the drug's inhibitory potential on HER2-positive cancer cells³⁰.

Material and Methods

Materials and Reagents: The analytical grade chemicals and reagents used in this study do not need any further purification. *Escherichia coli* cells with chemical competence Invitrogen (USA) supplied the tris-EDTA solution and One Shot(r) TOP10. HiMedia Laboratories (India) supplied Luria Bertani broth, Luria Bertani agar (HiVeg agar, Miller formulation) and other media used for microbiology. Ampicillin, potassium chloride (KCl), sodium hydroxide (NaOH), sodium dodecyl sulfate (SDS), agarose, bromophenol blue and phosphate buffered saline solution were used for bacterial culture and other molecular biology studies.

The cell culture and protein expression media and reagents were supplied by Gibco (Thermo Fisher Scientific, USA) and they included OptiPRO™ serum-free medium, ExpiCHO™ Expression medium, ExpiCHO-STM cells, ExpiFectamine™ CHO reagent, ExpiCHO™ feed and ExpiFectamine™ CHO enhancer. The plasmid was purified using the QIAprep Spin Miniprep kit supplied by Qiagen (Germany). The protein was purified using protein A magnetic beads. The analytical tests were carried out using general reagents.

Cell Line and Cell Culture Conditions: Monoclonal antibody trastuzumab is produced using ExpiCHO-S cells and recombinant DNA methods. The cells were rapidly thawed in a water bath at 37 °C. Frozen cells were aseptically transferred to a number of Erlenmeyer flasks containing ExpiCHO™ expression Medium at 37 °C. The flasks were placed in a 37 °C, 8% CO₂ and 125 rpm humidified incubator. Trypan blue viability test and Neubauer's hemocytometer were generally used to monitor cell growth. The cells were sub-cultured to a viable cell density of about 4-6 × 10² cells/mL to maintain optimum growth conditions. The culture was grown to a density required to carry out transient transfection and protein expression.

Drug Preparation (Trastuzumab Concentration): Using the cytomegalovirus (CMV) promoter as a model, the optimized coding sections of the heavy and light chains of the recombinant trastuzumab gene construct were incorporated into the mammalian expression vector pcDNA3.1(+). After the heat shock transformation of the chemically competent *E. coli* cells with the plasmid vector, the vector was grown in LB medium with 100 µg/mL of ampicillin in a nutrient agar plate.

The plasmid purification was carried out using a QIAprep Spin Miniprep kit and was done following the instructions provided by the manufacturer. A NanoDrop spectrophotometer was used to determine the quantity of purified plasmid. The purified plasmid was stored at -20 °C in a freezer for future use. The plasmid was transfected into ExpiCHO-S cells for transient expression using an ExpiFectamine™ CHO transfection reagent following the instructions provided by the manufacturer.

MTT Assay Procedure: The cytotoxicity of trastuzumab was assessed by using the MTT assay. It is one of the widely used methods for assessing cell survival and growth. Breast cancer cells were inoculated in appropriate cell density in a 96-well plate under normal growth conditions. The cells were allowed to adhere to the plate overnight and were treated with varying concentrations of trastuzumab. Each well was treated with MTT reagent to allow live cells to convert tetrazolium salt into insoluble crystals of formazan. The crystals formed were dissolved in solubilizing solution and read spectrophotometrically at 570 nm using a microplate reader.

Cell Viability Calculation: The MTT assay test's absorbance readings were used to calculate the cell viability. The treated cells' absorbance values were contrasted with those of the untreated cells, which served as controls. Cell vitality (%) is calculated by dividing the absorbance values of treated cells by the absorbance values of control cells and multiplying the result by 100. This experiment's computation revealed that trastuzumab inhibits the proliferation of breast cancer cells.

IC₅₀ Determination: Plotting the cell viability percentage in relation to the logarithmic concentration of the drug enabled the calculation of the half-maximum inhibitory concentration value for trastuzumab. Using appropriate statistical tools for data analysis, a dose-response curve for the drug was constructed. This helped in calculating the concentration required for reducing the cell viability by 50% in comparison to the control cells. This value is represented as the IC₅₀ value for trastuzumab in relation to the specified breast cancer cell line.

Statistical Analysis: All experiments were conducted in triplicate to ensure reproducibility and reliability of results. The results were expressed as mean $\pm$ standard deviation (SD). Statistical analysis for comparing treated and control group data was done using appropriate statistical tests. The results were considered statistically significant at $p < 0.05$.

Cancer cell lines

MTT Assay: MTT assay, a colorimetric method that measures cell viability by mitochondrial metabolic activity, was utilized to study the cytotoxic and antiproliferative activities of trastuzumab against MDA-MB-453 human breast cancer cells. MTT, a yellow tetrazolium compound, is reduced to purple formazan crystals by mitochondrial dehydrogenase enzymes present only in viable cells³¹. The quantity of formazan crystals formed is directly proportional to the number of viable cells, as this reaction only takes place in metabolically active cells. Absorbance of dissolved formazan crystals is then used to measure the vitality of cells spectrophotometrically compared to untreated cells³².

To grow MDA-MB-453 breast cancer cells, standard conditions were applied and then these cells were placed in a sterile 96-well plate with 7,000 cells per well, using a

complete growth medium. Before exposing these cells to any drug, they were allowed to attach properly to the plate, taking one day. After this, a new medium containing different concentrations of trastuzumab (100 nM to 1000 nM) was used to replace the growth medium. To ensure reproducibility, all experiments were done in triplicate.

MTT reagent was added to each well after 48 hours of drug exposure. The plates were incubated to allow the live cells to convert MTT to formazan crystals. The growth medium was carefully removed and the crystals were dissolved in the appropriate solvent. A microplate reader was used to test for absorbance at the suggested wavelength. The half-maximal inhibitory concentration (IC₅₀) of trastuzumab was ascertained from the dose-response curve. Cell viability (%) was calculated based on the comparison of the absorbance in the treated wells with that in the control wells which were assumed to be 100% viable.

Scratch (Wound Healing) Assay: The scratch (wound healing) assay was applied to determine the effect of trastuzumab on the migration of MDA-MB-453 human breast cancer cells. The scratch assay, in which the scratch in the confluent cell layer represents the wound, monitors the migration of cells into the wounded area over time, simulating the process of collective cell migration *in vitro*. The migratory ability of the cells is measured as the rate of wound healing³³. The assay was done in low serum conditions to ensure that the wound healing occurs as a result of cell migration rather than proliferation³⁴.

A near-confluent monolayer was obtained by seeding the cells at a density of 15,000 cells per well in 96-well culture plates and maintaining them under conventional growth conditions. Subsequently, a cell-free wound area was created by applying a scratch wound across the surface of the cells in each well of the 96-well culture plate using a sterile pipette tip. The cells were then removed by washing with phosphate-buffered saline solution and the cells were cultured in 0.1% fetal bovine serum to minimize the effect of proliferation-mediated wound closure.

The trastuzumab was added to the cells with varying concentrations of 600 nM, 700 nM and 800 nM. The control cells were not exposed to trastuzumab. To ensure that the experiment is replicable, each condition was repeated three times. The images were captured using an inverted microscope with the same settings at 24, 48, 72 and 96 hours after scratch wound creation at 0 h. The percentage wound closure was calculated relative to the initial wound area after measuring the wound area using ImageJ software with the plugin for wound healing.

AO/EtBr Dual Staining Assay: The acridine orange/ethidium bromide (AO/EtBr) dual labeling technique was utilized to qualitatively examine the induction of apoptosis by Trastuzumab on human breast cancer cells MDA-MB-453³⁵. The AO/EtBr technique is a fluorescence-

based cell viability assay that differentiates apoptotic cells from necrotic cells, thereby confirming whether cell death is through apoptosis or not³⁶.

In cells with intact membrane integrity, the cell-permeable fluorescent compound acridine orange (AO) binds to the nucleic acids and fluoresces green. On the contrary, the orange or red fluorescence of ethidium bromide (EtBr) bound to DNA is due to its inability to permeate the cell membrane and enter cells with intact membrane integrity, which is typical of viable cells but not apoptotic or necrotic cells. Apoptotic cells are brilliant green or yellow in fluorescence with chromatin condensation whereas necrotic cells are orange with nuclear disintegration and have intense red fluorescence.

The drug was given to MDA-MB-453 cells that were cultured in normal conditions and ranged in concentration from 100 nM to 800 nM. Cells that were not given the drug were used as the control. Phosphate-buffered saline solution (PBS) was used to gently wash the cells after the treatment period. This is done in order to remove any traces of the medium. After that, the cells were stained with a newly synthesized AO/EtBr staining solution and were given time to absorb the solution in the dark.

In order to reduce photobleaching, the cells were observed immediately using a fluorescent microscope at 10x. To ensure that representative results are observed, multiple areas were examined.

DNA Fragmentation Assay: Verification of apoptosis resulting from treatment with trastuzumab using a DNA fragmentation technique, wherein specific patterns of DNA fragmentation have been correlated with programmed cell death, was accomplished. One of the key biochemical hallmarks of apoptosis is DNA fragmentation, which results from the action of intracellular endonucleases cleaving cellular DNA at internucleosomal locations³⁷. The cellular DNA of non-apoptotic or live cells appears as a high molecular weight band on agarose gel electrophoresis whereas apoptotic cells display fragmented or smeared DNA. Hence, verification of apoptosis is confirmed by the appearance of fragmented or smeared DNA³⁸.

On the basis of previous cytotoxicity and apoptosis screening results, cells of MDA-MB-453 were treated with trastuzumab at specific concentrations³⁹. The cells were left untreated and used as the control. The cells were collected after treatment and proteins and cell debris were removed using normal extraction methods⁴⁰. The genomic DNA was isolated. The gel was prepared with 0.8% agarose gel and an appropriate buffer for electrophoresis. The gel was then filled with an equal amount of DNA from both the treated and untreated cells. To separate DNA of specific sizes, electrophoresis was performed for one hour at 100 V. To assess the migration of the DNA, it was visualized using a gel documentation system.

Results and Discussion

Molecular docking of Trastuzumab with HER2: The HADDOCK 2.4 server was used for the docking of the trastuzumab protein and the HER2 receptor, with the aim of identifying the most stable conformations for the complex formed by the two proteins and understanding the molecular basis of the protein-protein interaction. A number of docked protein complex structures were generated through the HADDOCK clustering analysis. The HADDOCK score, cluster size, RMSD and energy terms, such as the van der Waals and electrostatic interactions, are used for the arrangement of the docked complex structures, as presented in table 1.

Among all the docked complex structures generated through the docking analysis, the best docked complex was found in the second cluster, with the lowest HADDOCK score of -53.0 ± 6.9 and a significant negative Z-score of -1.3 compared to the entire docking population, showing significant statistical significance.

With a moderate cluster size of 13 structures and a low RMSD value of 5.5 ± 0.3 Å, this cluster showed good topological convergence and binding fidelity. The favorable van der Waals (-33.6 ± 3.5) and electrostatic energy (-79.8 ± 31.6) values ensured the binding. This shows the importance of hydrophobic and electrostatic interactions during the complex formation. The stability and uniqueness of the binding mode were ensured by cluster 3, which had a high HADDOCK score of -52.0 ± 3.5 , Z-score of -1.2 and low RMSD values of 4.3 ± 0.7 Å. This cluster also had high electrostatic interactions (-129.1 ± 20.4), showing the importance of electrostatic interactions during the binding of trastuzumab to the HER2 epitope.

However, a slightly lower amount of conformational sampling is indicated by the cluster size being slightly lower than that of cluster 2. Cluster 5 had the smallest RMSD value (3.1 ± 0.3 Å) and thus indicated a relatively constant docking orientation, despite being less favorable in terms of Z-score (-0.8) and weaker van der Waals connections. This cluster was ranked lower because it was less energetically favorable, even though it was structurally stable.

On the other hand, the clusters with positive Z-score values (Clusters 8, 11 and 12) had lower interaction energy, HADDOCK score and RMSD, which indicated that the binding conformations were not optimal. Thus, the probability of the clusters indicating strong interactions between the two molecules i.e. HER2 and Trastuzumab, is low. The therapeutic efficacy of trastuzumab in the treatment of HER2-positive breast cancer is a clear indicator of the strong and specific binding of the antibody to the HER2 receptor, which is supported by the results of the docking research. The thermodynamically stable complex formed by the antibody and the receptor is indicated by the high number of negative Z-scores and HADDOCK scores within the top-ranked clusters.

Of particular interest is the balance of electrostatic and van der Waals forces, which highlights the level of cooperation within the molecular interactions that have been at play during the trastuzumab-HER2 interaction. Figure 1A shows a scatter plot that highlights the correlation between the percentage of common connections (FCC) and the HADDOCK score of the docked Trastuzumab-HER2 complexes. Although the colored points with standard deviation error bars represent the cluster representatives, the points represent the docking solutions. More favorable docking processes are indicated by the HADDOCK score and FCC. In figure 1B, the electrostatic interaction energy of the Trastuzumab-HER2 docking clusters is represented as a function of the interface root mean square deviation.

Each data point represents a unique docking attitude and standard deviation error bars are given by the points. The stability and energy of the antibody-antigen complex are implied by the points with lower electrostatic energy and lower i-RMSD values. The graph showing the van der Waals interaction energy of the docked Trastuzumab-HER2 complexes for various values of i-RMSD is given on the correlation graph in figure 1C. The graph represents the average standard deviation of the clusters. The higher negative values of the van der Waals interaction energy at the interface and lower values of i-RMSD imply an ideal antibody-antigen complex with beneficial hydrophobic interactions.

Table 1 Calculated HADDOCK scores and related parameters of trastuzumab and HER2						
Clusters	HADDOCK Score	Cluster Size	RMSD	Van der Waals Energy	Electrostatic Energy	Z-score
1	-46.2 ± 0.9	43	6.7 ± 0.1	-32.0 ± 5.3	-81.8 ± 29.3	-0.1
2	-53.0 ± 6.9	13	5.5 ± 0.3	-33.6 ± 3.5	-79.8 ± 31.6	-1.3
3	-52.0 ± 3.5	11	4.3 ± 0.7	-26.0 ± 1.8	-129.1 ± 20.4	-1.2
4	-47.8 ± 5.0	10	7.3 ± 0.3	-24.0 ± 4.0	-137.6 ± 31.7	-0.4
5	-50.0 ± 7.2	8	3.1 ± 0.3	-17.3 ± 5.1	-172.4 ± 47.1	-0.8
6	-45.7 ± 5.0	7	3.4 ± 0.5	-14.8 ± 4.3	-177.9 ± 39.1	-0.0
7	-44.0 ± 3.1	7	8.8 ± 0.1	-34.4 ± 3.0	-68.5 ± 8.1	0.3
8	-41.7 ± 9.9	5	6.9 ± 0.3	-21.5 ± 4.7	-118.4 ± 49.2	0.7
11	-33.0 ± 4.1	4	8.6 ± 0.1	-28.1 ± 2.5	-58.0 ± 24.5	2.2
12	-41.3 ± 8.3	4	6.2 ± 0.1	-19.8 ± 5.8	-113.9 ± 35.0	0.7




Figure 1: (A) HADDOCK score versus fraction of common contacts (FCC). (B) Electrostatic energy versus interface RMSD. (C) Van der Waals energy versus interface RMSD. Cluster-wise distribution of (D) electrostatic interaction energies and (E) van der Waals interaction energies.

The electrostatic interaction energy distribution of the HADDOCK clusters is shown in figure 1D as a box and whisker graph. The box and whisker graph shows the interquartile range, the median and the outliers of the electrostatic interaction energy. The stability of the Trastuzumab-HER2 bond can be explained by the increased long-range electrostatic interaction, as shown by those with lower median electrostatic interaction energy. The box-whisker plot is a graphical method of displaying the van der Waals interaction energy for a given cluster of the HADDOCK clusters (Figure 1E). The variation in the median can be attributed to the different packing efficiencies at the interface. The findings provide a rationale for the use of trastuzumab as a targeted medication for HER2-positive cancer cells and provide molecular evidence of the targeted antibody specificity of the compound.

Construction of Trastuzumab vector: A short flexible linker sequence of 48 bp was employed to link the trastuzumab HC (469 amino acids; ~1407 bp) and LC (233 amino acids; ~699 bp). This ensured that there was proper coordination in transcription and translation. This approach enables the balanced expression of both chains of antibodies. This is important for the efficient assembly and secretion of antibodies in mammalian cells. The humanized monoclonal antibody heavy chain (HC) and light chain (LC), as a whole antibody, are produced simultaneously by creating the expression vector for trastuzumab. The expression of the humanized antibody can now be carried out on the created plasmid. The total size of the constructed plasmid is about 7508 bp.

The constructed plasmid has all the essential regulatory elements. These include transcriptional regulatory areas, polyadenylation signals, a strong promoter and enhancer region and indicators that confer antibiotic resistance to bacteria. Additionally, a high copy replication origin in the plasmid backbone allowed for replication and amplification of the constructed vector in *E. coli*. Figure 2 is a vector diagram representing the general architecture of the project. All the aforementioned results indicate that a stable and well-constructed trastuzumab expression vector has been successfully constructed.

Transformation and Colony formation: After constructing the vector, one shot TOP10 chemically competent *E. coli* cells were transformed by heat shock to incorporate the recombinant trastuzumab plasmid. In order to allow selective growth of *E. coli* containing the plasmid, transformed *E. coli* were grown on LB agar plates containing ampicillin. Figures 3A and 3B describe the formation of various distinct colonies on the plates after an overnight incubation at 37°C. The high density of colonies formed indicates the competency of the host cells as well as the stability of the recombinant vector produced, in addition to high transformation efficiency and uptake of the plasmid by *E. coli*. The shape of the colonies, boundaries and their constant size showed homogenous plasmid maintenance and

the lack of any growth inhibition effects resulting from the additional expression cassette of trastuzumab.

Twelve randomly selected colonies from the transformation plate underwent rapid (quick) plasmid DNA isolation. DNA bands of the expected size for the trastuzumab expression vector were observed when these separated plasmids underwent 1% agarose gel electrophoresis (Figure 3C). This was the screening phase for the stable maintenance of the recombinant plasmid.

Plasmid isolation and screening of positive clones: Based on the quality and intensity of the plasmid bands, colonies 1, 6, 9 and 12 were selected for further investigation as positive clones based on the agarose gel results. To ensure that the clones were pure and that the plasmids were being kept stable in the bacteria cells, the colonies were restreaked onto new LB agar plates containing ampicillin. Figure 4A to figure 4D shows the results of the colonies that contained the transformed bacteria growing strongly on the selective media. This suggested that the bacteria were being kept in a state of resistance to the antibiotic and that the plasmids were being kept stable.

Using the QIAprep Spin Miniprep kit, high-quality plasmid DNA was extracted from the four colonies that were selected in accordance with the manufacturer's instructions. The isolation process ensured that the plasmid DNA was intact with minimal degradation, as confirmed using agarose gel electrophoresis. From the presence of distinct bands in the gel electrophoresis results of all four colonies, it is safe to conclude that the recombinant trastuzumab expression vector was effectively lysed, purified and recovered (Figure 4E). To determine the quality and quantity of the purified plasmids, quantitative and subjective analysis using a NanoDrop Lite spectrophotometer was carried out.

From the results, the A260/A280 ratio of each of the four colonies was found to be 1.90, indicating minimal protein contamination and maximum purity of the plasmid. Since the plasmid concentrations were 1396.4 ng/μL, 1329.7 ng/μL, 1098.3 ng/μL and 911.1 ng/μL in colonies 1, 6, 9 and 12, respectively, the experiment demonstrated the amplification of the plasmids in the chosen colonies. The highest concentrations of plasmid were observed in colonies 1 and 6, which indicated a better copy number and/or growth rate. All these results indicated the successful transformation, growth and isolation of the recombinant trastuzumab expression plasmid.

ExpiCHO-STM cells morphology following Trastuzumab transfection: The morphological features of ExpiCHO-STM cells were examined on day 14 post-transfection to assess the viability and flexibility of ExpiCHO-STM cells following transitory expression of the trastuzumab construct (Figure 5A). Under microscopic examination using a 40x objective lens, ExpiCHO-STM cells were observed to maintain their characteristic

morphological features including being spherical in shape and smooth boundary and size distribution. Shear-sensitivity is a characteristic feature of CHO cell cultures and such cultures were successfully sustained for the entire duration of the prolonged expression phase, as evidenced by the presence of a majority of suspended ExpiCHO-STM cells in the culture medium without any sign of aggregation.

Moreover, a low degree of cell lysis was observed and a large number of cells were able to withstand the natural decrease in viability characteristic of the late stage of transient expression. The survivability of ExpiCHO-STM cells on day 14 post-transfection indicates that trastuzumab expression does not significantly produce harmful effects on ExpiCHO-STM cells and the ExpiCHO-STM procedure is capable of long-term expression of recombinant antibody. The outcomes are consistent with ExpiCHO-STM's demonstrated resilience for high-yield monoclonal antibody expression and the subsequent extraction of trastuzumab antibody from ExpiCHO-STM cell supernatant.

Purification and SEC-HPLC analysis of recombinant Trastuzumab: The purified and clarified fractions of Trastuzumab, after affinity purification using protein A magnetic beads, were analyzed using Size Exclusion

Chromatography with High-Performance Liquid Chromatography (SEC-HPLC). This technique is used to determine the molecular homogeneity, purity and aggregation of the protein. The analysis using SEC-HPLC, which is based on the measurement of the UV absorbance of the protein at 280 nm (Figure 5B), is a sensitive technique and can be used to monitor the proteins that are being eluted from the column. The chromatogram has indicated that 95% of the total peak area is comprised of a large peak with a retention time of 12.3 minutes, corresponding to the monomer of the trastuzumab protein, as expected due to the molecular size.

The small peaks that were found to elute early (8.7–10.2 minutes) constituted less than 5% of the total area, implying that they were minute contaminants or traces of high molecular weight substances. Most importantly, the antibody was found to be largely monodisperse after synthesis and purification, as evidenced by the lack of aggregation peaks. The large area percentage of the main peak is a clear indication of the high level of purity and integrity of the expressed recombinant trastuzumab. The lack of significant low molecular weight degradation products is a clear indication of proteolytic cleavage during expression.




Figure 2: General framework of well-constructed Trastuzumab expression vector

The figure shows a circular Trastuzumab expression vector (~7508 bp) designed for monoclonal antibody production. It contains CMV promoter/enhancer elements for high-level expression of heavy and light chain genes. The vector includes signal peptides, poly-A signals and dual expression cassettes to ensure proper transcription and secretion. Selectable markers (AmpR) and origins of replication (SV40 ori, bacterial ori) enable propagation in host systems. Multiple restriction sites are present for cloning and genetic modifications.




Figure 3: The formed well-isolated colonies on the plates of (A) Control and (B) Trastuzumab transformed cells. (C) The DNA band separation of twelve colonies observed in 1% agarose gel electrophoresis.

Selection of purified Trastuzumab for SDS-PAGE analysis: Based on the SEC-HPLC analysis that indicated significant homogeneity and low levels of aggregation in the purified antibody, the purified trastuzumab containing the main monomer peak was selected for further analysis using SDS-PAGE. In addition, the results obtained using SEC-HPLC were critical in ensuring that the integrity and purity of the antibody were guaranteed prior to analysis using other methods. Therefore, the integrity and purity of the antibody are clearly demonstrated using affinity purification, SEC-HPLC and SDS-PAGE. This is critical in ensuring the accuracy of the recombinant trastuzumab analysis.

SDS-PAGE analysis of purified Trastuzumab under reducing and non-reducing conditions: The purified recombinant trastuzumab was obtained using the ExpiCHO-STM cell culture supernatant. The structural integrity and purity of the recombinant trastuzumab were analyzed using sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) in both reducing (R) and non-reducing (NR) conditions (Figure 5C). A pre-stained molecular weight

protein marker (Lane M1) with a molecular weight range of 10-250 kDa was used for the determination of the molecular weight of the purified trastuzumab bands. Under non-reducing conditions, the purified trastuzumab was observed as a single, distinct band with an apparent molecular weight of approximately 150 kDa. This band corresponds to the entire immunoglobulin G molecule, with the heavy and light chains still covalently attached by interchain disulfide bonds.

The clear and distinct appearance of this band is an affirmation of the native antibody structure during handling, purification and expression, which indicates a high degree of molecular homogeneity. Interestingly, no additional high molecular weight bands were detected above 150 kDa, which indicates that neither multimerization of the antibody, nor antibody aggregation, occurred during the experiment. In addition, proteolytic degradation is minimal when low molecular weight bands are not detected. Our findings are in agreement with the SEC-HPLC data, which detected a strong monomer peak with minimal aggregation.




Figure 4: Robust development of the transformed colonies on LB agar plates (A) Colony 1; (B) Colony 6; (C) Colony 9 and (D) Colony 12. (E) Clear and distinct plasmid bands seen in all four colonies.

As such, trastuzumab separated into its heavy and light chains as expected since the disulfide bonds in the structure of the antibody were broken during the analysis. The immunoglobulin heavy chain, as represented by the primary band, has an approximate molecular weight of 50 kDa, while the immunoglobulin light chain, as represented by the second band, has an approximate molecular weight of 25 kDa.

The distinct separation of the immunoglobulin heavy and light chains, as well as the appropriate molecular weights, confirms the proper structure of the antibodies, as well as the proper folding, assembly and post-translational processing of the ExpiCHO-STM expression system. The similarity in the band intensity of the heavy and light immunoglobulin chains also confirms that these two chains are expressed in the proper balance, as they must be in order to assemble efficiently into an IgG antibody. Lack of additional bands corresponding to truncated products, misfolded proteins and

host cell proteins indicates the high purity of the proteins achieved by affinity chromatography with protein A.

The SDS-PAGE profile clearly indicates the successful synthesis of structurally intact, full-length proteins of trastuzumab under reducing as well as non-reducing conditions. The presence of a single band of ~150 kDa under non-reducing conditions indicates strong disulfide bonding between chains, resulting in native structure whereas the appearance of distinct bands of ~50 kDa and ~25 kDa under reducing conditions indicates the sufficient synthesis of heavy and light chains of the antibody. The SDS-PAGE findings confirm the high level of purity and structural integrity of the purified recombinant trastuzumab, as expressed in ExpiCHO STM cells, in line with the SEC-HPLC findings which confirm the high level of molecular homogeneity and low level of aggregation in the purified trastuzumab, indicating that the purified trastuzumab would be suitable for further downstream applications.

MTT Assay: In addition, after the treatment of cells with the drug for 48 hours, trastuzumab treatment leads to a dose-dependent reduction in the viability of MDA-MB-453 cells. The dose of the drug, when raised to values above or equal to 600 nM, significantly reduced the number of viable cells. In contrast, lower values of the dose (in the range of 100-400 nM) showed a gradual reduction in metabolic activity. The cytotoxic effect of the drug was almost complete when the dose of the drug was in the range of micromolar values. The IC₅₀ value of the drug, as indicated by the dose-response curve, was about 700 nM.

The findings were further supported by morphological studies, as the cells treated with the drug showed rounded cells, reduced adhesions and lower cell density when compared to untreated cells. In addition, after the treatment of cells with the drug for 48 hours, trastuzumab treatment leads to a dose-dependent reduction in the viability of MDA-MB453 cells. The dose of the drug, when raised to values above or equal to 600 nM, significantly reduced the number of viable cells. In contrast, lower values of the dose (in the range of 100-400 nM) showed a gradual reduction in metabolic activity. The cytotoxic effect of the drug was almost complete when the dose of the drug was in the range of micromolar values.

The IC₅₀ value of the drug, as indicated by the dose-response curve, was about 700 nM. The findings were further supported by morphological studies, as the cells treated with the drug showed rounded cells, reduced adhesions and lower cell density when compared to untreated cells. The reliability of the assay method was further supported by the uniformity of the triplicates as shown in fig. 6.

Scratch (Wound healing) Assay: With time, the control wells indicated progressive wound closure, which was expected as a normal migratory response. On the other hand, the migration of cells was inhibited in a concentration-dependent manner after the cells were treated with trastuzumab. A partial inhibition of wound closure was observed at 600 nM. The migration of cells was significantly reduced at 700 nM and with time, very little cell migration was observed into the wound area. Strong cytotoxic effects were observed at 800 nM, as evidenced by the extension of the wound area with increased cell detachment. The percent of wound closure in the treated cells was significantly lower when compared with the control cells, as analyzed by ImageJ software, with the greatest effect observed at the higher dosages as shown in fig. 7.




Figure 5: (A) Morphological alterations in the ExpiCHO-STM cells post Trastuzumab transfection on Day 0 and Day 14. (B) SEC-HPLC analysis for protein selection based on size. (C) Sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) in both reducing (R) and non-reducing (NR) conditions




Figure 6: Dose-dependent cytotoxic effect of Trastuzumab on MDA-MB-453 cells. Representative phase-contrast microscopic images illustrating morphological alterations in MDA-MB-453 cells following treatment with increasing concentrations of Trastuzumab: (A) 100 nM, (B) 200 nM, (C) 300 nM, (D) 400 nM, (E) 500 nM, (F) 600 nM, (G) 700 nM, (H) 800 nM and (I) 1000 nM. Progressive loss of cell integrity, rounding and reduced cell density were observed with increasing concentrations. (J) Dose-response curve demonstrating the cytotoxic effect of Trastuzumab on MDA-MB-453 cells, from which the IC_{50} value was estimated to be approximately 700 nM. Data are presented as mean $\pm$ SD of triplicate experiments. Statistical significance was assessed using Student's t-test compared with the control ($p < 0.05$, $p < 0.01$, $*p < 0.001$; ns, not significant).




Figure 7: Scratch assay analysis of MDA-MB-453 cells treated with Trastuzumab. Representative images showing wound closure at 0, 24, 48, 72 and 96 h under 0.1% FBS conditions for control, 600 nM, 700 nM and 800 nM treatments. Images were analyzed using ImageJ software with the Wound Healing plugin (Montpellier R. Imaging Resources, GitHub). Wound closure was quantified to evaluate the effects of Trastuzumab on cell migration and proliferation in a time- and concentration-dependent manner.




Figure 8: AO/EtBr Dual Staining Assay
Representative images of acridine orange/ethidium bromide (AO/EtBr) dual staining in MDA-MB-453 cells following treatment with Trastuzumab. Panels A–I depict fluorescence images corresponding to control and increasing concentrations (100–800 nM), visualized under an inverted fluorescence microscope at 10× magnification. Panels J–R represent images of the scratched wound region from the respective treatment groups, captured under the same magnification. AO-positive viable cells exhibit green fluorescence, whereas EtBr-positive apoptotic or necrotic cells display orange to red fluorescence, indicating treatment-induced cytotoxic and apoptotic effects across concentrations.

Table 2
Quantitative analysis of wound closure area at different time intervals following treatment with Trastuzumab

Time (hrs)	Control Area	% Change	600 nM Area	% Change	700 nM Area	% Change	800 nM Area	% Change
0	414,460	—	335,002	—	422,401	—	229,577	—
24	6,750,426	—	5,324,943	—	5,285,388	—	3,056,046	—
48	6,596,269	−2.28%	5,142,583	−3.43%	4,131,067	−21.84%	3,245,782	+6.21%
72	6,119,328	−7.23%	4,057,425	−21.09%	3,512,376	−14.97%	3,420,503	+5.38%
96	6,008,452	−1.81%	5,000,910	+23.25%	3,292,240	−6.27%	3,647,595	+6.64%




Figure 9: Agarose gel electrophoresis showing DNA fragmentation patterns in MDA-MB-453 cells treated with Trastuzumab. DNA samples were analyzed from control cells and cells treated with 600 nM, 700 nM and 800 nM concentrations. The presence of DNA banding patterns indicates treatment-dependent DNA fragmentation compared with the control.

The quantitative analysis of wound closure area at different time intervals following treatment with Trastuzumab was shown in table 2. Images were analyzed using ImageJ software with the Wound Healing plugin (Montpellier R. Imaging Resources, GitHub) to calculate the wound area and assess the extent of wound closure over time.

AO/EtBr Dual Staining Assay: This was evident from the consistent fluorescence and morphology of the control cells. On the other hand, the apoptotic markers increased in the cells treated with trastuzumab in a concentration-dependent manner. Early apoptotic markers such as chromatin condensation and high-intensity green fluorescence in the nuclei were observed at lower concentrations (100–300 nM). More cells exhibited orange fluorescence at intermediate concentrations (400–600 nM), indicating the progression of apoptosis to the later stages. Significant fragmentation of the nuclei and orange fluorescence were observed at higher concentrations (700–800 nM), indicating apoptosis and secondary necrosis. The proportion of apoptotic cells increased with the gradual increase in the concentration of trastuzumab, which was consistent with the cytotoxic effects observed in scratch and MTT assay was shown in fig. 8.

DNA fragmentation Assay: A distinct undamaged band of high molecular weight in the DNA of the control cells indicated that the genome integrity is preserved. In contrast, the DNA of the cells treated with Trastuzumab exhibited distinct smearing and fragmentation, indicating internucleosomal DNA fragmentation. These patterns were consistent in the treated cells despite the low amount of DNA yield, indicating that apoptosis is occurring with no random degradation of DNA. There was a correlation between the amount of DNA fragmentation and increasing concentrations of Trastuzumab as shown in fig. 9.

Conclusion

This study utilizes the techniques: docking, molecular biology, mammalian expression and biochemical studies, which collectively create an effective process for the production of recombinant trastuzumab. The docking study, which utilized the HADDOCK docking method, created an initial framework by utilizing an investigation of the energetically favorable antibody-antigen interactions as confirmed by the low RMSD values, electrostatic and van der Waals contributions, which created confidence in the antibody design before its experimental confirmation. The successful creation, transformation and amplification of the Trastuzumab expression vector in *E. coli* created an effective method of plasmid production. By screening randomly selected colonies, extracting the plasmid and assessing its purity, the design's integrity and appropriateness for mammalian expression were verified.

Transient transfection into ExpiCHO-STM cells produced strong antibodies while maintaining good cell shape and high vitality over extended culture periods. Protein A-based affinity purification resulted in highly concentrated antibody preparations with minimal host cell protein contamination. SEC-HPLC quality assessment revealed that the pure trastuzumab primarily existed in a monomeric state with very little aggregation or degradation. This chromatographic screening allowed for the advancement of only structurally homogeneous protein fractions for electrophoretic analysis.

Although the reduction conditions resolved the antibody into its correct heavy and light chains, verifying correct folding, disulfide bond formation and assembly, the non-reducing SDS-PAGE also confirmed the correct assembly of the IgG molecule. The agreement between the experiment and the prediction is evident and confirms the reliability of the

integrated approach. This study emphasizes the importance of combining robust biochemical analysis with *in silico* prediction and validates the construction of high-quality recombinant trastuzumab molecules for future functional, biophysical and therapeutic applications.

The MTT assay showed that trastuzumab has significant cytotoxicity and anti-proliferative effects on MDA-MB-453 breast cancer cells after 24 hours of cell growth and 48 hours of drug treatment. The low nanomolar IC₅₀ value (~700 nM) indicates that the compound is highly effective. On this basis, it was determined that the IC₅₀ concentration would be used in all functional studies.

The scratch test showed that trastuzumab efficiently and dose-dependently prevents MDA-MB-453 cells from migrating. Higher doses caused cytotoxicity, which resulted in wound enlargement, but intermediate concentrations mainly decreased migratory capacity. These results validate trastuzumab's anti-migratory and anticancer properties. Trastuzumab causes apoptosis-mediated cell death in MDA-MB-453 breast cancer cells in a dose-dependent manner, according to AO/EtBr staining data.

Apoptosis is a key mechanism underlying the cytotoxic effects of trastuzumab, according to results on nuclear condensation, membrane integrity and fluorescence change from green to orange/red. Significant biochemical evidence that trastuzumab causes apoptosis-mediated cell death in MDA-MB-453 breast cancer cells, has been shown by the DNA fragmentation test. Apoptosis is a key mechanism underlying the cytotoxic effects of trastuzumab, according to results on DNA fragmentation patterns.

References

1. Akbarzadeh-Sharbat S., Yakhchali B., Minuchehr Z., Shokrgozar M.A. and Zeinali S., Expression enhancement in trastuzumab therapeutic monoclonal antibody production using genomic amplification with methotrexate, *Avicenna J. Med. Biotechnol.*, **5**, 87 (2013)
2. Amiri M.M., Bahadori T., Soltantoyeh T., Hosseini G.R., Golsaz S.F., Jeddi T.M. and Shokri F., Development of a novel inhibitory chimeric anti-HER2 monoclonal antibody, *Iranian Journal of Immunology*, **16**(1), 26–42 (2019)
3. Beck A., Wagner-Rousset E., Ayoub D., Van Dorsselaer A. and Sanglier-Cianferani S., Characterization of therapeutic antibodies and related products, *Anal. Chem.*, **85**, 7157–7166, <https://doi.org/10.1021/ac3032355> (2013)
4. Cernosek T., Dalphin M., Jain N., Richter N., Beard S., Wang J. and Wunderli P., Analytical quality by design as applied to the development of a SEC-HPLC platform procedure for the determination of monoclonal antibody purity without mobile phase additives, *J. Pharm. Biomed. Anal.*, **246**, 116220 (2024)
5. Chowdhury P.S. and Pastan I., Improving antibody affinity by mimicking somatic hypermutation *in vitro*, *Nat. Biotechnol.*, **17**, 568–572 (1999)
6. Dakshinamurthy P.K. and Gothandam K.M., Comprehensive analytical characterization of the proposed biosimilar trastuzumab, *Indian J. Pharm. Educ. Res.*, **56**, 438–447 (2022)
7. Dias K.E., Sourris K.C., Jha J.C., Jandeleit-Dahm K. and Wood B.R., Rapid, precise and clinically relevant quantification of urinary albumin and creatinine using a NanoDrop UV/Vis spectrophotometer, *Sensors*, **25**, 3307 (2025)
8. Fekete S., Beck A., Veuthey J.L. and Guilleme D., Theory and practice of size exclusion chromatography for the analysis of protein aggregates, *J. Pharm. Biomed. Anal.*, **101**, 161–173 (2014)
9. Green M.R. and Sambrook J., A guide to cloning the products of polymerase chain reactions, *Cold Spring Harb. Protoc.*, **2021**(9), pdb-top101345 (2021)
10. Gustafsson C., Govindarajan S. and Minshull J., Codon bias and heterologous protein expression, *Trends Biotechnol.*, **22**, 346–353 (2004)
11. Honorato R.V., Koukos P.I., Jiménez-García B., Tsaregorodtsev A., Verlato M., Giachetti A. and Bonvin A.M., Structural biology in the clouds: The WeNMR-EOSC ecosystem, *Front. Mol. Biosci.*, **8**, 729513 (2021)
12. Hudis C.A., Trastuzumab—Mechanism of action and use in clinical practice, *N. Engl. J. Med.*, **357**, 39–51, DOI: 10.1056/NEJMra043186 (2019)
13. Kayatani H., Ohashi K., Ninomiya K., Makimoto G., Nishii K., Higo H. and Kiura K., Beneficial effect of erlotinib and trastuzumab emtansine combination in lung tumors harboring EGFR mutations, *Biochem. Biophys. Res. Commun.*, **532**, 341–346 (2020)
14. Kunert R. and Reinhart D., Advances in recombinant antibody manufacturing, *Appl. Microbiol. Biotechnol.*, **100**, 3451–3461 (2016)
15. Leonard B., Sharma V. and Smider V., Co-expression of antibody Fab heavy and light chain genes in *E. coli*, *J. Immunol. Methods*, **317**, 56–63 (2006)
16. Li F., Vijayasankaran N., Shen A., Kiss R. and Amanullah A., Cell culture processes for monoclonal antibody production, *mAbs*, **2**, 466–479 (2010)
17. Loibl S. and Gianni L., HER2-positive breast cancer, *Lancet*, **389**, 2415–2429 (2017)
18. Lu Y., Zhou Q., Han Q., Wu P., Zhang L., Zhu L. and Zhang B., Inactivation of deubiquitinase CYLD enhances therapeutic antibody production in CHO cells, *Appl. Microbiol. Biotechnol.*, **102**, 6081–6093 (2018)
19. Marx N., Otte A.M., Leitner K., Sitepu R., Berger T., Schäpertöns V. and Borth N., Characterization of large transgene integrations in CHO cells, *Biotechnol. Prog.*, **41**, e3524 (2025)
20. Ménard S., Pupa S.M., Campiglio M. and Tagliabue E., Biologic and therapeutic role of HER2 in cancer, *Oncogene*, **22**, 6570–6578 (2003)

21. Pernas S. and Tolaney S.M., HER2-positive breast cancer: New therapeutic frontiers, *Ther. Adv. Med. Oncol.*, **11**, 1–16 (2019)
22. Pirzer T., Becher K.S., Rieker M., Meckel T., Mootz H.D. and Kolmar H., Generation of potent antiHER1/2 immunotoxins using split inteins, *ACS Chem. Biol.*, **13**, 2058–2066 (2018)
23. Roshani A., Mohammadi M., Bahadori T., Ahmadi Zare H., Judaki M.A., Mobini M. and Shokri F., Comparison of transient gene expression systems for anti-HER2 antibody production, *DARU J. Pharm. Sci.*, **31**, 221–231 (2023)
24. Rosano G.L. and Ceccarelli E.A., Recombinant protein expression in *Escherichia coli*: Advances and challenges, *Front. Microbiol.*, **5**, 172 (2014)
25. Slamon D.J., Clark G.M., Wong S.G., Levin W.J., Ullrich A. and McGuire W.L., Human breast cancer correlation with HER-2/neu amplification, *Science*, **235**, 177–182 (1987)
26. Soo B.P., Tay J., Ng S., Ho S.C., Yang Y. and Chao S.H., Correlation between recombinant protein expression and H3K4Me3 enhancer abundance, *Mol. Biotechnol.*, **59**, 315–322 (2017)
27. Sung H., Ferlay J., Siegel R.L., Laversanne M., Soerjomataram I., Jemal A. and Bray F., Global cancer statistics 2020, *CA Cancer J. Clin.*, **71**, 209–249 (2021)
28. Swain S.M., Baselga J., Kim S.B., Ro J., Semiglazov V., Campone M. and Cortés J., Pertuzumab, trastuzumab and docetaxel in HER2-positive breast cancer, *N. Engl. J. Med.*, **372**, 724–734 (2015)
29. UniProt Consortium, UniProt: The universal protein knowledgebase, *Nucleic Acids Res.*, **46**, 2699 (2018)
30. Van Zundert G.C.P., Rodrigues J.P.G.L.M., Trellet M., Schmitz C., Kastiris P.L., Karaca E. and Bonvin A.M.J.J., HADDOCK2.2 web server, *J. Mol. Biol.*, **428**, 720–725 (2016)
31. Walsh G., Biopharmaceutical benchmarks 2018, *Nat. Biotechnol.*, **36**, 1136–1145 (2018)
32. Waks A.G. and Winer E.P., Breast cancer treatment: A review, *JAMA*, **321**, 288–300 (2019)
33. Woo J., Kim J.B., Cho T., Yoo E.H., Moon B.I., Kwon H. and Lim W., Selenium inhibits growth of trastuzumab-resistant cancer cells, *PLoS One*, **16**, e0257298 (2021)
34. Yamashita-Kashima Y., Yoshimura Y., Fujimura T., Shu S., Yanagisawa M., Yoroze K., Furugaki K., Higuchi R., Shoda J. and Harada N., Molecular targeting of HER2-overexpressing biliary tract cancer cells with trastuzumab emtansine, an antibody-cytotoxic drug conjugate, *Cancer Chemother. Pharmacol.*, **83**, 659–671 (2019).

(Received 06th May 2026, accepted 09th June 2026)