

Extensive physicochemical and biological characterization of BP02, a trastuzumab biosimilar, and comparative assessment with Herceptin

Sanjeeva Metikala · Maheswara Rao · Venu Kumar Vemula · D. R. Yashwanth Kumar · Yash Patel · Babul Reddy · Naga Satish · Vuchuru Nagarjuna Reddy · Kambagiri Swamy Kayala · Pujitha Yarlagadda · Raghu K. Chitta · Srilalitha Kuruganti · Disha Dadke

Received: 27 April 2026 / Accepted: 8 September 2026
© The Author(s) 2026

Abstract

Trastuzumab is an IgG1 monoclonal antibody that targets the extracellular HER2 domain on HER2+ cancer cells. CuraTeQ biologics has developed a trastuzumab biosimilar, code named BP02, aimed at treating early and metastatic breast cancer and metastatic gastric cancer. This biosimilar has been thoroughly characterized through physicochemical and biological methods in accordance with established guidelines. The primary, secondary, and tertiary structures of were evaluated using analytical techniques, which indicated that BP02 shares structural similarities with Herceptin. Evaluations of size and charge variants revealed that BP02 possesses comparable levels of both high and low molecular weight species. Although minor differences in charge variants were observed in BP02 when compared to Herceptin, these differences were not deemed biologically significant. Furthermore, glycan heterogeneity and post-translational modifications were comparable in both BP02 and Herceptin. Biological studies were conducted to assess the inhibition of cancer cell proliferation, the primary mechanism of action of trastuzumab. Additional known mechanisms of trastuzumab, including antibody-dependent cellular cytotoxicity and antibody-dependent cellular phagocytosis, were also evaluated. BP02 demonstrated comparable biological activities across all assays and was found to be on par with Herceptin. Moreover, the binding activities and affinities of the HER2 receptor, FcRn, and other Fc receptors were assessed and determined to be similar. In summary, BP02 was found to exhibit similarities in both physicochemical and biological properties compared to Herceptin.

This Article in Press is shared early to give you faster access to new research. It is citable and carries a permanent DOI. The final edited version will replace it automatically.



Keywords Trastuzumab · Breast cancer · Herceptin · Biosimilars · Comparative assessment

Introduction

BP02, a trastuzumab biosimilar product developed by CuraTeQ Biologics, is a recombinant humanized IgG1 monoclonal antibody produced in Chinese Hamster Ovary (CHO) cells. It is approved in the European Union, UK and India by the name Dazublys[®] for early and metastatic breast cancer, metastatic gastric cancer and is a biosimilar to Roche Diagnostics' Herceptin[®].

Trastuzumab is directed against the human epidermal growth factor receptor 2 (HER2). It has played a critical role in advancing targeted therapy for solid tumors. Since its initial approval by the U.S. Food and Drug Administration (FDA) in 1998, trastuzumab has become the basis of treatment for HER2-positive breast cancer, which is overexpressed in 15–30% of overall breast cancer tumors¹, and later expanded to HER2-positive gastric and gastroesophageal junction cancers². Its clinical success has also driven the development of biosimilars to enhance accessibility and reduce treatment costs.

Trastuzumab is known to have multiple mechanisms of action. It binds with high affinity to the extracellular domain IV of the HER2 receptor, thereby inhibiting ligand-independent HER2 signaling and downstream oncogenic pathways such as PI3K/AKT and MAPK resulting in the inhibition of cellular proliferation^{3–7}. It also prevents HER2 homodimers and HER2 heterodimers with HER1/HER3/HER4 receptors⁸. Trastuzumab also promotes receptor internalization and degradation⁹, and reduces HER2 extracellular domain shedding¹⁰. It is also shown to mediate antibody dependent cell-cytotoxicity (ADCC)^{3–6,9} and antibody dependent cell-phagocytosis (ADCP)^{11,12} by recruiting immune effector cells to eliminate HER2-overexpressing tumor cells. It, however, lacks complement dependent cytotoxicity (CDC)^{12–14}.

The pharmacokinetics, safety, tolerability, and immunogenicity in Phase I, and efficacy and safety in Phase III clinical studies showed that BP02 is similar to EU-approved Herceptin[®] and US-licensed Herceptin[®]^{15,16}. This manuscript focusses on the comparative analytical assessment (CAA) of BP02 with EU-approved Herceptin[®] and US-licensed Herceptin[®] (hereafter referred to as EU-Herceptin and US-Herceptin or Herceptin). For the CAA, we employed a range of physicochemical and functional methods to characterize our trastuzumab biosimilar, BP02, and to compare it with the EU-Herceptin and the US-Herceptin. The physicochemical techniques were selected to thoroughly characterize trastuzumab, while the biological methods were utilized to evaluate the known mechanisms of action of trastuzumab. The combined evidence from physicochemical and biological methods ensured that no residual uncertainty remained with respect to biosimilarity of BP02 with Herceptin. All methods have been qualified according to established guidelines^{17–19} or have been demonstrated to be suitable for their intended purpose. The biosimilarity assessment was conducted across multiple campaigns utilizing batches of BP02 manufactured by CuraTeQ Biologics in India along with Roche's Herceptin batches sourced from the European Union and the United States of America to cover the product's shelf-life as comprehensively as possible. The demonstration of biosimilarity adhered to the existing EMA and FDA guidelines^{20–22}.

The clinical success of reference trastuzumab (Herceptin) has fundamentally transformed the therapeutic landscape for patients with human epidermal growth factor receptor 2 (HER2) positive breast and gastric malignancies. However, high treatment costs and financial constraints often restrict global access to this life-saving biologic, presenting a critical public health challenge. To address this disparity, BP02 was developed as a biosimilar to Herceptin with the aim of ensuring that this cancer treatment is as effective and efficacious as Herceptin, while also being affordable for patients in need. Because therapeutic monoclonal antibodies are highly complex, manufacturing variations can influence critical quality attributes (CQAs). Consequently, clinical adoption and therapeutic trust rest heavily on demonstrating structural and functional identity through head-to-head analytical benchmarking.

While regulatory approvals provide a baseline of safety and efficacy, publishing detailed, head-to-head physicochemical and in vitro biological characterisation data is a critical requirement for building trust among healthcare professionals, clinical investigators, and the public. Through this manuscript, we aim to make the product quality and in vitro biological activity data available to the public, patients, and healthcare professionals to make informed decisions when treating patients. Our findings indicate that BP02 exhibits a high degree of similarity to Herceptin concerning its physicochemical properties and its capacity to perform biological functions.

Results

A comprehensive panel of state-of-the-art analytical methodologies was employed to conduct a comparative assessment of BP02 against EU and US-Herceptin. Analytical similarity was demonstrated through rigorous characterization of the primary amino acid sequence, molecular and charge heterogeneity, glycosylation profiles, post-translational modifications, higher-order structural attributes, and relevant biological activities.

Quality attributes were chosen based on the available literature and known mechanisms of action of trastuzumab as per ICH Q8 (R2)²³. For this evaluation, BP02 drug product (DP) batches manufactured from multiple drug substance (DS) lots were included. All comparative analyses were performed in a direct, head-to-head manner utilizing the full suite of analytical techniques to ensure a robust and comprehensive similarity assessment. For similarity assessment, $\mu \pm 3\sigma$ (mean of the EU-Herceptin or US-Herceptin ± 3 *standard deviation) approach was taken for the quality attributes which are amenable to statistics.

Physicochemical characterization

Primary structure

The primary structure of trastuzumab was analyzed using multiple orthogonal methods. The peptide chromatograms of trypsin digested BP02, and Herceptin products showed similar peaks. No additional peaks were observed in BP02 compared to US and EU-Herceptin (Fig. 1). The amino acid composition of BP02 and Herceptin was also identical, and the sequence coverage was 100% (data not shown). Furthermore, the intact, reduced and




Fig. 1 Peptide mapping. Chromatograms showing tryptic peptide map profiles of BP02 and Herceptin by RP-HPLC.

deglycosylated molecular mass of BP02 is similar to Herceptin and match with the theoretical masses (Table 1; Fig. 2). The extinction coefficient of BP02 was also found to be within the quality ranges of Herceptin (Table 1).

Higher order structure

The secondary and tertiary structures were assessed using multiple spectrophotometric techniques, including far-UV circular dichroism (CD), near-UV CD, intrinsic fluorescence (FLR), nuclear magnetic resonance (NMR), size-exclusion chromatography with multi-angle light scattering (SEC-MALS) and Fourier transform infrared (FTIR) spectroscopy. Further, the conformational stability (thermal properties and structural-phase transitions) of BP02 and Herceptin were assessed by differential scanning calorimetry (DSC).

The analysis by far-UV CD and FTIR showed that the secondary structure profiles for BP02 and Herceptin are comparable, and the major structural feature is beta-sheet, which is consistent with the known IgG1 mAb structure (Table 1; Fig. 3A, B). The tertiary structure of BP02 and Herceptin was assessed using near-UV CD and FLR. The profiles are visually comparable, and the emission maxima data obtained after excitation at 280 nm and 295 nm indicate that BP02 and Herceptin have comparable tertiary structure. (Table 1; Fig. 3C, D).

The differential energy requirement for phase transitions and the unfolding of sample was measured using DSC. The thermal unfolding transition midpoints (Tm1 and Tm2) were found to be comparable, indicating similar thermal stability and in turn a similar higher order structure and folding of BP02 and Herceptin (Table 1; Fig. 4A).

The light scatter profiles, molar mass (kDa), hydrodynamic radii (Rh) and polydispersity of BP02 and Herceptin were assessed using SEC-MALS. The polydispersity index indicated that BP02 and Herceptin have a narrow molecular weight distribution which confirms similarity of BP02 with innovator HMW/aggregate species with molar mass distribution were also comparable, indicating that the mean monomer molecular weight and hydrodynamic radius of BP02 and Herceptin are similar confirming comparable oligomeric/higher order structure (Table 1; Fig. 4B).

The higher order structure was also studied using 1D ^1H NMR. The spectra exhibited characteristic fingerprints of methyl/methylene groups (<0.5 ppm) from a folded hydrophobic core and backbone amide resonances (>8.5 ppm) associated with ordered secondary structure and aromatic side chains (Fig. 5). These spectral regions, when visually compared for BP02 and EU-approved Herceptin, indicate highly similar overall folding (Fig. 1; Table 1).

The disulphide bond assignment was studied using non reduced RP-UPLC-LC-ESI-MS/MS. The overlay of the chromatogram profiles demonstrated 8 disulphide bonded linked peptides in BP02 and Herceptin. (Fig. 6).

Overall, the higher order structures (secondary, tertiary) structures of BP02 are similar to Herceptin.

Molecular heterogeneity

Molecular heterogeneity of trastuzumab was evaluated using multiple analytical techniques using BP02 drug product batches against EU-Herceptin and US-Herceptin batches. SE-HPLC was employed to assess purity, AUC was used to quantify high molecular weight species (HMWs), CE-SDS was used to characterize low molecular weight species (LMWs), and CE-SDS under reduced conditions was used for non-glycosylated heavy chain species (NGHC) and light-chain and heavy-chain (LC+HC). The SE-HPLC profiles showed distinct monomer peaks with the %HMWs comparable between BP02 and Herceptin (Table 1; Fig. 7). AUC analysis further confirmed that the HMW species were comparable between BP02 and Herceptin (Table 1; Fig. 8). Additionally, CE-SDS profiles showed that the LMWs were comparable between BP02 and Herceptin. (Table 1; Fig. 9A – C). The level of LC+HC content was also similar across BP02 and Herceptin and NGHC content was comparatively lower in BP02 (Table 1; Fig. 9D – F). Overall, the purity, HMW and LMW profiles of BP02 are similar to Herceptin.

Table 1 Physicochemical characterization data summary.

Product Quality Attribute	Analytical Methodology	BP02 Mean, SD (min, max), <i>N</i>	EU-Herceptin Mean, SD (min, max), <i>N</i>	US-Herceptin Mean, SD (min, max), <i>N</i>
Product molecular mass, intact	LC-ESI-MS	148055.0, (148052.8, 148058.1), 12	148054.9, (148053.3, 148057.0), 12	148056.2, (148053.6, 148058.6), 10
Product molecular mass, reduced, heavy chain	LC-ESI-MS	50601.1, (50600.5, 50602.2), 12	50601.1, (50600.7, 50602.2), 12	50600.7, (50600.1, 50601.3), 10
Product molecular mass, reduced, light chain	LC-ESI-MS	23442.4, (23442.3, 23442.7), 12	23442.5, (23442.4, 23442.6), 12	23442.5, (23442.4, 23442.6), 10
Product molecular mass, deglycosylated intact	LC-ESI-MS	145164.7, (145161.4, 145166.3) 12	145165.3, (145163.0, 145167.2) 12	145164.6, (145161.6, 145168.3) 10
Extinction Coefficient	UV-Spectroscopy	1.47, (1.47, 1.47) 03	1.47, (1.46, 1.47) 03	1.48, (1.47, 1.48) 03
Secondary Structure	CD maxima by Far-UV CD	202.0, (201.7, 202.4), 12	202.1, (201.5, 202.6), 12	202.0, (201.8, 202.2), 10
	CD minima by Far-UV CD	217.3, (216.7, 217.7), 12	217.3, (216.9, 217.7), 12	217.4, (216.7, 217.9), 10
	Wavenumber/ %T dip (cm ⁻¹) by FTIR	1637.0, (1637.3, 1637.3), 12	1637.5, (1637.3, 1639.2), 12	1637.5, (1637.3, 1639.2), 10
	CD maxima by Near-UV CD	296.4, (295.5, 297.2), 12	296.4, (295.8, 297.1), 12	296.7, (296.0, 297.5), 10
Tertiary Structure	Emission Maxima (nm) (Ex 280 nm) by intrinsic fluorescence (FLR)	329, (329, 331), 12	330, (329, 330), 12	330, (328, 331), 10
	Emission Maxima (nm) (Ex 295 nm) by intrinsic fluorescence (FLR)	330, (329, 333), 12	330, (329, 331), 12	330, (329, 331), 10
Melting Temperature (T _m)	T _m 1 °C (peak 1) by DSC	82.6, (82.0, 83.8), 12	82.4, (82.0, 83.8), 12	82.8, (82.0, 83.8), 10
	T _m 2 °C (peak 2) by DSC	71.6, (70.6, 73.4), 12	71.0, (70.6, 73.1), 12	71.6, (70.6, 72.9), 10
Aggregates/ HMWs	SEC-MALS	332.3, 37.6 (300.4, 426.0), 12	335.4, 30.9 (310.8, 407.8), 12	330.6, 35.1 (291.5, 393.4), 10
Free Thiol	Ellman's Assay	0.6, 0.1 (0.5, 0.7), 10	0.5, 0.1 (0.4, 0.5), 4	0.5, 0.0 (0.4, 0.5), 10
Monomer	SE-HPLC	99.0, 0.48 (98.3, 99.6) 12	99.2, 0.08 (99.1, 99.3) 12	99.3, 0.59 (97.7, 99.7) 10
HMWs		0.90, 0.44 (0.38, 1.60), 12	0.62, 0.05 (0.50, 0.67), 12	0.63, 0.54 (0.29, 2.10), 10
LMWs		0.09, 0.09 (0.02, 0.27), 12	0.17, 0.08 (0.02, 0.26), 12	0.08, 0.08 (0.02, 0.22), 10
Aggregates	SV-AUC	1.4, 0.2 (1.2, 1.6), 03	1.4, 0.8 (0.7, 2.2), 03	0.9, 0.2 (0.7, 1.1), 03
Intact IgG % area	CE-SDS (NR)	97.1, 0.83 (95.5, 97.9), 12	97.2, 0.48 (96.8, 98.0), 12	97.4, 0.48 (96.8, 98.1), 10
LMWs % area		2.7, 0.99 (1.7, 4.6), 12	2.8, 0.52 (1.9, 3.2), 12	2.5, 0.54 (1.9, 3.2), 10
Purity - LC+HC % area	CE-SDS (R)	98.8, 0.18 (98.4, 99.0), 12	98.7, 0.11 (98.6, 98.9), 12	98.7, 0.18 (98.5, 99.1), 10
Purity - NGHC % area		0.51, 0.16 (0.35, 0.80), 12	0.67, 0.03 (0.64, 0.73), 12	0.67, 0.06 (0.52, 0.79), 10
Without CpB digestion - Acidic peak % area	CEX-HPLC	31.7, 2.3 (27.3, 34.6), 12	28.4, 0.5 (27.7, 28.9), 12	27.9, 1.1 (26.2, 29.3), 10
Without CpB digestion - Main peak % area		54.1, 2.6 (50.0, 57.5), 12	57.9, 0.6 (56.9, 58.9), 12	58.5, 1.8 (56.0, 61.4), 10
Without CpB digestion - Basic peak % area		14.2, 1.3 (12.5, 17.0), 12	13.7, 0.3 (13.4, 14.4), 12	13.6, 0.8 (12.3, 14.7), 10
With CpB digestion - Acidic peak % area		34.0, 2.5 (30.1, 37.3) 12	30.8, 0.6 (29.7, 31.5), 12	30.5, 2.0 (27.6, 33.6), 10
With CpB digestion - Main peak % area		54.1, 2.6 (50.0, 57.9), 12	57.0, 0.5 (56.3, 58.2), 12	57.6, 2.2 (54.1, 60.9), 10
With CpB digestion - Basic peak % area		11.6, 1.1 (10.3, 13.5), 12	12.2, 0.5 (11.4, 12.8), 12	11.9, 0.6 (11.1, 13.0), 10

Table 1 (continued)

Product Quality Attribute	Analytical Methodology	BP02 Mean, SD (min, max), <i>N</i>	EU-Herceptin Mean, SD (min, max), <i>N</i>	US-Herceptin Mean, SD (min, max), <i>N</i>
Isoelectric point	icIEF	8.9, (8.8, 8.9), 12	8.9, (8.8, 8.9), 10	8.9, (8.8, 8.9), 12
Total afucose % (G0+G1+G1'+G2)	HILIC	7.0, 0.9 (6.0, 8.7), 12	6.8, 0.6 (5.1, 7.7), 12	6.7, 0.3 (6.1, 7.1) 10
Total mannose % (M5+M6)		2.1, 0.5 (1.6, 3.2), 12	3.0, 0.5 (2.5, 4.2), 12	3.3, 0.8 (2.2, 4.5), 10
Total fucose % (G0F+G1F+G1F'+G2F)		88.2, 0.7 (86.6, 89.1), 12	86.4, 1.2 (83.1, 88.3), 12	86.0, 0.9 (84.5, 87.1), 10
NANA/NeuAc (Moles NeuAc per mole of protein)	RP-UPLC	0.056, 0.006 (0.047, 0.066), 10	0.046, 0.004 (0.041, 0.049), 6	0.045, 0.005 (0.039, 0.049), 6
NGNA/NeuGc (Moles NeuGc per mole of protein)		0.001, 0.000 (0.001, 0.002), 10	0.004, 0.001 (0.002, 0.005), 6	0.004, 0.002 (0.002, 0.006), 6
Total Sialylation (NANA+NGNA) in moles per mole of protein		0.057, 0.006 (0.048, 0.067), 10	0.049, 0.004 (0.044, 0.054), 6	0.049, 0.006 (0.041, 0.055), 6
Fucose monosaccharide composition	GC-MS	5.9, 1.0 (5.0, 6.9), 3	5.5, 0.7 (4.8, 6.1), 3	5.2, 0.5 (4.7, 5.7), 3
Mannose monosaccharide composition		20.0, 1.0 (19.0, 21.0), 3	19.8, 1.6 (18.2, 21.4), 3	20.8, 1.6 (19.0, 21.9), 3
Galactose monosaccharide composition		2.7, 0.7 (2.5, 2.9), 3	3.6, 0.2 (3.4, 3.8), 3	3.6, 0.1 (3.6, 3.7), 3
GLcNAc monosaccharide composition		16.1, 1.6 (14.4, 17.5), 3	15.2, 2.6 (12.3, 17.1), 3	16.4, 0.7 (15.8, 17.2), 3




Fig. 2 Molecular Mass. Deconvoluted profiles for A) intact mass, C), E) reduced mass (heavy and light chain), and G) deglycosylated intact mass.

Charge heterogeneity

Trastuzumab was assessed using icIEF to determine the isoelectric point and CEX-HPLC (with and without CpB treatment) to assess charge variants. The isoelectric points were between 8.8 and 8.9 across BP02 and Herceptin (Table 1). Charge variants were evaluated using CEX-HPLC technique with and without CpB digestion. The percentages of main peak, acidic peaks and basic peaks were similar with minor differences (Table 1; Fig. 10) in BP02 drug product batches. These minor differences do not have any impact on biological activity.

Glycan heterogeneity

Glycan species of trastuzumab was assessed by HILIC analysis using 2-AB labelling. The chromatographic profiles are shown in Fig. 11. Glycan groups were categorized into % afucosylation, % fucosylation and % mannose as these groups are relevant for Fc-effector functions. % fucosylation and % mannose fall within the Herceptin




Fig. 3 Secondary and Tertiary Structure. (A) Far-UV CD profiles of BP02 and Herceptin. (B) FTIR Profiles of BP02 and Herceptin. (C) Near UV-CD profiles of BP02 and Herceptin. (D) Intrinsic fluorescence profiles of BP02 and Herceptin.




Fig. 4 Higher Order Structure. (A) DSC profiles of BP02 and Herceptin showing major and minor peaks. (B) SEC-MALS profiles of BP02 and Herceptin showing monomer and HMW peaks.

quality ranges. Minor difference of 0.1% was observed in % afucosylation level of 1 drug product batch out of 12 BP02 batches which is negligible. The overall glycan levels were comparable (Table 1; Fig. 11B - D). The ADCC activities confirmed that these minor differences did not affect the ADCC activity. The mannose levels from monosaccharide analysis using GC-MS showed that BP02 mannose levels were within the Herceptin quality ranges (data not shown). Two major Sialic acids, NANA/Neu5Ac and NGNA/NeuGc, were measured. Although NANA/Neu5Ac levels appear to be slightly higher in two BP02 drug substance batches, total sialylation levels of BP02 were similar to Herceptin (Table 1; Fig. 11E - G). Difference in sialylation can affect FcRn binding and ADCC activity²⁴. This minor difference in sialic acid levels did not have any effect on the FcRn binding affinities or ADCC activity (Figs. 13D, E, G and H and 14A and B).




Fig. 5 NMR Spectra. Zoomed in spectra exhibited characteristic fingerprints of (A) backbone amide resonances (>8.5 ppm) associated with ordered secondary structure and aromatic side chains, and (B) methyl/methylene groups (<0.5 ppm) from a folded hydrophobic core.




Fig. 6 Disulphide Mapping Profiles. Disulphide bond assignment of BP02 and Herceptin by ESI LC-MS/MS (Non-reduced RP-UPLC).

Post-translational modifications

Post-translational modifications (PTMs) were determined by RP-UPLC tryptic peptide mapping method for deamidation and oxidation. Peptide masses of the expected modifications such as oxidation, deamidation, N-terminal pyroglutamate, and C-terminal lysine variants were derived from the MS and MS/MS spectra and were compared to the theoretical mass (Fig. 12). The amino acid sequences of the peptides were compared between BP02 and Herceptin. All PTMs of BP02 fall within the quality ranges of Herceptin with minor differences that do not affect the biological activity (Table 1).




Fig. 7 Size Heterogeneity by SE HPLC. (A) SE HPLC profiles of BP02 and Herceptin and (B), (C) the distribution of monomer and HMW % areas. Whiskers represent minimum and maximum values. Solid brown lines on y-axis represents 3σ of EU-Herceptin and dashed blue lines represent 3σ of US-Herceptin.

Functional characterization

Inhibition of cellular proliferation

Trastuzumab binds to the subunit 4 of the surface HER2 receptor's extra-cellular domain thereby inhibiting HER2-HER2 homodimerization. This inhibition results in reduced activation of PI3K/AKT signalling pathway ultimately reducing the cellular proliferation^{3,8}. To compare the inhibition of cellular proliferation, BT474 cells with high HER2 surface expression were used^{25,26}. Upon treating the cells with BP02, EU-Herceptin and US-Herceptin, the cellular proliferation was inhibited. The relative potencies of BP02 fall within the Herceptin quality range (Table 2; Fig. 13A). The results demonstrate that BP02's potency to inhibit proliferation of cancer cells is similar to the reference medicinal product.

HER2 binding

Trastuzumab exerts its function by binding to the HER2 receptor^{3,6} on the surface of the cancer cells. The binding of BP02, EU-Herceptin and US-Herceptin to HER2 was evaluated using BT474 cell-based ELISA. The relative binding activities of BP02 fall within the Herceptin quality range (Table 2; Fig. 13B). Real-time binding affinities of BP02, EU-Herceptin and US-Herceptin to HER2 were evaluated using surface plasmon resonance (SPR). The relative binding affinities of BP02 fall within the Herceptin quality range (Table 2; Fig. 13C). These results demonstrate that BP02's HER2 binding is highly similar to the reference medicinal product.




Fig. 8 Size Heterogeneity by SV-AUC. (A) SV-AUC profiles of BP02 and Herceptin and (B), (C) the distribution of monomer and HMW % areas. Whiskers represent minimum and maximum values. Solid brown lines on y-axis represents 3σ of EU-Herceptin and dashed blue lines represent 3σ of US-Herceptin.

Antibody dependent cell-cytotoxicity (ADCC)

Trastuzumab is known to exhibit antibody dependent cell-cytotoxicity (ADCC) mediated by NK cells through Fc receptors. The Fc portion of trastuzumab binds to both high and low affinity FcγRIIIA receptors (V176 and F176)^{6,9}.

The ADCC activity of BP02, EU-Herceptin and US-Herceptin to HER2 was evaluated using BT474 cells as target cells, and luciferase reporter gene and FcγRIIIA (V176 or F176) receptor expressing Jurkat cells as effector cells. The ADCC activities, for both FcγRIIIA receptor variants, of BP02 fall within the Herceptin quality range (Table 2; Fig. 13D, E). The ADCC activity was also evaluated using CHO cells expressing HER2 receptor as target cells and PBMCs as effector cells. The ADCC activities, for both FcγRIIIA receptor variants, of BP02 fall within the Herceptin quality range (Table 2; Fig. 13G, H). These results show that BP02 is similar to Herceptin in inducing ADCC activity.

Antibody dependent cell-phagocytosis (ADCP)

Trastuzumab is known to exhibit ADCP mediated by macrophage activation through Fc receptors. The Fc portion of trastuzumab binds to FcγRIIA receptor^{11,12}.

The ADCP activity of BP02, EU-Herceptin and US-Herceptin to HER2 was evaluated using BT474 cells as target cells and luciferase reporter gene and FcγRIIA receptor expressing Jurkat cells as effector cells. The ADCP




Fig. 9 Size Heterogeneity by CE SDS. (A) Non-reduced CE SDS profiles of BP02 and Herceptin and (B), (C) the distribution of intact IgG and LMW % areas. D) Reduced CE SDS profiles of BP02 and Herceptin and (E), (F) the distribution of LC+HC % area and NGHC % area. Whiskers represent minimum and maximum values. Solid brown lines on y-axis represents 3σ of EU-Herceptin and dashed blue lines represent 3σ of US-Herceptin.




Fig. 10 Charge Heterogeneity by CEX. (A) Charge variant profiles of BP02 and Herceptin and (B) - (D) the distribution of main acidic and basic peak % areas using box plots. E) Charge variant profiles of BP02 and Herceptin and (F) - (H) the distribution of LC+HC % area and NGHC % area using box plots obtained by CpB digested samples. Whiskers represent minimum and maximum values. Solid brown lines on y-axis represents 3σ of EU-Herceptin and dashed blue lines represent 3σ of US-Herceptin.

activities of BP02 fall within the Herceptin quality range (Table 2; Fig. 13F). The results show that BP02 exerts ADCP activity similar to Herceptin.

Complement-dependent cytotoxicity (CDC)

In CDC assay, the Fab portion of a monoclonal antibody binds to the antigen on the target cell. The complement factor, C1q, binds to the Fc portion of this monoclonal antibody causing opsonization of the target cell. This triggers the complement cascade, resulting in the formation of a membrane attack complex (MAC) on the surface of target cells, further resulting in a classical pathway of complement activation ensuring target cell lysis. Trastuzumab does not exhibit CDC activity^{12–14}.




Fig. 11 Glycan Heterogeneity. (A) N-glycosylation profiles of BP02 and Herceptin and (B) - (D) the distribution of % afucosylation, % fucosylation and % total mannose using box plots. E) – G) distribution of NANA/NeuAc, NGNA/NeuGc and total sialylation / NANA+NGNA using box plots. Whiskers represent minimum and maximum values. Solid brown lines on y-axis represents 3σ of EU-Herceptin and dashed blue lines represent 3σ of US-Herceptin.

The absence of CDC activity of BP02, EU-Herceptin and US-Herceptin to HER2 was evaluated using baby rabbit complement (BRC). Rituximab, along with CD20 expressing WIL2-S cells, was used as a positive assay control, whereas trastuzumab, along with HER2 expressing BT474 cells, was used to test for CDC activity. At increasing Rituximab doses, CDC activities ranged from 1.11 to 2.53-fold compared to the BRC normalized control taken as 1. CDC activities for BP02, EU-Herceptin and US-Herceptin were not observed at any of the doses of trastuzumab (Table 2; Fig. 13I).

FcRn binding, Fc receptors binding and C1q binding

Relative binding affinities of neonatal FcRn, Fc receptors and C1q to BP02, EU-Herceptin and US-Herceptin were assessed using surface plasmon resonance (SPR) and bio-layer interferometry (BLI). FcRn binding of an IgG protects it from degradation and increases its half-life. FcγRIIA and FcγRIIIA are low affinity receptors involved in ADCP and ADCC activities. FcγRIIB is an inhibitory receptor. Even though trastuzumab does not exhibit CDC activity, it still binds to C1q. Therefore, this binding was assessed^{13,27,28}.

The BP02 relative binding affinities of FcRn K_{D1} and K_{D2} fall within the Herceptin quality ranges (Table 2; Fig. 14A, B). The BP02 relative binding affinities of FcγRI fall within the Herceptin quality range (Table 2; Fig. 14C). The BP02 relative binding affinities of FcγRIIA H167 and R167 fall within the Herceptin quality range (Table 2; Fig. 14D, E). The relative binding affinities of FcγRIIB were also evaluated. All BP02 batches, except two drug product batches, fall within the Herceptin quality range (Table 2; Fig. 14F). FcγRIIB is considered an inhibitory receptor^{27,28}. However, the ADCP activity of this batch was similar to Herceptin suggesting that the difference observed is negligible.




Fig. 12 Identity. Tryptic peptide map profiles (UV 214) by (A) RP UPLC and (B) LC-MS/MS.

The relative binding affinities of all BP02 batches for FcγRIIIA V176 and 13 out of 14 batches analyzed for FcγRIIIA F176 fall within the Herceptin ranges (Table 2; Fig. 14G, H). One drug product batch is outside the Herceptin quality ranges. FcγRIIIA F176 is considered to have weaker binding affinity and forms a less stable complex compared to V176^{29,30}. Furthermore, the ADCC activities in our experiments using RGA and PBMC based assays showed that the potencies of all BP02 batches are similar to Herceptin (Fig. 13D, E, G, H). All batches of BP02 for FcγRIIIB relative binding affinities fall within the Herceptin quality range (Table 2; Fig. 14I). Lastly, BP02 relative binding affinities of C1q fall within the Herceptin quality range (Table 2; Fig. 14J).

The results demonstrate that BP02 is highly similar to the reference medicinal product in its

binding affinities to FcRn, Fc receptors and C1q with minor differences that did not show significant effect in biological activities.

Table 2 Biological characterization data summary.

Product Quality Attribute	Analytical Methodology	Reporting parameter	BP02 Mean, SD (min, max), <i>N</i>	EU-Herceptin Mean, SD (min, max), <i>N</i>	US-Herceptin Mean, SD (min, max), <i>N</i>
Inhibition of cellular proliferation	Potency Estimation by BT474 Proliferation Inhibition Assay	% relative potency	102, 7 (90, 118), 16	101, 7 (88, 117), 19	104, 7 (93, 114), 10
HER2 binding	Target Binding Activity- HER2 binding ELISA	% relative binding activity	101, 8 (88, 119), 12	100, 10 (88, 126), 13	97, 9 (84, 112), 10
	HER2 Binding by SPR	% relative binding affinity	97, 7 (85, 110), 14	100, 5 (87, 105), 13	99, 3 (92, 104), 10
ADCC	ADCC V176 activity by RGA	% relative potency	103, 8 (86, 114), 14	102, 10 (88, 124), 13	103, 5 (97, 111), 10
	ADCC F176 activity by RGA	% relative potency	105, 11 (80, 120), 14	104, 11 (90, 127), 13	103, 10 (86, 116), 10
	ADCC VV176 activity by PBMC	% relative potency	105, 12 (84, 121), 7	96, 15 (68, 108), 7	86, 16 (59, 105), 7
	ADCC FF176 activity by PBMC	% relative potency	102, 12 (88, 114), 7	96, 10 (83, 108), 7	88, 13 (60, 98), 7
ADCP	ADCP activity by RGA	% relative potency	101, 9 (88, 116), 14	112, 5 (104, 109), 13	103, 9 (91, 119), 10
CDC	Negative CDC activity	fold change	NA	NA	NA
FcRn binding	FcRn binding by SPR	% relative binding affinity of K _{D1}	98, 2 (94, 103), 14	90, 3 (85, 96), 13	91, 4 (85, 98), 10
		% relative binding affinity of K _{D2}	101, 8 (89, 118), 14	97, 4 (91, 103), 13	101, 7 (87, 109), 10
FcγRI binding	FcγRI binding by SPR	% relative binding affinity	101, 6 (91, 110), 12	96, 3 (91, 102), 13	98, 4 (89, 104), 10
FcγRIIA binding	FcγRIIA H167 binding by BLI	% relative binding affinity	100, 14 (78, 117), 12	98, 9 (81, 109), 13	89, 10 (78, 101), 10
	FcγRIIA R167 binding by BLI	% relative binding affinity	95, 15 (73, 121), 12	88, 8 (79, 101), 13	87, 11 (70, 107), 10
FcγRIIB binding	FcγRIIB binding by BLI	% relative binding affinity	103, 12 (80, 125), 12	99, 6 (88, 109), 13	91, 9 (75, 102), 10
FcγRIIIA binding	FcγRIIIA V176 binding by SPR	% relative binding affinity	105, 9 (93, 123), 14	99, 8 (86, 115), 13	104, 9 (90, 121), 10
	FcγRIIIA F176 binding by SPR	% relative binding affinity	105, 6 (94, 120), 14	102, 5 (95, 111), 13	102, 4 (97, 111), 10
FcγRIIIB binding	FcγRIIIB binding by BLI	% relative binding affinity	105, 8 (91, 115), 12	102, 7 (89, 116), 13	98, 11 (81, 110), 10
C1q binding	C1q binding by BLI	% relative binding affinity	103, 4 (95, 108), 12	103, 3 (100, 109), 13	100, 4 (95, 105), 10

Discussion

A comprehensive analytical similarity assessment was conducted to compare BP02, a trastuzumab biosimilar, with EU-approved and US-licensed Herceptin reference products. Multiple structural and functional attributes were evaluated using an extensive set of orthogonal techniques to minimize uncertainty and ensure robust conclusions regarding biosimilarity. All studies adhered to relevant regulatory guidelines, including ICH Q2 (R1) and USP Chaps. 1032, 1033^{17–19}, and employed qualified methods.

Quality ranges for critical attributes were established based on the mean ± 3 standard deviations of the reference product data, providing scientifically justified acceptance criteria. Across all evaluated parameters, BP02 consistently fell within these ranges.




Fig. 13 Functional characterization of BP02, EU-Herceptin and US-Herceptin. Box plots showing relative potencies of BP02, EU-Herceptin and US-Herceptin for (A) inhibition of cellular proliferation, (B) cell-based HER2 binding ELISA, (C) HER2 binding by SPR, (D, E) ADCC V176 and F176 activity by RGA, (F) ADCP activity by RGA, (G, H) ADCC VV176 and FF176 activity using PBMC based assay and (I) Bar graphs showing CDC activity. Whiskers represent minimum and maximum values. Solid brown lines on y-axis represents 3σ of EU-Herceptin and dashed blue lines represent 3σ of US-Herceptin. Black dotted line in I represent the fold-change obtained using the lowest concentration of the positive control.

Structural characterization confirmed that the primary sequence, amino acid composition, molecular mass, and extinction coefficient of BP02 were identical to those of the reference products. Higher-order structural analyses, including secondary and tertiary structure assessments and thermal stability studies, revealed no significant differences. NMR-based evaluations further supported structural similarity. Purity profiles, including main peak, high molecular weight species (HMWs), and low molecular weight species (LMWs), were comparable among all products using orthogonal techniques. Charge variant analysis showed minor differences in acidic, main, and basic peaks; however, these variations were not functionally relevant.

Given the role of Fc-mediated effector functions in trastuzumab's mechanism of action, glycan heterogeneity was examined. BP02 demonstrated fucosylation, mannose, and sialylation levels similar to Herceptin. No differences in post-translational modifications were detected, supporting consistency in glycan-dependent activities.

Functional similarity was confirmed across all critical attributes. HER2 binding was demonstrated using ELISA and SPR, and inhibition of BT-474 breast cancer cell proliferation showed highly similar antiproliferative activity. Fc-effector functions were assessed through interactions with FcγRIIA and FcγRIIIA receptors using BLI and SPR, confirming comparable antibody dependent cell-cytotoxicity (ADCC) and antibody dependent cell-phagocytosis (ADCP) activities in reporter gene and PBMC-based assays. Complement-dependent cytotoxicity (CDC) was absent, consistent with trastuzumab's known mechanism of action^{3,4,6,7,9}. Finally, FcRn binding studies indicated similar affinities across all products, supporting equivalent half-life predictions.

Although BP02, EU-Herceptin and US-Herceptin are recombinant proteins produced from CHO cells, heterogeneity is inherent to proteins produced in living organisms. Therefore, minor differences in PTMs, glycan profile and charge variants are expected. However, for the sponsor's product to be considered a biosimilar to the innovator product, it is expected that these differences do not affect the biological effect of the product. Minor differences were observed in BP02 compared to EU-Herceptin or US-Herceptin in terms of charge variants, sialic acid content, FcγRIIB, FcγRIIIA (F176) binding affinities. However, the biological activities were within the




Fig. 14 Binding of FcRn, Fc receptors and C1q to BP02, EU-Herceptin and US-Herceptin. Box plots showing relative binding affinities of BP02, EU-Herceptin and US-Herceptin for (A, B) FcRn, (C) FcγRI, (D, E) FcγRIIA H167 and R167, (F) FcγRIIB, (G, H) FcγRIIIA V176 and F176, (I) FcγRIIIB and (J) C1q. Whiskers represent minimum and maximum values. Solid brown lines on y-axis represents 3σ of EU-Herceptin and dashed blue lines represent 3σ of US-Herceptin.

EU-Herceptin and US-Herceptin quality ranges. Therefore, the results from this comparative assessment showed that despite minor differences, the biological activity was not affected thereby supporting similarity.

Collectively, these findings demonstrate that BP02 is highly similar to EU-approved and US-licensed Herceptin® in terms of structural integrity, glycan profile, and functional activity, thereby confirming its biosimilarity.

Conclusion

BP02 demonstrates physicochemical and biological sameness with EU-Herceptin and US-Herceptin. Comprehensive analytical assessments confirm that BP02 meets all predefined quality ranges and exhibits comparable HER2 binding, antiproliferative activity, and Fc-mediated effector functions. These findings collectively establish BP02 as a biosimilar to Herceptin.

Methods

Intact and reduced mass analysis

For intact mass analysis, test samples were separated on a MAbPac™ RP column using Waters ACQUITY UPLC H-Class system connected to Orbitrap Q Exactive Plus mass spectrometer (Thermo Fisher Scientific). For reduced mass analysis, test samples were denatured (7 M GuHCl, and 0.1 M tris, pH 7.0) and reduced with

5 mM DTT. Q Exactive Plus Orbitrap mass spec settings: Full MS in positive mode, spray voltage 4.2 kV, capillary temperature 350 °C (intact) and 320 °C (reduced). MS spectra were collected in range of 1000–5000 m/z (intact) and 500–4000 m/z (reduced) at a resolution of 17,500 and the AGC target in MS data acquisition was set to 1×10^6 at 100 ms injection time. Formic acid and trifluoroacetic acid in water was used as mobile phase A and acetonitrile, formic acid and trifluoroacetic acid was used as mobile phase B. Thermo BioPharma Finder (V 1.0) software was used to process and interpret the collected LC-MS data.

Peptide mapping analysis

Test samples were denatured (7 M Urea, 2 mM EDTA, 0.1 M Tris, pH 8.0), disulfide bonds (-S-S-) were reduced with 50 mM DTT and alkylated with 125 mM iodoacetic acid. Samples were then desalted using PD-10 desalting columns and digested overnight with trypsin at 1:30 ratio. Digested and desalted peptide sample was separated on ACQUITY UPLC-BEH column (130 Å, 1.7 µm, 2.1 mm X 150 mm) with C18 stationary phase, using ACQUITY UPLC H-Class system hyphenated to Orbitrap Q Exactive Plus mass spectrometer (Thermo Fisher Scientific).

The Q Exactive Plus benchtop Orbitrap mass spec settings: Tryptic peptides were analyzed in data dependent acquisition mode (dd-MS2 Top12) in positive mode. MS1 spectra were collected in the scan range of 150–2200 m/z at 70,000 resolution. Data dependent MS2 experiments for top 12 ions were carried out using 23, 25, 27 kV stepped (N) normalized collision energy at 17,500 resolution, 100 ms injection time and isolation window (m/z = 2). Water containing trifluoroacetic acid and acetonitrile containing trifluoroacetic acid were used as mobile phase A and B, respectively. The mass spectra raw data was analyzed using Thermo Xcalibur, Qual Browser (v 4.1.31.9) and BioPharma Finder 1.0.

Disulfide mapping analysis

Protein samples were denatured (7 M Urea, 2 mM EDTA, 0.1 M Tris, pH 8.0) and alkylated (125 mM iodoacetic acid) to cap any free cysteine residues. Samples were then desalted using PD-10 desalting columns with 2 M Urea and 0.5 M Tris, pH 7.8 and subsequently digested overnight with trypsin (1:30). MS instrument method parameters used for peptide mapping were used for disulfide mapping as well. The mass spectra raw data was analysed using Thermo Xcalibur, Qual Browser (V 4.1.31.9).

Extinction coefficient

The molar extinction coefficient ($\epsilon_{(280)}$) was determined using the Edelhoch method, which estimates ϵ based on the absorbance contributions of aromatic residues under fully denaturing conditions. According to established values derived from Edelhoch method, $\epsilon_{(280)}$ was calculated using 5500 M⁻¹ cm⁻¹ for tryptophan, 1490 M⁻¹ cm⁻¹ for tyrosine, and 125 M⁻¹ cm⁻¹ for cystine residues.

Peptide mapping by RP-HPLC

Peptide mapping was performed using reversed-phase high-performance liquid chromatography. Samples were denatured in 8 M urea (pH 8.0), reduced with 2.5 M dithiothreitol at 50 °C for 60 min and alkylated with 2.5 M sodium iodoacetate at 25 °C for 30 min. Denatured samples were desalted using PD10 columns and digested overnight using Trypsin (Promega) at 20:1 (sample: Trypsin) ratio at 25 °C. The digestion was stopped by the addition of 2% (v/v) TFA and samples were chromatographically separated on a Zorbax 300 SB C18 column using a HPLC system coupled to a PDA detector (Waters). Data was acquired and processed by Empower (Waters) software.

Far-UV circular dichroism (CD) spectroscopy

Test samples diluted to 0.2 mg/mL in formulation buffer were analysed using a JASCO CD spectrophotometer. Spectra were recorded over 190–260 nm with five accumulations at a scan speed of 100 nm/min. Data acquisition and processing were performed using Spectra Manager software.

Attenuated total reflectance fourier transform infrared spectroscopy (ATR-FTIR)

Test samples diluted to 10.0 mg/mL in formulation buffer were analysed using an ATR cell on a JASCO FTIR instrument. Spectra were recorded over 1800–1400 cm^{-1} with 100 scans at a resolution of 4 cm^{-1} . Data acquisition and processing were performed using Spectra Manager software.

Near-UV circular dichroism (CD) spectroscopy

Test samples diluted to 4.0 mg/mL in formulation buffer were analysed using a JASCO CD spectrophotometer. Spectra were recorded over 250–350 nm with three accumulations at a scan speed of 100 nm/min. Data acquisition and processing were performed using Spectra Manager software.

Intrinsic fluorescence spectroscopy

Test samples diluted to 0.1 mg/mL in formulation buffer were analysed using a Shimadzu RF-6000 fluorescence spectrometer. Spectra were recorded with excitation at 280 nm and emission collected from 300 to 450 nm, using 3 nm excitation and emission bandwidths at a scan speed of 200 nm/min. Data acquisition and processing were performed using LabSolutions software.

Differential scanning calorimetry (DSC)

Test samples diluted to 3.0 mg/mL in formulation buffer. DSC measurements were performed on a Nano DSC instrument (TA Instruments) using Nanorun for experiment setup and NanoAnalyze for data processing. Conditions: scan range 20–100 °C, scan rate 1 °C/min and equilibration time 600 s.

Free Thiol estimation by Ellman's assay

Free thiol analysis was performed using a Shimadzu UV-1800 UV-Visible spectrophotometer equipped with quartz cuvettes. L-cysteine calibration standards ranging from 0 to 160 μM were prepared from a 1500 μM stock solution. Test samples and standards were prepared in 100 mM phosphate buffer containing 1 mM EDTA (pH 7.4) and mixed with 25 mM DTNB following incubation with guanidinium hydrochloride to facilitate thiol exposure. All standards and samples were combined with DTNB and incubated for 30 min at room temperature in the dark. Absorbance was measured at 412 nm. A blank solution containing all assay reagents but excluding the protein and cysteine standards was used to auto-zero the instrument. All measurements were performed in duplicate, and the average value was used for calculation of free thiol content.

Size-exclusion chromatography with multi-angle light scattering (SEC-MALS)

Test samples were analysed using a TSKgel G3000SWxl column at 25 °C on a Waters HPLC system coupled with Wyatt MALS detection. UV detection was performed at 280 nm with a flow rate of 1.0 mL/min for 20 min. Experiment sequences were configured using Empower software. Post-run, signal alignment and peak definition were performed in Astra software for molecular mass and aggregation analysis.

Size variants by size-exclusion high-performance liquid chromatography (SE-HPLC)

For SE analysis, sample was diluted to 5 mg/mL in mobile phase (phosphate buffer, pH 7.0) and injected on TSK-gel 3000 SWXL using Waters HPLC system. Samples were separated isocratically, and the eluent was monitored at 280 nm.

Sedimentation velocity analytical ultracentrifugation (SV-AUC)

Test samples were diluted in 1x PBS and further adjusted to an OD₂₈₀ of ~0.8. SV-AUC measurements were performed using an Optima Analytical Ultracentrifuge (Beckman Coulter) with an AN50Ti 8-hole rotor and Beckman cells with sapphire windows. Cell alignment was carried out using the ZentriForce Pharma tool. Samples were equilibrated at 20 °C for 1 h and analysed at 40,000 rpm using interference and absorbance optics (280 nm). Data was processed using Ultrascan software (V4.0, rev. 6187).

Low molecular weight species (LMWs) by capillary electrophoresis sodium dodecyl sulphate under non reduced condition (CE SDS - NR)

Test samples were prepared in SDS-MW sample buffer with 10 kDa internal marker and iodoacetamide. Samples were then heated at 70 °C for 5 min and electrophoretically separated using a bare fused-silica capillary (AB Sciex) on an PA 800 Plus (Beckman coulter) using UV detection at 220 nm. Data was acquired and processed using 32 Karat software (Beckman coulter).

Non-glycosylated heavy chain (NgHC) fragment by capillary electrophoresis sodium Dodecyl sulphate under reduced condition (CE SDS-R)

Samples were prepared by reducing the disulfide bridges of the antibody by β -mercaptoethanol and sodium dodecyl sulphate (SDS-MW sample buffer, AB Sciex). The fragments (NgHC, HC, LC) were then subjected to analysis. Deglycosylated samples were prepared in 2 mM Tris-HCl (pH-7.0) and digested overnight with N-Glycanase at 37 °C. Digested samples were mixed with SDS-MW sample buffer, 10 kDa internal marker and 2-mercaptoethanol (Sigma) and incubated at 85 °C for 10 min. Samples were electrophoretically separated using a bare fused-silica capillary (AB Sciex) on an PA 800 Plus (Beckman coulter) using UV detection at 220 nm. Data was acquired and processed by 32 Karat software (Beckman coulter).

Image capillary isoelectric focusing (icIEF)

The icIEF analysis for evaluating charge variants was performed on an image capillary isoelectric focusing instrument (Maurice, Protein Simple). For CpB treatment, samples were digested with CpB (Sigma) at a 20:1 (sample: CpB) ratio at 37 °C for 2 h. Non CpB treated samples were diluted to 2 mg/mL with purified water as intermediate stock. Prepared reaction mix with 2.5 M Urea, 0.35% Methyl Cellulose, 4% Pharmalyte pH 3–10, 15 mM Arginine-, 15 mM Iminodiacetic acid, 1% pI Marker 4.05, 1% pI Marker 9.99, purified water, 2 mg/mL sample was added to the reaction mixture. Both set of samples were transferred to the 96 well plate and centrifuged at 4000 rpm for 10 min. Separation was performed on an image capillary isoelectric focusing instrument (Maurice, Protein Simple) for 11 min.

Charge variants by cation exchange chromatography (CEX-HPLC)

CEX-HPLC was used for the assessment of charge variants of test samples with and without carboxypeptidase B (CpB) treatment. For CpB digestion, samples were prepared in sodium phosphate buffer (pH 7.6) and digested

with CpB (Sigma) at a 20:1 (sample: CpB) ratio at 37 °C for 2 h. After incubation diluted the sample to 1.0 mg/mL with 10 mM Sodium Phosphate (pH 7.6) buffer and injected into the column. Non CpB digested samples were prepared in sodium phosphate buffer (pH 7.6) and injected into the column. Both sets of samples were chromatographically separated on an Agilent Bio WCX Column, NP10, PK using a HPLC system (Waters). Data was acquired and processed by Empower (Waters) software.

Oligosaccharide profiling of N-Glycans

The glycosylation profile was measured by 2-AB labelling and hydrophilic interaction liquid chromatography with fluorescence detection (HILIC-FL) analysis. Briefly, test samples were digested overnight with PNGase F at 37 °C, spun down in ice cold ethanol and vacuum dried. 2-AB labelling was done as per the manufacturer's instructions. 2-AB-labeled samples were analyzed by HILIC-FL on a TSKgel Amide-80 column and detected using a fluorescence detector.

Total sialic acid

Sialic acid content (N-Acetylneuraminic acid (NANA) and N-glycolylneuraminic acid (NGNA)) was analysed using RP-UHPLC-FLR. Samples were prepared with 50 pmol of internal standard (KDN, Ketodeoxynonulosonic acid). The sialic acids were released from the sugar chains by acidic hydrolysis (HCl) and labelled using the fluorescent dye 1,2-diamino-4,5-methylenedioxybenzene (DMB). Samples were then analysed by UHPLC/FLR using Waters H-Class Bio Plus instrument with Fluorescence detection. Data based on NANA and NGNA standard (Sigma) calibration curves were acquired and processed by Empower (Waters).

Monosaccharide composition analysis (neutral and amino sugars)

Samples were prepared by spiking with internal standards, Arabitol and N-Acetylmannosamine, and methanolysed, re-N-acetylated and TMS derivatised and Samples were then examined by gas chromatography / mass Spectrometry (GC-MS) using Perkin Elmer Clarus 680 SQ8T GC-MS instrument.

Nuclear magnetic resonance (NMR)

Test samples were supplemented with 10% D₂O and transferred into NMR tubes. One-dimensional (¹H) NMR spectra were recorded at 303 K on a Bruker DRX800 spectrometer equipped with a triple-resonance cryoprobe.

Inhibition of cellular proliferation

HER2 positive BT474 cells (ATCC, Catalog number: HTB-20) were used to assess the inhibition of cellular proliferation. Trastuzumab internal reference standard and test samples were serially diluted to different concentrations and incubated with BT474 cells. After incubation, resazurin was added. Resazurin is reduced by NADPH or NADH in metabolically active cells to a fluorescent compound, resorufin. The fluorescence is directly proportional to the number of live cells. Fluorescence was read using a microplate reader. The relative potency was calculated from nonlinear 4-PL curves of reference standard and test samples.

HER2 binding activity by ELISA

HER2 positive BT474 cells (ATCC, Catalog number: HTB-20) were used to assess binding activity of trastuzumab to HER2 receptor. BT474 cells were seeded in 96 well plates and incubated overnight. After incubation, cells were fixed using formaldehyde. Trastuzumab internal reference standard and test samples were serially

diluted to different concentrations and incubated with fixed BT474 cells. Anti-human IgG-Fc antibody conjugated with HRP was used as detection antibody. TMB was used as the substrate and the reaction was stopped using a stop solution before reading the OD in a microplate reader. The relative potency was calculated from nonlinear 4-PL curves of reference standard and test samples.

HER2 binding affinity by SPR

Recombinant human HER2 was immobilized on a CM5 sensor chip. Trastuzumab internal reference standard and test samples were serially diluted to different concentrations and were passed over the chip with immobilized rhHER2. 1X HBS EP+ buffer was used to dissociate HER2-trastuzumab complex. The experiment was carried out at 37 °C. Bivalent binding model was used to evaluate the HER2-trastuzumab interaction. The association (K_a) and dissociation (K_d) rate constants were obtained via kinetic fitting, and the equilibrium dissociation constant (K_D) was derived by taking the ratio of dissociation rate constant to the association rate constant. The % relative binding affinity (%RBA) was calculated by taking the mean K_D of reference standard / mean K_D of sample X 100.

ADCC activity by reporter gene assay (RGA)

ADCC activity was determined using engineered Jurkat cells as effector cells and HER2 positive BT474 cells (ATCC, Catalog number: HTB-20) as target cells. Engineered Jurkat cells overexpress FcγRIIIa v or f allelic form (BPS Bioscience, USA, Catalogue number: 60541 and 60540 respectively) on the surface and luciferase reporter gene under control of NFAT transactivation elements. Trastuzumab internal reference standard and test samples were serially diluted to different concentrations and incubated with both Jurkat and BT474 cells. The expression of luciferase was determined using ONE-Step™ Luciferase Assay System. The luciferase expression was read using a microplate reader. Percent relative ADCC activity was estimated by detecting luciferase expression in reference standard and test samples and calculated using nonlinear 4-PL curves.

ADCC activity by PBMC based assay

ADCC activity was also determined using peripheral blood mononuclear cells (PBMC) as effector cells and HER2 expressing CHO luciferase target cells (HER2 CHO-k1 Luc2 cell line custom engineered by Innoprot, Spain). PBMCs with FF and VV genotypes were used separately for sample analysis. Briefly, target cell and PBMC preparations were combined and transferred to assay plates. Serially diluted trastuzumab was added to the cell preparations and incubated. Finally, Steady Glo™ reagent was added, and the luciferase signal was read. The signal is directly proportional to the cytotoxicity. Percent relative potency was estimated by a 4-PL restricted curve using the luciferase expression in reference standard and test samples.

ADCP activity by reporter gene assay (RGA)

ADCP activity was determined using engineered Jurkat cells as effector cells and HER2 positive BT474 cells (ATCC, Catalog number: HTB-20) as target cells. Engineered Jurkat cells overexpress FcγRIIIa h allelic form (BPS Bioscience, USA, Catalogue number: 71273) on the surface and luciferase reporter gene under control of NFAT transactivation elements. Trastuzumab internal reference standard and test samples were serially diluted to different concentrations and incubated with both Jurkat and BT474 cells. The expression of luciferase was determined using ONE-Step™ Luciferase Assay System. The luciferase expression was read using a microplate reader. Percent relative ADCP activity was estimated by detecting luciferase expression in reference standard and test samples and calculated using nonlinear 4-PL curves.

FcγRIa binding affinity by SPR

Anti-His antibody was immobilized on CM5 chip using amine coupling chemistry. His-tagged recombinant human FcγRIa was captured on the anti-His antibody. Trastuzumab internal reference standard and test samples were serially diluted to different concentrations and passed over the chip with captured FcγRIa. 1X HBS EP+ buffer was used to dissociate FcγRIa-trastuzumab complex. The experiment was carried out at 25 °C. 1:1 binding model was used to evaluate the FcγRIa-trastuzumab interaction. The association (K_a) and dissociation (K_d) rate constants were obtained via kinetic fitting, and the equilibrium dissociation constant (K_D) was derived by taking the ratio of dissociation rate constant to the association rate constant. The % relative binding affinity (%RBA) was calculated by taking the mean K_D of reference standard / mean K_D of sample X 100.

FcγRIIA binding by BLI

Biolayer Interferometry (BLI) technology was used to determine the binding affinity of trastuzumab to FcγRIIA (H167) and FcγRIIA (R167) receptors. For assessing the binding affinity of trastuzumab, His-tagged FcγRIIA Receptor (ligand) was loaded on Anti-Penta-Histidine biosensors (HISK1), and trastuzumab (analyte) was bound over the captured FcγRIIA. 1X HBS EP+ buffer was used to dissociate FcγRIIA-trastuzumab complex. 1:1 Binding model was used to evaluate the FcγRIIA-trastuzumab interaction. The association (K_a) and dissociation (K_d) rate constants were obtained via kinetic fitting, and the equilibrium dissociation constant (K_D) was derived by taking the ratio of dissociation rate constant to the association rate constant. The % relative binding affinity (%RBA) was calculated by taking the mean K_D of reference standard / mean K_D of sample X 100.

FcγRIIB binding by BLI

For assessing the binding affinity of trastuzumab, His-tagged FcγRIIB Receptor (ligand) was loaded on Anti-Penta-Histidine biosensors (HISK1), and trastuzumab (analyte) was bound over the captured FcγRIIB. 1X HBS EP+ buffer was used to dissociate FcγRIIB-trastuzumab complex. 1:1 Binding model was used to evaluate the FcγRIIB-trastuzumab interaction. The association (K_a) and dissociation (K_d) rate constants were obtained via kinetic fitting, and the equilibrium dissociation constant (K_D) was derived by taking the ratio of dissociation rate constant to the association rate constant. The % relative binding affinity (%RBA) was calculated by taking the mean K_D of reference standard / mean K_D of sample X 100.

FcγRIIIA binding by SPR

Anti-His antibody was immobilized on CM5 chip using amine coupling chemistry. His-tagged recombinant human FcγRIIIA (V176) and FcγRIIIA (F176) were captured on the anti-His antibody separately. Trastuzumab internal reference standard and test samples were serially diluted to different concentrations and passed over the chip with captured FcγRIIIA. 1X HBS EP+ buffer was used to dissociate FcγRIIIA-trastuzumab complex. The experiment was carried out at 25 °C. Two state binding model was used to evaluate the FcγRIIIA-trastuzumab interaction. The association (K_a) and dissociation (K_d) rate constants were obtained via kinetic fitting, and the equilibrium dissociation constant (K_D) was derived by taking the ratio of dissociation rate constant to the association rate constant. The % relative binding affinity (%RBA) was calculated by taking the mean K_D of reference standard / mean K_D of sample X 100.

FcγRIIIB binding by BLI

For assessing the binding affinity of trastuzumab, His-tagged FcγRIIIB Receptor (ligand) was loaded on Anti-Penta-Histidine biosensors (HISK1), and trastuzumab (analyte) was bound over the captured FcγRIIIB. 1X HBS

EP+ buffer was used to dissociate Fc γ RIIIB-trastuzumab complex. 1:1 Binding model was used to evaluate the Fc γ RIIIB-trastuzumab interaction. The association (K_a) and dissociation (K_d) rate constants were obtained via kinetic fitting, and the equilibrium dissociation constant (K_D) was derived by taking the ratio of dissociation rate constant to the association rate constant. The % relative binding affinity (%RBA) was calculated by taking the mean K_D of reference standard / mean K_D of sample X 100.

FcRn binding by SPR

rhFcRn (ligand) was immobilized on the CM5 sensor chip. Trastuzumab internal reference standard and test samples were serially diluted to different concentrations in PBST (pH 5.7) buffer and passed over the immobilized FcRn. 1X PBS (pH 7.4) buffer was used to dissociate FcRn-trastuzumab complex. The experiment was carried out at 25 °C. Heterogeneous Ligand binding model was used to evaluate the FcRn-trastuzumab interaction. The association (K_{a1} , K_{a2}) and dissociation (K_{d1} , K_{d2}) rate constants were obtained via kinetic fitting, and the equilibrium dissociation constant (K_{D1} , K_{D2}) was derived by taking the ratio of dissociation rate constant to the association rate constant. The % relative binding affinities (%RBA) was calculated by taking the mean K_D of reference standard / mean K_D of sample X 100.

C1q binding by BLI

For assessing the binding affinity, trastuzumab was loaded on FAB2G biosensors, and serially diluted C1q (analyte) was bound over the captured trastuzumab. 1X HBS EP+ buffer was used to dissociate C1q-trastuzumab complex. 1:1 Binding model was used to evaluate the C1q-trastuzumab interaction. The association (K_a) and dissociation (K_d) rate constants were obtained via kinetic fitting, and the equilibrium dissociation constant (K_D) was derived by taking the ratio of dissociation rate constant to the association rate constant. The % relative binding affinity (%RBA) was calculated by taking the mean K_D of reference standard / mean K_D of sample X 100.

Acknowledgements We thank our colleagues for their support during sample testing, data analysis and verification. We thank RoukenBio, Scotland for their support in developing the PBMC-based ADCC methodology and sample testing.

Author contributions Sanjeeva Metikala: Conceptualization, Methodology, Validation, Formal analysis, Investigation, Data Curation, Writing - Original Draft, Visualization, Supervision. Maheswara Rao B: Methodology, Validation, Formal analysis, Investigation, Data Curation, Visualization, Supervision. Venu Kumar Vemula: Methodology, Validation, Formal analysis, Investigation, Data Curation, Visualization, Supervision. D R Yashwanth Kumar: Methodology, Validation, Formal analysis, Investigation, Data Curation, Visualization, Supervision. Yash Patel: Methodology, Validation, Formal analysis, Investigation, Data Curation, Visualization, Supervision. Babul Reddy: Methodology, Validation, Formal analysis, Investigation, Data Curation, Visualization. Naga Satish: Methodology, Validation, Formal analysis, Investigation, Data Curation, Visualization. Vuchuru Nagarjuna Reddy: Methodology, Validation, Formal analysis, Investigation, Data Curation, Visualization. Kambagiri Swamy Kayala: Methodology, Validation, Formal analysis, Investigation, Data Curation, Visualization. Pujitha Yarlagadda: Methodology, Validation, Formal analysis, Investigation, Data Curation, Visualization. Raghu K Chitta: Conceptualization, Validation, Investigation, Writing - Review & Editing, Supervision. Srilalitha Kuruganti: Conceptualization, Validation, Investigation, Writing - Review & Editing, Supervision. Disha Dadke: Conceptualization, Validation, Investigation, Writing - Review & Editing, Supervision.

Data availability The data generated as part of this study is confidential. The data that support the findings of this study are not openly available due to reasons of sensitivity and are available from the corresponding author upon reasonable request.

Declarations

Competing interest Disha Dadke has patent on C1Q BINDING ASSAY pending to CuraTeQ Biologics Private Limited. Other authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Ethical approval All cells were handled in accordance with RCGM guidelines, and all methods were carried out as per ICH Q2 (R1)/(R2), USP 1033 and USP 1034 guidelines. All experimental protocols were approved by internal quality assurance according to good laboratory practices (GLP). All cell lines were used in the study after Institutional Bio-Safety Committee (IBSC) approval. All donor Human PBMC material collections were conducted in full compliance with applicable local, national, and international guidelines, regulations, and ethical standards governing the procurement and use of human biological materials. All collections adhered to relevant legal requirements, including informed consent from all donors prior to participation. All protocols governing the collection of donor materials were reviewed and approved by the appropriate institutional review board (IRB), no NHS REC review was required. Informed consent was obtained prior to the collection of leukapheresis for PBMC isolation. Donor anonymity was maintained, with data processed in line with UK and EG GDPR regulations.

Open Access This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Carlson, R. W. et al. HER2 Testing in Breast Cancer: NCCN Task Force Report and Recommendations. *J. Natl. Compr. Canc. Netw.* **4**, S–1 (2006).
2. US FDA & Herceptin, F. D. A. Labeling and Prescribing Information. www.fda.gov/medwatch. (2010).
3. Hudis, C. A. Trastuzumab — Mechanism of Action and Use in Clinical Practice. *N. Engl. J. Med.* **357**, 39–51 (2007).
4. Pan, L. et al. HER2/PI3K/AKT pathway in HER2-positive breast cancer A review. (2024). <https://doi.org/10.1097/MD.00000000000038508> doi:10.1097/MD.00000000000038508.
5. Mayfield, S., Vaughn, J. P. & Kute, T. E. DNA strand breaks and cell cycle perturbation in herceptin treated breast cancer cell lines. *Breast Cancer Res. Treat.* **70**, 123–129 (2001).
6. Maadi, H., Soheilifar, M. H., Choi, W. S., Moshtaghian, A. & Wang, Z. Trastuzumab Mechanism of Action; 20 Years of Research to Unravel a Dilemma. *Cancers (Basel)* **13**, (2021).
7. Spector, N. L. & Blackwell, K. L. Understanding the mechanisms behind trastuzumab therapy for human epidermal growth factor receptor 2-positive breast cancer. *J. Clin. Oncol.* **27**, 5838–5847 (2009).
8. Brennan, P. J., Kumogai, T., Berezov, A., Murali, R. & Greene, M. I. HER2/neu: Mechanisms of dimerization/oligomerization. *Oncogene* **19**, 6093–6101 (2000).
9. Vu, T. & Claret, F. X. Trastuzumab: updated mechanisms of action and resistance in breast cancer. (2012). <https://doi.org/10.3389/fonc.2012.00062> doi:10.3389/fonc.2012.00062.
10. Molina, M. A. et al. Trastuzumab (Herceptin), a Humanized Anti-HER2 Receptor Monoclonal Antibody, Inhibits Basal and Activated HER2 Ectodomain Cleavage in Breast Cancer Cells. *Med. Oncol. Service.* **61**, 4744–4749 (2001).
11. Shi, Y. et al. Trastuzumab Triggers Phagocytic Killing of High HER2 Cancer Cells In Vitro and In Vivo by Interaction with Fcγ Receptors on Macrophages. *J. Immunol.* **194**, 4379–4386 (2015).
12. Petricevic, B. et al. *Trastuzumab Mediates Antibody-Dependent Cell-Mediated Cytotoxicity and Phagocytosis to the Same Extent in Both Adjuvant and Metastatic HER2/Neu Breast Cancer Patients.* (2013). <http://www.translational-medicine.com/content/11/1/307>
13. Mamidi, S. et al. Lipoplex mediated silencing of membrane regulators (CD46, CD55 and CD59) enhances complement-dependent anti-tumor activity of trastuzumab and pertuzumab. (2013). <https://doi.org/10.1016/j.molonc.2013.02.011> doi:10.1016/j.molonc.2013.02.011.
14. Prang, N. et al. Cellular and complement-dependent cytotoxicity of Ep-CAM-specific monoclonal antibody MT201 against breast cancer cell lines. *Br. J. Cancer.* **92**, 342–349 (2005).
15. Mohan, M. V. T. K. et al. Efficacy and Safety of BP02 (Trastuzumab Biosimilar) in HER2-Positive Metastatic Breast Cancer: A Multicenter Phase III Study. *Clin. Drug Investig.* **44**, 513–525 (2024).
16. Schwabe, C. et al. Safety, Tolerability, and Immunogenicity of BP02 (Trastuzumab Biosimilar) Compared to EU- and US-Approved Trastuzumab in Healthy Adult Male Volunteers: A Phase 1, Randomized, Double-Blind Study. *Oncol. Ther.* **12**, 477–490 (2024).
17. ICH Harmonized Tripartite Guideline. Validation of Analytical Procedures: Text and Methodology Q2(R1). (2005).

18. The United States Pharmacopeial Convention. USP Chap. 1033: Biological Assay Validation. (2016).
19. The United States Pharmacopeial Convention. USP Chap. 1032: Design and Development of Biological Assays. (2016).
20. US FDA. Quality Considerations in Demonstrating Biosimilarity of a Therapeutic Protein Product to a Reference Product Guidance for Industry. <https://www.fda.gov/drugs/guidance-compliance-regulatory-information/guidances-drugs>
21. US FDA. Development of Therapeutic Protein Biosimilars: Comparative Analytical Assessment and Other Quality-Related Considerations Guidance for Industry DRAFT GUIDANCE. <https://www.fda.gov/vaccines-blood-biologics/guidance-compliance-regulatory-information-biologics/biologics-guidances>
22. Committee for Medicinal Products for Human Use (CHMP). Guideline on similar biological medicinal products. (2014). www.ema.europa.eu/contact
23. International conference on harmonisation of technical requirements for registration of pharmaceuticals for human use pharmaceutical development Q8(R2). (2009).
24. Raju, T. S. Terminal sugars of Fc glycans influence antibody effector functions of IgGs. *Curr. Opin. Immunol.* **20**, 471–478 (2008).
25. Ram, S., Kim, D., Ober, R. J. & Ward, E. S. The level of HER2 expression is a predictor of antibody-HER2 trafficking behavior in cancer cells. *MAbs* **6**, 1211–1219 (2014).
26. Neve, R. M. et al. A collection of breast cancer cell lines for the study of functionally distinct cancer subtypes. *Cancer Cell.* **10**, 515–527 (2006).
27. Gogesch, P. et al. The Role of Fc Receptors on the Effectiveness of Therapeutic Monoclonal Antibodies. (2021). <https://doi.org/10.3390/ijms22168947> doi:10.3390/ijms22168947.
28. Wöhner, M. & Nimmerjahn, F. Cytotoxic IgG: Mechanisms, functions, and applications. *Immunity* **58**, 1378–1395 (2025).
29. Kremer, P. G. & Barb, A. W. The weaker-binding Fc γ receptor IIIa F158 allotype retains sensitivity to N-glycan composition and exhibits a destabilized antibody-binding interface. *J. Biol. Chem.* **298**, 102329 (2022).
30. Koene, H. R. et al. Fc γ RIIIa-158V/F Polymorphism Influences the Binding of IgG by Natural Killer Cell Fc γ RIIIa, Independently of the Fc γ RIIIa-48L/R/H Phenotype. *Blood* **90**, 1109–1114 (1997).

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Authors and Affiliations

Sanjeeva Metikala¹ · Maheswara Rao¹ · Venu Kumar Vemula¹ · D. R. Yashwanth Kumar¹ · Yash Patel¹ · Babul Reddy¹ · Naga Satish¹ · Vuchuru Nagarjuna Reddy¹ · Kambagiri Swamy Kayala¹ · Pujitha Yarlagadda¹ · Raghu K. Chitta¹ · Srilalitha Kuruganti¹ · Disha Dadke¹

✉ Disha Dadke
Disha.Dadke@curateqbio.com

¹ CuraTeQ Biologics Pvt. Ltd, Survey no 77&78, Indrakarn Village, Kandi Mandal, Sangareddy Dist., Hyderabad, Telangana 502329, India