

GUIDED LECTURE NOTES
for
PROOF, SET THEORY, AND LOGIC
MATH 330

Part I: Logic and Proof

Textbook materials by
Richard P. Millspaugh
University of North Dakota

Guided notes revised and prepared by
Bryce A. Christopherson
University of North Dakota

Fall 2026

These guided notes accompany the Fall 2026 revised course edition of *Elementary Set Theory: Proof, Set Theory, and Logic*. They are based on Richard P. Millspaugh's original textbook materials and lecture materials prepared and revised by Bryce A. Christopherson.

The blank spaces in the student copy are completed during class. Each blank occupies exactly the same space as the corresponding typed completion in the instructor copy, so the two packets have identical pagination. Definitions, theorem statements, and the surrounding explanations are provided so that class time can focus on examples, proof decisions, and the mathematical steps that connect them.

Copyright © 2018 Richard P. Millspaugh.

The original work is licensed under the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. A copy of the license is available at creativecommons.org/licenses/by-nc-sa/4.0/.

This Fall 2026 guided-note edition revises and adapts the original materials and is distributed under the same CC BY-NC-SA 4.0 license. Richard P. Millspaugh has also confirmed Bryce A. Christopherson's permission to use and revise the book on which these notes are based.

CONTENTS OF PART I

1. Elementary Logic
2. Logical Arguments

CHAPTER 1

ELEMENTARY LOGIC

1.1 PROPOSITIONS

DEFINITION 1.1.

A *proposition* is a statement that is either true or false, but not both. The *truth value* of a proposition is true (T) if the sentence is true and false (F) if the sentence is false.

EXAMPLE 1.2.

Determine whether the following are propositions:

- *Two plus two equals four.*
- *Seven minus three equals twelve.*
- *It will snow in Grand Forks on March 17, 2525.*
- *Go clean your room.*

- *Is it raining outside?*

- *This sentence is false.*

- $x + 3 = 12.$

We will use uppercase letters, frequently P or Q, to stand for variable propositions in much the same way that you might use x or y to represent variable numbers in algebra.

1.2 CONNECTIVES AND TRUTH TABLES

Many of the propositions we are interested in are made up of more elementary propositions. For example, the proposition S : *today is Tuesday and it's raining* is made up of the two propositions P : *today is Tuesday* and Q : *it's raining*. The word *and* in S is called a connective since it gives us a way to connect two propositions and form another. In this section we will look at several kinds of connectives.

1.2.1 CONJUNCTION

DEFINITION 1.3.

The *conjunction* of P and Q , denoted $P \wedge Q$, is true exactly when both P and Q are true.

We will sometimes keep track of truth values in a *truth table*, where we list the truth value of a proposition in terms of the truth values of its elementary propositions.

EXAMPLE 1.4.

Construct a truth table for the conjunction $P \wedge Q$.

P	Q	$P \wedge Q$
T	T	
T	F	
F	T	
F	F	

Note that in this case there are four rows in the table. For a proposition involving n distinct elementary propositions, there are 2^n possible truth assignments and hence 2^n rows in its truth table. This makes it cumbersome to construct truth tables for very complicated propositions. Nevertheless, we will find them to be a convenient tool.

1.2.2 DISJUNCTION

We next consider propositions of the form P *or* Q . The word *or* can be ambiguous in English: sometimes it includes the possibility that both conditions hold, while in other contexts only one option is intended. In logic we use *or* inclusively. Thus $P \vee Q$ is true when at least one of P and Q is true.

DEFINITION 1.5.

The *disjunction* of P and Q , denoted $P \vee Q$, is true whenever at least one of P or Q is true.

EXAMPLE 1.6.

Construct a truth table for the proposition $P \vee Q$.

P	Q	$P \vee Q$
T	T	
T	F	
F	T	
F	F	

1.2.3 IMPLICATION

Most mathematical propositions have the form *If P, then Q*. Of course, either of P or Q might be propositions made of several other propositions.

DEFINITION 1.7.

The proposition *If P, then Q* is called an *implication* and is denoted $P \Rightarrow Q$. It is false exactly when P is true and Q is false.

EXAMPLE 1.8.

Construct a truth table for $P \Rightarrow Q$.

P	Q	$P \Rightarrow Q$
T	T	
T	F	
F	T	
F	F	

Notice: To say that $P \Rightarrow Q$ is true **does not** mean that P is true or that Q is true. It **does** mean that there is a relationship between the truth values of these two propositions. The implication $P \Rightarrow Q$ is false when P is true and Q is false; it is true when either P is false or Q is true.

There are several other implications related to the implication $P \Rightarrow Q$ that we will occasionally find useful.

DEFINITION 1.9.

For the implication $P \Rightarrow Q$:

- The *converse* is $Q \Rightarrow P$.
- The *contrapositive* is $\neg Q \Rightarrow \neg P$.
- The *inverse* is $\neg P \Rightarrow \neg Q$.

1.2.4 LOGICAL EQUIVALENCE

DEFINITION 1.10.

The *biconditional* $P \Leftrightarrow Q$, read “P if and only if Q,” is true exactly when P and Q have the same truth value under the given assignment. Two propositional forms P and Q are *logically equivalent* if they have the same truth value under every assignment of truth values to their elementary propositions. Equivalently, $P \Leftrightarrow Q$ is a tautology.

EXAMPLE 1.11.

Construct a truth table for $P \Leftrightarrow Q$.

P	Q	$P \Leftrightarrow Q$
T	T	
T	F	
F	T	
F	F	

1.2.5 NEGATION

DEFINITION 1.12.

The *negation* of P, denoted $\neg P$ and sometimes read “not P,” is the proposition whose truth value is always opposite that of P.

Negation is a unary connective: it acts on one proposition, whereas conjunction, disjunction, implication, and biconditional are binary connectives.

EXAMPLE 1.13.

Construct a truth table for P and $\neg P$.

P	$\neg P$
T	
F	

1.2.6 TAUTOLOGY AND CONTRADICTION

DEFINITION 1.14.

A *tautology* is a proposition that is always true, regardless of the truth values of the elementary propositions that make it up.

DEFINITION 1.15.

A *contradiction* is a proposition that is always false.

EXAMPLE 1.16.

Show the proposition $P \vee \neg P$ is a tautology and that the proposition $P \wedge \neg P$ is a contradiction using a truth table.

P	$\neg P$	$P \vee \neg P$	$P \wedge \neg P$
T			
F			

EXAMPLE 1.17.

Use a truth table to show that $P \Rightarrow (P \vee Q)$ is a tautology.

P	Q	$P \vee Q$	$P \Rightarrow (P \vee Q)$
T	T		
T	F		
F	T		
F	F		

EXAMPLE 1.18.

Use a truth table to show that $(P \Rightarrow Q) \Leftrightarrow (\neg Q \Rightarrow \neg P)$ is a tautology.

P	Q	$\neg P$	$\neg Q$	$P \Rightarrow Q$	$\neg Q \Rightarrow \neg P$
T	T				
T	F				
F	T				
F	F				

1.3 PREDICATES, INSTANTIATION, AND QUANTIFICATION

We return our attention now to statements that involve variables, like $x + 3 = 12$. Such a statement is called a *predicate* and we will usually indicate it with $P(x)$ rather than P . Before a predicate takes on a truth value, we need to know something about the variable. We noted before that if $x = 0$, then the statement is false; but if $x = 9$, then the statement is true. What if x represents my desk? In that case the statement is complete nonsense, but you probably feel like there's something a little bit unfair about that choice for x . We see an equation and usually think that the variable must represent a number, right? The first thing you should know about any variable in a predicate is what it can represent. This is called the *universe* or the *domain of discourse*. In mathematics, we will be very explicit about what universe we are talking about.

1.3.1 INSTANTIATION

Suppose the universe for the preceding predicate $P(x)$ is $\mathbb{R}$. That still doesn't turn $P(x)$ into a proposition because, as noted above, $P(x)$ doesn't take on a truth value until we know which real number x indicates. One way to do this is called *instantiation*, which means choosing a particular value for the variable(s). Consider for example the sentence: *Let $x = 2$, then $x + 3 = 12$* . In this case the sentence is a proposition with truth value F. The other option is *quantification* of the variable(s). We will discuss two kinds of quantifiers, *universal* and *existential*. You might occasionally encounter other quantifiers, but they are not necessary.

1.3.2 UNIVERSAL QUANTIFICATION

DEFINITION 1.19.

A *universal quantifier* indicates that the predicate is true for all instances of the variable in the given universe. It is typically indicated by using the phrase “for every” or “for all,” and can be denoted symbolically by $\forall$ in symbolic statements.

The following sentences are universally quantified propositions. Only the second is true.

- *For every real number x , $x + 3 = 12$.*
- *The square of x is nonnegative for all real x .*
- *The square of x is real for all complex x .*

The third statement is false; for example, $(1 + i)^2 = 2i$, which is not real.

1.3.3 EXISTENTIAL QUANTIFICATION

DEFINITION 1.20.

An *existential quantifier* indicates that the predicate is true for at least one instance of the variable in the given universe. It is typically indicated by the phrase “for some” or “there exists,” and can be denoted by $\exists$ in symbolic statements.

The following sentences are existentially quantified. All three of these are true.

- *There is a real number x so that $x + 3 = 12$.*
- *The square of x is nonnegative for some real x .*
- *There is a complex number x whose square is -1 .*

1.3.4 SYLLOGISMS AND VENN DIAGRAMS

A *syllogism* is a form of logical argument that deduces a conclusion based on two premises. For example:

All men are mortal. Socrates is a man. Hence Socrates is mortal.

This is a valid syllogism. If we assume that the premises are true, then the conclusion must follow. One tool we can use to help think about syllogisms is a *Venn diagram*, which in its simplest form is a collection of circles that represent the categories in the premises.

EXAMPLE 1.21.

Construct a Venn diagram to determine if the following syllogism is valid: *All men are mortal. Socrates is a man. Hence Socrates is mortal.*

EXAMPLE 1.22.

Determine whether the following syllogism is valid:

Some students are freshmen. Pat is a student. Hence Pat is a freshman.

EXAMPLE 1.23.

Determine whether the following syllogism is valid:

No cats are dogs. Jade is a cat. Hence Jade is not a dog.

EXAMPLE 1.24.

Determine whether the following syllogism is valid:

Some cats are psychopaths. Jawa is a cat. Hence Jawa is a psychopath.

EXAMPLE 1.25.

Determine whether the following syllogism is valid:

Some Billywiggles are Bleepzigs. Some Bleepzigs are Jabberwoks. Hence some Billywiggles are Jabberwoks.

1.3.5 QUANTIFICATION OF SEVERAL VARIABLES

A predicate can have two or more variables. In order to use the predicate in a proposition, a universe must be declared for each variable and each variable must be instantiated or quantified. We will discuss the various ways a two-variable predicate might be quantified by considering an example.

EXAMPLE 1.26.

Consider the predicate $P(x, y)$ given by $x + y = 0$ and assume that the universe for both x and y is the set of real numbers. Determine whether the following proposition is true: $\forall x \forall y P(x, y)$

EXAMPLE 1.27.

Consider the predicate $P(x, y)$ given by $x + y = 0$ and assume that the universe for both x and y is the set of real numbers. Determine whether the following proposition is true: $\forall y \forall x P(x, y)$.

EXAMPLE 1.28.

Consider the predicate $P(x, y)$ given by $x + y = 0$ and assume that the universe for both x and y is the set of real numbers. Determine whether the following proposition is true: $\exists x \exists y P(x, y)$.

EXAMPLE 1.29.

Consider the predicate $P(x, y)$ given by $x + y = 0$ and assume that the universe for both x and y is the set of real numbers. Determine whether the following proposition is true: $\forall x \exists y P(x, y)$.

EXAMPLE 1.30.

Consider the predicate $P(x, y)$ given by $x + y = 0$ and assume that the universe for both x and y is the set of real numbers. Determine whether the following proposition is true: $\exists y \forall x P(x, y)$.

1.4 NEGATIONS

The negation of a proposition is the proposition with the opposite truth value. In this section we are going to discuss the negations of compound propositions, which can be an important step in some kinds of proofs. We begin with two theorems that tell how to find the negations of propositions involving connectives.

THEOREM 1.31.

Given any propositions P and Q , $\neg(P \Rightarrow Q) \Leftrightarrow (P \wedge \neg Q)$.

Proof. Complete the truth table below and compare its final two columns.

P	Q	$\neg Q$	$P \Rightarrow Q$	$\neg(P \Rightarrow Q)$	$P \wedge \neg Q$
T	T				
T	F				
F	T				
F	F				

THEOREM 1.32 (De Morgan's Laws).

Given any propositions P and Q :

(i) $\neg(P \wedge Q) \Leftrightarrow (\neg P \vee \neg Q)$

(ii) $\neg(P \vee Q) \Leftrightarrow (\neg P \wedge \neg Q)$

You will prove this theorem in the homework.

EXAMPLE 1.33.

Negate the proposition $P \Rightarrow (Q \wedge R)$.

EXAMPLE 1.34.

Find the negation of $P \wedge (Q \Rightarrow R)$.

1.4.1 NEGATING QUANTIFIED STATEMENTS

The quantifiers $\forall$ and $\exists$ are interchanged by negation. This is the quantified analogue of De Morgan's Laws.

THEOREM 1.35 (Quantified Negation). *Let $P(x)$ be a predicate on a specified domain. Then*

$$\neg(\forall x P(x)) \iff \exists x \neg P(x)$$

and

$$\neg(\exists x P(x)) \iff \forall x \neg P(x).$$

Proof.

For nested quantifiers, move the negation inward one quantifier at a time. For example,

$$\neg(\forall x \exists y Q(x, y)) \iff \exists x \forall y \neg Q(x, y).$$

Notice that in $\forall x \exists y Q(x, y)$ the witness y may depend on x ; reversing the quantifiers would assert the existence of one choice of y that works for every x .

EXAMPLE 1.36.

Find the negation of $(P \vee Q) \wedge (P \vee R)$.

CHAPTER 2

LOGICAL ARGUMENTS

2.1 RULES OF INFERENCE

We accept the following two basic rules of thought:

The *Law of Excluded Middle* says that given any proposition P , we may deduce $P \vee \neg P$. In words, either P is true or $\neg P$ is true.

The *Law of non-Contradiction* says that given any proposition P , we may deduce $\neg(P \wedge \neg P)$. In words, P and $\neg P$ cannot both be true.

A *rule of inference* is a logical rule consisting of a hypothesis and a conclusion. Whenever the hypothesis is true, the conclusion must also be true. For our purposes this means that once we have deduced all of the hypotheses, we may also deduce the conclusion. Rules of inference are closely related to tautologies. In fact, they are essentially the same thing. While every tautology gives rise to a rule of inference, in practice the seven listed in the table below will usually suffice.

Table 2.1: Rules of Inference

Name	Hypotheses	Conclusion
Modus ponens	P and $P \Rightarrow Q$	Q
Modus tollens	$\neg Q$ and $P \Rightarrow Q$	$\neg P$
Hypothetical syllogism	$P \Rightarrow Q$ and $Q \Rightarrow R$	$P \Rightarrow R$
Addition	P	$P \vee Q$
Simplification	$P \wedge Q$	P
Conjunction	P and Q	$P \wedge Q$
Disjunctive syllogism	$P \vee Q$ and $\neg P$	Q

The Rule of Addition, for example, comes from the tautology $P \Rightarrow (P \vee Q)$. Since the implication is always true, knowing that the hypothesis is true will tell us that the conclusion must also be true. We can verify each of the Rules of Inference in Table 2.1 by constructing a truth table for the associated tautology.

2.1.1 PROPOSITIONAL ARGUMENTS

Each of our Rules of Inference is actually a *propositional argument*, where accepting the hypotheses forces us to accept the conclusion. We use the following notation for propositional arguments, where the propositions above the line are hypotheses and the proposition(s) below the line are the conclusion(s). We precede the conclusion(s) with the symbol $\therefore$, which is read “therefore.”

EXAMPLE 2.1.

Here is the rule of modus tollens as a propositional argument:

$$\begin{array}{c} \neg Q \\ P \Rightarrow Q \\ \hline \therefore \neg P \end{array}$$

EXAMPLE 2.2.

Express the rule of modus ponens as a propositional argument.

EXAMPLE 2.3.

Express the rule of hypothetical syllogism as a propositional argument.

DEFINITION 2.4.

A propositional argument is said to be *valid* if accepting the hypotheses forces us to accept the conclusion(s).

How do we determine whether or not a propositional argument is valid? One option is to construct a truth table, but that can be very tedious. Our preferred option is to construct a logical argument which begins by assuming each of the hypotheses is true and deduces consequences using our rules of inference. Each deduced proposition is a step in our proof that can be justified as either a hypothesis, a tautology, a consequence of previous steps and a rule of inference, or a proposition that is logically equivalent to a previous step. The proof is complete when we have deduced the desired conclusion.

EXAMPLE 2.5.

Prove that the following argument is valid:

$$\begin{array}{c} P \\ P \Rightarrow Q \\ (Q \vee R) \Rightarrow S \\ \hline \therefore S \end{array}$$

EXAMPLE 2.6.

Prove that the following argument is valid:

$$\frac{P \Rightarrow (Q \wedge \neg Q)}{\therefore \neg P}$$

The argument in Example 2.6 says that if a proposition P implies a contradiction, then $\neg P$ must be true. This result is the basis for proofs by contradiction.

2.1.2 ARGUMENTS WITH QUANTIFIERS

Arguments involving quantified propositions require rules for instantiation and generalization. The *rules for instantiation* let us deduce statements about particular elements of the domain from quantified statements. The *rules for generalization* tell us when deductions about elements of the domain justify quantified statements. We give these rules in Table 2.2.

Table 2.2: Rules of Quantification

Name	Rule
Universal Instantiation	$\forall x P(x) : P(c)$ whenever c is in the domain of x
Existential Instantiation	$\exists x P(x) : P(c)$ for some particular but unspecified c in the domain of x
Universal Generalization	$P(c)$ for <i>arbitrary</i> c in the domain of x : $\forall x P(x)$
Existential Generalization	$P(c)$ for <i>some</i> c in the domain of x : $\exists x P(x)$

The restrictions in this table matter. In existential instantiation, the letter c denotes one particular but unspecified value whose existence is guaranteed by the existential statement. We do not get to choose which value it is, and we may not assume anything about it beyond what the existential statement tells us. The letter c should therefore not already be in use elsewhere in the argument. In universal generalization, by contrast, c represents an arbitrary element of the domain.

EXAMPLE 2.7.

Prove that the following argument is valid:

$$\begin{array}{l} \forall x (P(x) \wedge Q(x)) \\ \exists x (P(x) \Rightarrow R(x)) \\ \hline \therefore \exists x (Q(x) \wedge R(x)) \end{array}$$

2.2 AXIOMS AND FORMAL MATHEMATICAL THEORIES

The rules of inference developed above tell us which deductive steps are valid because of their logical form. To develop a particular area of mathematics, however, we also need starting statements about the objects we intend to study. A *formal mathematical theory* consists of a specified language, logical principles and rules of inference, and a collection of *axioms*. An axiom is a statement assumed throughout the theory rather than proved from earlier statements in that theory.

Axioms should be distinguished from hypotheses and definitions. An axiom is a standing assumption of the entire theory, while a hypothesis is usually a temporary assumption made within a particular argument. A definition assigns meaning to a new term or symbol; it does not, by itself, guarantee that an object having the stated properties exists.

A *theorem* is a statement that can be deduced from the axioms using the accepted rules of inference. For example, if P and $P \Rightarrow Q$ are axioms or previously proved theorems, then modus ponens allows us to deduce Q . Once proved, Q may itself be used in later arguments. In a completely formal proof, every such step is written down and justified.

Calling a statement an axiom does not mean that it is self-evident or true in every possible mathematical setting. It describes the role that the statement plays within a particular theory. Different choices of axioms can lead to different theories and therefore to different collections of theorems.

Ordinary mathematical proofs do not include every elementary formal step. Instead, we write in mathematical English, make the important deductions explicit, and suppress routine applications of logic. Such proofs are nevertheless intended to be formalizable: in principle, the omitted steps could be supplied within an appropriate axiomatic theory. In the next chapter we will briefly examine ZFC, the standard axiomatic framework underlying most ordinary work with sets.

2.3 PROVING MATHEMATICAL THEOREMS

Most mathematical statements are implications of the form $P \Rightarrow Q$, where P or Q might be compound propositions. In this section we will look at some methods for proving implications. Note that we are using some definitions and results from other areas of mathematics that we have not explicitly stated. We will not allow ourselves to make those kinds of assumptions once we begin our study of set theory.

2.3.1 DIRECT PROOFS

A direct proof of the implication $P \Rightarrow Q$ assumes that the hypothesis P is true, then uses definitions, previously proved theorems, and the rules of inference to deduce that the conclusion Q must also be true.

EXAMPLE 2.8.

Prove the following theorem:

THEOREM 2.9.

If n is an even integer, then n^2 is even.

Proof.

Although the two-column proof above is valid, it is not written in a format that mathematicians usually find pleasing. We prefer proofs that are written using complete sentences, paragraphs, and correct grammar and punctuation. Here is a better presentation for the previous proof.

EXAMPLE 2.10.

Prove Theorem 2.9 in plain language.

Proof.

EXAMPLE 2.11.

Prove the following theorem:

THEOREM 2.12.

The square of an odd integer is odd.

Proof.

2.3.2 PROVING THE CONTRAPOSITIVE

As you have shown in a homework exercise, the contrapositive of an implication is logically equivalent to the implication. Sometimes it is easier to prove the contrapositive of the implication we are interested in, as we will show here.

EXAMPLE 2.13.

Prove the following theorem:

THEOREM 2.14.

If n is an integer and n^2 is even, then n is even.

Proof.

EXAMPLE 2.15.

Prove the following theorem:

THEOREM 2.16. *If n is an integer and n^2 is odd, then n is odd.*

Proof.

2.3.3 PROOF BY CONTRADICTION

A proof by contradiction, or *reductio ad absurdum*, begins by assuming that the result we are trying to prove is false, then deriving a contradiction from that assumption. It is easy to confuse this with a direct proof of the contrapositive, which makes different beginning assumptions. To prove the implication $P \Rightarrow Q$ by proving the contrapositive, you assume $\neg Q$ and use that to prove $\neg P$. A proof by contradiction assumes $P \wedge \neg Q$ and derives a contradiction $R \wedge \neg R$. Here is a classic example of a proof by contradiction.

EXAMPLE 2.17.

Prove the following theorem:

THEOREM 2.18.

There is no rational number x such that $x^2 = 2$.

Proof.

2.3.4 PROOF BY CASES

To prove the implication $P \Rightarrow Q$ by cases, we make use of the following argument:

$$\begin{array}{c} P \Rightarrow (R \vee S) \\ R \Rightarrow Q \\ S \Rightarrow Q \\ \hline \therefore P \Rightarrow Q \end{array}$$

You will be asked to show that this argument is valid in the homework.

Here is a very simple example of a proof by cases. We make use of some of the order properties of the real numbers. In particular, we know that multiplying both sides of an inequality by a positive number preserves the inequality, and multiplying both sides of an inequality by a negative number reverses the inequality.

EXAMPLE 2.19.

Prove the following theorem:

THEOREM 2.20.

If x is a real number, then $x^2 \geq 0$.

Proof.

2.4 PROVING QUANTIFIED STATEMENTS

2.4.1 UNIVERSALLY QUANTIFIED STATEMENTS

On the simplest level, a universally quantified statement has the form $\forall x P(x)$. To prove it, let x be an arbitrary element of the required domain and prove $P(x)$ without using any special property of x beyond its membership in that domain. It is frequently helpful to think of such a statement as an implication whose hypothesis is that x belongs to the required domain and whose conclusion is $P(x)$. Since universally quantified statements can be viewed as implications, we can use the same proof techniques.

2.4.2 EXISTENTIALLY QUANTIFIED STATEMENTS

Existentially quantified statements are significantly different than implications and universally quantified statements. To prove the statement $\exists x, P(x)$ we must show that there is at least one x in the domain that satisfies $P(x)$. One way to do this is simply to find a single particular example.

EXAMPLE 2.21.

Prove the following theorem:

THEOREM 2.22.

There is a positive integer N such that N , $N + 1$, $N + 2$, $N + 3$, and $N + 4$ are all composite (not prime).

Proof.

There are some instances where it is easier to show that some x satisfies $P(x)$ without actually finding a particular x that works. Here is an elementary example that works by demonstrating that one of two possible choices must work, without determining which it is.

EXAMPLE 2.23.

Prove the following theorem:

THEOREM 2.24.

There exist irrational numbers a and b so that a^b is rational.

Proof.

2.4.3 COUNTEREXAMPLES

An example showing that a universally quantified statement is false is called a *counterexample* to that statement.

EXAMPLE 2.25.

Disprove the following proposition:

CONJECTURE 2.26.

Every integer is even.

2.5 MATHEMATICAL INDUCTION

There is another type of proof that is frequently used to prove statements about the natural numbers, i.e., the numbers $1, 2, 3, \dots$

THEOREM 2.27 (The Principle of Mathematical Induction).

Let $P(n)$ be a statement about the natural number n . If $P(1)$ is true and if $P(k) \implies P(k + 1)$ for every natural number k , then $P(n)$ is true for every natural number n .

All proofs by induction use the following basic outline. To prove $P(n)$ for all natural numbers n ,

1. Base Case: Show that $P(1)$ is true.
2. Inductive Step: Prove the implication $P(k) \implies P(k + 1)$ for all $k \in \mathbb{N}$. Note: in proving that $P(k) \implies P(k + 1)$, the assumption that $P(k)$ is true is often referred to as the inductive hypothesis.
3. It is considered good form to clearly state that the statement $P(n)$ is now true for all natural numbers n by induction.

EXAMPLE 2.28.

Prove the following theorem:

THEOREM 2.29.

For each natural number n , $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.

Proof.