

Gray-type monoidal product and Bifunctor Theorem for double categories

Bojana Femić

Virtual Double Categories Workshop

29 November 2022

Mathematical Institute of
Serbian Academy of Sciences and Arts
Belgrade (Serbia)

Overview of the talk

Introductory part:

- ▶ double categories (as specific internal categories)

Overview of the talk

Introductory part:

- ▶ double categories (as specific internal categories)
- ▶ double category of monads
(why “vertical morphisms” of monads are useful)

Overview of the talk

Introductory part:

- ▶ double categories (as specific internal categories)
- ▶ double category of monads
(why “vertical morphisms” of monads are useful)

Gray-type tensor product on double categories

- ▶ Gray-type tensor product on (strict-strict) double categories

Overview of the talk

Introductory part:

- ▶ double categories (as specific internal categories)
- ▶ double category of monads
(why “vertical morphisms” of monads are useful)

Gray-type tensor product on double categories

- ▶ Gray-type tensor product on (strict-strict) double categories
- ▶ Bifunctor Theorem for (lax-hop) double categories

Overview of the talk

Introductory part:

- ▶ double categories (as specific internal categories)
- ▶ double category of monads
(why “vertical morphisms” of monads are useful)

Gray-type tensor product on double categories

- ▶ Gray-type tensor product on (strict-strict) double categories
- ▶ Bifunctor Theorem for (lax-hop) double categories
 - ▶ (no) Gray-type tensor product on (strict-lax) double categories
 - ▶ Bifunctor Theorem
 - ▶ “(Un)currying” 2-functors
 - ▶ application to monads in double categories

Double categories

Double categories

Bicategory:

- 0-cells
- 1-cells
- 2-cells

Double categories

Bicategory:

- 0-cells
- 1-cells
- 2-cells

Double category:

- 0-cells
- vertical 1-cells
- horizontal 1-cells
- squares (2-cells)

Double category as internal category

A **double category** is an internal category in the category \mathbf{Cat}_1 .

Double category as internal category

A **double category** is an internal category in the category \mathbf{Cat}_1 .

There are:

- categories C_0, C_1 (0-cells in \mathbf{Cat}_1)

Double category as internal category

A **double category** is an internal category in the category \mathbf{Cat}_1 .

There are:

- categories C_0, C_1 (0-cells in \mathbf{Cat}_1)
- functors (1-cells in \mathbf{Cat}_1)

$$s, t : C_1 \rightarrow C_0, \quad u : C_0 \rightarrow C_1 \quad \text{and} \quad c : C_1 \times_{C_0} C_1 \rightarrow C_1$$

Double category as internal category

A **double category** is an internal category in the category \mathbf{Cat}_1 .

There are:

- categories C_0, C_1 (0-cells in \mathbf{Cat}_1)
- functors (1-cells in \mathbf{Cat}_1)

$s, t : C_1 \rightarrow C_0$, $u : C_0 \rightarrow C_1$ and $c : C_1 \times_{C_0} C_1 \rightarrow C_1$
s.t.

- c is associative and unital.

Double category as internal category

A **double category** is an internal category in the category \mathbf{Cat}_1 .

There are:

- categories C_0, C_1 (0-cells in \mathbf{Cat}_1)
- functors (1-cells in \mathbf{Cat}_1)


$s, t : C_1 \rightarrow C_0$, $u : C_0 \rightarrow C_1$ and $c : C_1 \times_{C_0} C_1 \rightarrow C_1$
s.t.

- c is associative and unital.

- 0-cells

- vertical 1-cells

- horizontal 1-cells



- squares (2-cells)

Double category as internal category

A **double category** is an internal category in the category \mathbf{Cat}_1 .

There are:

- categories C_0, C_1 (0-cells in \mathbf{Cat}_1)
- functors (1-cells in \mathbf{Cat}_1)


$s, t : C_1 \rightarrow C_0$, $u : C_0 \rightarrow C_1$ and $c : C_1 \times_{C_0} C_1 \rightarrow C_1$
s.t.

- c is associative and unital.

- 0-cells

- vertical 1-cells

- horizontal 1-cells



- squares (2-cells)

C_0 : 0-cells and 1v-cells, C_1 : 1h-cells and 2-cells.

Pseudodouble category as internal category

A **pseudodouble category** is a
pseudocategory internal in the 2-category \mathbf{Cat}_2 .

Pseudodouble category as internal category

A **pseudodouble category** is a
pseudocategory internal in the 2-category \mathbf{Cat}_2 .
(an internal category in \mathbf{Cat}_2)

Pseudodouble category as internal category

A **pseudodouble category** is a
pseudocategory internal in the 2-category \mathbf{Cat}_2 .
(an internal category in \mathbf{Cat}_2)

There are:

- categories C_0, C_1 (0-cells in \mathbf{Cat}_2)
- functors (1-cells in \mathbf{Cat}_2)

$$s, t : C_1 \rightarrow C_0, \quad u : C_0 \rightarrow C_1 \quad \text{and} \quad c : C_1 \times_{C_0} C_1 \rightarrow C_1$$

Pseudodouble category as internal category

A **pseudodouble category** is a
pseudocategory internal in the 2-category \mathbf{Cat}_2 .
 (an internal category in \mathbf{Cat}_2)

There are:

- categories C_0, C_1 (0-cells in \mathbf{Cat}_2)
- functors (1-cells in \mathbf{Cat}_2)

$$s, t : C_1 \rightarrow C_0, \quad u : C_0 \rightarrow C_1 \quad \text{and} \quad c : C_1 \times_{C_0} C_1 \rightarrow C_1$$

- natural transformations (2-cells in \mathbf{Cat}_2)

$$\alpha : c \otimes (id_{C_1} \times_{C_0} c) \Rightarrow c \otimes (c \times_{C_0} id_{C_1})$$

$$\lambda : c \otimes (u \times_{C_0} id_{C_1}) \Rightarrow id_{C_1}$$

$$\rho : c \otimes (id_{C_1} \times_{C_0} u) \Rightarrow id_{C_1}$$

which satisfy a pentagon and a triangle.

Bicategories and pseudodouble categories

A **bicategory** known to everyone:

0: algebras A, B, \dots , 1: A - B -bimodules, 2: bimodule morphisms.

Bicategories and pseudodouble categories

A **bicategory** known to everyone:

0: algebras A, B, \dots , 1: A - B -bimodules, 2: bimodule morphisms.

Pseudodouble category:

- 0: algebras A, B
- 1v: alg. morphisms
- 1h: A - B -bimodules
- 2: bimodule morphisms

Bicategories and pseudodouble categories

A **bicategory** known to everyone:

0: algebras A, B, \dots , 1: A - B -bimodules, 2: bimodule morphisms.

Pseudodouble category:

- 0: algebras A, B

- 1v: **alg. morphisms**

- 1h: A - B -bimodules

- 2: bimodule morphisms

$$\begin{array}{ccc} B & \xrightarrow{M} & A \\ f \downarrow & \boxed{\alpha} & \downarrow g \\ B' & \xrightarrow{N} & A' \end{array}$$

$\alpha : M \rightarrow N$ A - B -bimodule morphism

$$a \cdot n \cdot b := g(a) \cdot n \cdot f(b)$$

Bicategories and pseudodouble categories

- ▶ In every pseudodouble category there is a bicategory:

Bicategories and pseudodouble categories

- In every pseudodouble category there is a bicategory:
for vertical morphisms take only identities,

Bicategories and pseudodouble categories

- In every pseudodouble category there is a bicategory:
for vertical morphisms take only identities,
and correspondingly “globular 2-cells”.

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow & & \downarrow \\ A & \xrightarrow{g} & B \end{array}$$

Bicategories and pseudodouble categories

- In every pseudodouble category there is a bicategory:
for vertical morphisms take only identities,
and correspondingly “globular 2-cells”.

$$\begin{array}{ccc}
 A & \xrightarrow{f} & B \\
 \downarrow & & \downarrow \\
 A & \xrightarrow{g} & B
 \end{array}
 \begin{array}{c}
 = \\
 \\
 =
 \end{array}$$

(This is the *horizontal bicategory* $\mathcal{H}(\mathbb{D})$ of the pseudodouble category \mathbb{D} .)

- In every pseudodouble category there is a bicategory:
for vertical morphisms take only identities,
and correspondingly “globular 2-cells”. f

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow & & \downarrow \\ A & \xrightarrow{g} & B \end{array}$$

(This is the *horizontal bicategory* $\mathcal{H}(\mathbb{D})$ of the pseudodouble category \mathbb{D} .)

- Each bicategory can be embedded into a pseudodouble category.

Why
the 2-category $\mathbf{Mnd}(\mathcal{K})$
of 2-monads
isn't enough

The 2-category $\mathbf{Mnd}(\mathcal{K})$ of monads in \mathcal{K}

0-cells:

2-monads $(\mathcal{A}, T : \mathcal{A} \rightarrow \mathcal{A}, \mu_T : TT \rightarrow T, \eta_T : \mathrm{Id}_{\mathcal{A}} \rightarrow T)$

1-cells: pairs $(X, \psi) : (\mathcal{A}, T) \rightarrow (\mathcal{A}', T')$ where $X : \mathcal{A} \rightarrow \mathcal{A}'$ is a 1-cell and $\psi : T'X \Rightarrow XT$ a 2-cell s.t.

$$\begin{array}{c} T' \quad T' \quad X \\ \text{---} \boxed{\psi} \text{---} \\ \text{---} \boxed{\psi} \text{---} \\ X \quad T \end{array} = \begin{array}{c} T' \quad T' \quad X \\ \text{---} \text{---} \text{---} \\ \text{---} \boxed{\psi} \text{---} \\ X \quad T \end{array} ; \quad \begin{array}{c} X \\ \text{---} \text{---} \text{---} \\ \text{---} \boxed{\psi} \text{---} \\ X \quad T \end{array} = \begin{array}{c} X \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ X \quad T \end{array}$$

2-cells: $(X, \psi) \Rightarrow (Y, \psi')$ are given by 2-cells $\zeta : X \rightarrow Y$ in \mathcal{K} satisfying:

$$\begin{array}{c} T' \quad X \\ \text{---} \boxed{\psi} \text{---} \\ \text{---} \boxed{\zeta} \text{---} \\ Y \quad T \end{array} = \begin{array}{c} T' \quad X \\ \text{---} \text{---} \text{---} \\ \text{---} \boxed{\psi'} \text{---} \\ Y \quad T \end{array}$$

Monads and monad morphisms

Categories, operads, multicategories and T-multicategories

Monads and monad morphisms

Categories, operads, multicategories and T-multicategories **can all be seen as monads in appropriate bicategories.**

Monads and monad morphisms

Categories, operads, multicategories and T-multicategories **can all be seen as monads in appropriate bicategories.**

However, morphisms between them are not monad maps.

Monads and monad morphisms

Categories, operads, multicategories and T-multicategories **can all be seen as monads in appropriate bicategories.**

However, morphisms between them are not monad maps.

- internal categories (with a discrete object of objects) are monads in $\mathbf{Span}(\mathcal{V})$ (resp. in $\mathbf{Span}_d(\mathcal{V})$)

Monads and monad morphisms

Categories, operads, multicategories and T-multicategories **can all be seen as monads in appropriate bicategories.**

However, morphisms between them are not monad maps.

- internal categories (with a discrete object of objects) are monads in $\mathbf{Span}(\mathcal{V})$ (resp. in $\mathbf{Span}_d(\mathcal{V})$)
- enriched categories are monads in $\mathcal{V}\text{-Mat}$.

Monads and monad morphisms

Categories, operads, multicategories and T-multicategories **can all be seen as monads in appropriate bicategories.**

However, morphisms between them are not monad maps.

- internal categories (with a discrete object of objects) are monads in $\mathbf{Span}(\mathcal{V})$ (resp. in $\mathbf{Span}_d(\mathcal{V})$)
- enriched categories are monads in $\mathcal{V}\text{-Mat}$.

BUT:

- monad morphisms between monads on the bicategories $\mathbf{Span}_d(\mathcal{V})$ and $\mathcal{V}\text{-Mat}$ are not functors of categories internal in \mathcal{V} , resp. of categories enriched over \mathcal{V} .

The double category $\mathbf{Mnd}(\mathbb{D})$ of (double) monads

To address this issue [Fiore, Gambino, Kock: “Monads in double categories” (2010)] defined the **double category** $\mathbf{Mnd}(\mathbb{D})$ **of monads**:

The double category $\mathbf{Mnd}(\mathbb{D})$ of (double) monads

To address this issue [Fiore, Gambino, Kock: “Monads in double categories” (2010)] defined the **double category** $\mathbf{Mnd}(\mathbb{D})$ **of monads**:

- (double) monads
- vertical monad maps
- horizontal monad maps
- squares (2-cells)

The double category $\mathbf{Mnd}(\mathbb{D})$ of (double) monads

To address this issue [Fiore, Gambino, Kock: “Monads in double categories” (2010)] defined the **double category** $\mathbf{Mnd}(\mathbb{D})$ **of monads**:

- (double) monads
- vertical monad maps
- horizontal monad maps
- squares (2-cells)

This allows to **describe mathematical structures** and **morphisms between them**

The double category $\mathbf{Mnd}(\mathbb{D})$ of (double) monads

To address this issue [Fiore, Gambino, Kock: “Monads in double categories” (2010)] defined the **double category** $\mathbf{Mnd}(\mathbb{D})$ **of monads**:

- (double) monads
- **vertical monad maps**
- horizontal monad maps
- **squares (2-cells)**

This allows to **describe mathematical structures** and **morphisms between them** as monads and vertical monad maps in appropriate double categories.

The double category $\mathbf{Mnd}(\mathbb{D})$ of (double) monads

1v-cells:

$$\begin{array}{ccc}
 \begin{array}{c}
 \mathcal{A} \xrightarrow{T} \mathcal{A} \xrightarrow{T} \mathcal{A} \\
 \downarrow u \quad \boxed{\alpha} \quad \downarrow u \quad \boxed{\alpha} \quad \downarrow u \\
 \mathcal{A}' \xrightarrow{T'} \mathcal{A}' \xrightarrow{T'} \mathcal{A}' \\
 \downarrow = \quad \boxed{\mu'} \quad \downarrow = \\
 \mathcal{A}' \xrightarrow{T'} \mathcal{A}'
 \end{array}
 & = &
 \begin{array}{c}
 \mathcal{A} \xrightarrow{T} \mathcal{A} \xrightarrow{T} \mathcal{A} \\
 \downarrow = \quad \boxed{\mu} \quad \downarrow = \\
 \mathcal{A} \xrightarrow{T} \mathcal{A} \\
 \downarrow u \quad \boxed{\alpha} \quad \downarrow u \\
 \mathcal{A}' \xrightarrow{T'} \mathcal{A}'
 \end{array} \\
 \\
 \begin{array}{c}
 \mathcal{A} \xrightarrow{=} \mathcal{A} \\
 \downarrow u \quad \boxed{\text{Id}^u} \quad \downarrow u \\
 \mathcal{A}' \xrightarrow{=} \mathcal{A}' \\
 \downarrow = \quad \boxed{\eta'} \quad \downarrow = \\
 \mathcal{A}' \xrightarrow{T'} \mathcal{A}'
 \end{array}
 & = &
 \begin{array}{c}
 \mathcal{A} \xrightarrow{=} \mathcal{A} \\
 \downarrow = \quad \boxed{\eta} \quad \downarrow = \\
 \mathcal{A} \xrightarrow{T} \mathcal{A} \\
 \downarrow u \quad \boxed{\alpha} \quad \downarrow u \\
 \mathcal{A}' \xrightarrow{T'} \mathcal{A}'
 \end{array}
 \end{array}$$

The double category $\mathbf{Mnd}(\mathbb{D})$ of (double) monads

1v-cells:

$$\begin{array}{ccc}
 \begin{array}{c}
 \mathcal{A} \xrightarrow{T} \mathcal{A} \xrightarrow{T} \mathcal{A} \\
 \downarrow u \quad \boxed{\alpha} \quad \downarrow u \quad \boxed{\alpha} \quad \downarrow u \\
 \mathcal{A}' \xrightarrow{T'} \mathcal{A}' \xrightarrow{T'} \mathcal{A}' \\
 \downarrow = \quad \boxed{\mu'} \quad \downarrow = \\
 \mathcal{A}' \xrightarrow{T'} \mathcal{A}'
 \end{array}
 & = &
 \begin{array}{c}
 \mathcal{A} \xrightarrow{T} \mathcal{A} \xrightarrow{T} \mathcal{A} \\
 \downarrow = \quad \boxed{\mu} \quad \downarrow = \\
 \mathcal{A} \xrightarrow{T} \mathcal{A} \\
 \downarrow u \quad \boxed{\alpha} \quad \downarrow u \\
 \mathcal{A}' \xrightarrow{T'} \mathcal{A}'
 \end{array} \\
 \\
 \begin{array}{c}
 \mathcal{A} \xrightarrow{=} \mathcal{A} \\
 \downarrow u \quad \boxed{\text{Id}^u} \quad \downarrow u \\
 \mathcal{A}' \xrightarrow{=} \mathcal{A}' \\
 \downarrow = \quad \boxed{\eta'} \quad \downarrow = \\
 \mathcal{A}' \xrightarrow{T'} \mathcal{A}'
 \end{array}
 & = &
 \begin{array}{c}
 \mathcal{A} \xrightarrow{=} \mathcal{A} \\
 \downarrow = \quad \boxed{\eta} \quad \downarrow = \\
 \mathcal{A} \xrightarrow{T} \mathcal{A} \\
 \downarrow u \quad \boxed{\alpha} \quad \downarrow u \\
 \mathcal{A}' \xrightarrow{T'} \mathcal{A}'
 \end{array}
 \end{array}$$

\Rightarrow 1v-cells in $\mathbf{Mnd}(\mathbf{Span}(\mathcal{V}))$ are internal functors in \mathcal{V}

The double category $\mathbf{Mnd}(\mathbb{D})$ of (double) monads

1v-cells:

$$\begin{array}{ccc}
 \begin{array}{c}
 \mathcal{A} \xrightarrow{T} \mathcal{A} \xrightarrow{T} \mathcal{A} \\
 \downarrow u \quad \boxed{\alpha} \quad \downarrow u \quad \boxed{\alpha} \quad \downarrow u \\
 \mathcal{A}' \xrightarrow{T'} \mathcal{A}' \xrightarrow{T'} \mathcal{A}' \\
 \downarrow = \quad \boxed{\mu'} \quad \downarrow = \\
 \mathcal{A}' \xrightarrow{T'} \mathcal{A}'
 \end{array}
 & = &
 \begin{array}{c}
 \mathcal{A} \xrightarrow{T} \mathcal{A} \xrightarrow{T} \mathcal{A} \\
 \downarrow = \quad \boxed{\mu} \quad \downarrow = \\
 \mathcal{A} \xrightarrow{T} \mathcal{A} \\
 \downarrow u \quad \boxed{\alpha} \quad \downarrow u \\
 \mathcal{A}' \xrightarrow{T'} \mathcal{A}'
 \end{array} \\
 \\
 \begin{array}{c}
 \mathcal{A} \xrightarrow{=} \mathcal{A} \\
 \downarrow u \quad \boxed{\text{Id}^u} \quad \downarrow u \\
 \mathcal{A}' \xrightarrow{=} \mathcal{A}' \\
 \downarrow = \quad \boxed{\eta'} \quad \downarrow = \\
 \mathcal{A}' \xrightarrow{T'} \mathcal{A}'
 \end{array}
 & = &
 \begin{array}{c}
 \mathcal{A} \xrightarrow{=} \mathcal{A} \\
 \downarrow = \quad \boxed{\eta} \quad \downarrow = \\
 \mathcal{A} \xrightarrow{T} \mathcal{A} \\
 \downarrow u \quad \boxed{\alpha} \quad \downarrow u \\
 \mathcal{A}' \xrightarrow{T'} \mathcal{A}'
 \end{array}
 \end{array}$$

\Rightarrow 1v-cells in $\mathbf{Mnd}(\mathbf{Span}(\mathcal{V}))$ are internal functors in \mathcal{V}

\Rightarrow 1v-cells in $\mathbf{Mnd}(\mathbf{V}\text{-Mat})$ are \mathcal{V} -enriched functors.

Gray tensor product on double categories

Gray tensor product

Gray tensor product gives a (left and right) closed monoidal structure on a category:

Gray tensor product

Gray tensor product gives a (left and right) closed monoidal structure on a category:

$$\mathcal{C}(X \otimes Y, Z) \cong \mathcal{C}(Y, [X, Z])$$

isomorphism natural in $X, Y, Z \in \mathcal{C}$, and $(X \otimes -, [X, -])$ is an adjoint pair of endofunctors on \mathcal{C} .

Gray tensor product

Gray tensor product gives a (left and right) closed monoidal structure on a category:

$$\mathcal{C}(X \otimes Y, Z) \cong \mathcal{C}(Y, [X, Z])$$

isomorphism natural in $X, Y, Z \in \mathcal{C}$, and $(X \otimes -, [X, -])$ is an adjoint pair of endofunctors on \mathcal{C} .

- There is composition: $[B, C] \times [A, B] \rightarrow [A, C]$, a morphism in \mathcal{C} defined via ev 's.

Gray tensor product

Gray tensor product gives a (left and right) closed monoidal structure on a category:

$$\mathcal{C}(X \otimes Y, Z) \cong \mathcal{C}(Y, [X, Z])$$

isomorphism natural in $X, Y, Z \in \mathcal{C}$, and $(X \otimes -, [X, -])$ is an adjoint pair of endofunctors on \mathcal{C} .

- ▶ There is composition: $[B, C] \times [A, B] \rightarrow [A, C]$, a morphism in \mathcal{C} defined via ev 's.

When $\mathcal{C} = 2\text{-Cat}$ (2-categories and 2-functors), the natural candidate for an inner hom is the 2-category $\mathbf{Fun}(\mathcal{A}, \mathcal{B})$ for 2-categories \mathcal{A}, \mathcal{B} (2-functors, lax natural transformations, modifications).

Gray tensor product

Gray tensor product gives a (left and right) closed monoidal structure on a category:

$$\mathcal{C}(X \otimes Y, Z) \cong \mathcal{C}(Y, [X, Z])$$

isomorphism natural in $X, Y, Z \in \mathcal{C}$, and $(X \otimes -, [X, -])$ is an adjoint pair of endofunctors on \mathcal{C} .

- There is composition: $[B, C] \times [A, B] \rightarrow [A, C]$, a morphism in \mathcal{C} defined via ev 's.

When $\mathcal{C} = 2\text{-Cat}$ (2-categories and 2-functors), the natural candidate for an inner hom is the 2-category $\mathbf{Fun}(\mathcal{A}, \mathcal{B})$ for 2-categories \mathcal{A}, \mathcal{B} (2-functors, lax natural transformations, modifications).

BUT:

the composition $c : \mathbf{Fun}(\mathcal{B}, \mathcal{C}) \times \mathbf{Fun}(\mathcal{A}, \mathcal{B}) \rightarrow \mathbf{Fun}(\mathcal{A}, \mathcal{C})$ isn't a 2-functor:

Gray tensor product

Gray tensor product gives a (left and right) closed monoidal structure on a category:

$$\mathcal{C}(X \otimes Y, Z) \cong \mathcal{C}(Y, [X, Z])$$

isomorphism natural in $X, Y, Z \in \mathcal{C}$, and $(X \otimes -, [X, -])$ is an adjoint pair of endofunctors on \mathcal{C} .

- There is composition: $[B, C] \times [A, B] \rightarrow [A, C]$, a morphism in \mathcal{C} defined via ev 's.

When $\mathcal{C} = 2\text{-Cat}$ (2-categories and 2-functors), the natural candidate for an inner hom is the 2-category $\mathbf{Fun}(\mathcal{A}, \mathcal{B})$ for 2-categories \mathcal{A}, \mathcal{B} (2-functors, lax natural transformations, modifications).

BUT:

the composition $c : \mathbf{Fun}(\mathcal{B}, \mathcal{C}) \times \mathbf{Fun}(\mathcal{A}, \mathcal{B}) \rightarrow \mathbf{Fun}(\mathcal{A}, \mathcal{C})$ isn't a 2-functor: although 1-cells (lax transformations) compose nicely, the 2-functor condition on them for c translates into the "interchange law": $\frac{\beta}{\beta'} \circ \frac{\alpha}{\alpha'} = \frac{\beta \circ \alpha}{\beta' \circ \alpha'}$

Gray tensor product

Gray tensor product gives a (left and right) closed monoidal structure on a category:

$$\mathcal{C}(X \otimes Y, Z) \cong \mathcal{C}(Y, [X, Z])$$

isomorphism natural in $X, Y, Z \in \mathcal{C}$, and $(X \otimes -, [X, -])$ is an adjoint pair of endofunctors on \mathcal{C} .

- ▶ There is composition: $[B, C] \times [A, B] \rightarrow [A, C]$, a morphism in \mathcal{C} defined via ev 's.

When $\mathcal{C} = 2\text{-Cat}$ (2-categories and 2-functors), the natural candidate for an inner hom is the 2-category $\mathbf{Fun}(\mathcal{A}, \mathcal{B})$ for 2-categories \mathcal{A}, \mathcal{B} (2-functors, lax natural transformations, modifications).

BUT:

the composition $c : \mathbf{Fun}(\mathcal{B}, \mathcal{C}) \times \mathbf{Fun}(\mathcal{A}, \mathcal{B}) \rightarrow \mathbf{Fun}(\mathcal{A}, \mathcal{C})$ isn't a 2-functor: although 1-cells (lax transformations) compose nicely, the 2-functor condition on them for c translates into the "interchange law": $\frac{\beta}{\beta'} \circ \frac{\alpha}{\alpha'} = \frac{\beta \circ \alpha}{\beta' \circ \alpha'}$

and the interchange does not hold in general on lax transformations (strictness of transformations is needed).

Gray tensor product on 2-Cat by generators and relations

One looks for a 2-category $\mathcal{A} \otimes \mathcal{B}$ s.t.:

$$2\text{-Cat}(\mathcal{A} \otimes \mathcal{B}, \mathcal{C}) \cong 2\text{-Cat}(\mathcal{B}, \text{Fun}(\mathcal{A}, \mathcal{C})).$$

Gray tensor product on 2-Cat by generators and relations

One looks for a 2-category $\mathcal{A} \otimes \mathcal{B}$ s.t.:

$$2\text{-Cat}(\mathcal{A} \otimes \mathcal{B}, \mathcal{C}) \cong 2\text{-Cat}(\mathcal{B}, \text{Fun}(\mathcal{A}, \mathcal{C})).$$

- A description of $\mathcal{A} \otimes \mathcal{B}$ is obtained by seeing what a 2-functor $F : \mathcal{B} \rightarrow \text{Fun}(\mathcal{A}, \mathcal{C})$ is.

Gray tensor product on 2-Cat by generators and relations

One looks for a 2-category $\mathcal{A} \otimes \mathcal{B}$ s.t.:

$$2\text{-Cat}(\mathcal{A} \otimes \mathcal{B}, \mathcal{C}) \cong 2\text{-Cat}(\mathcal{B}, \text{Fun}(\mathcal{A}, \mathcal{C})).$$

- ▶ A description of $\mathcal{A} \otimes \mathcal{B}$ is obtained by seeing what a 2-functor $F : \mathcal{B} \rightarrow \text{Fun}(\mathcal{A}, \mathcal{C})$ is.
- ▶ One obtains “quasi-functor of two variables” $H : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$ defined by relations among $F(B)(A)$, $A \in \mathcal{A}$, $B \in \mathcal{B}$,

Gray tensor product on 2-Cat by generators and relations

One looks for a 2-category $\mathcal{A} \otimes \mathcal{B}$ s.t.:

$$2\text{-Cat}(\mathcal{A} \otimes \mathcal{B}, \mathcal{C}) \cong 2\text{-Cat}(\mathcal{B}, \text{Fun}(\mathcal{A}, \mathcal{C})).$$

- ▶ A description of $\mathcal{A} \otimes \mathcal{B}$ is obtained by seeing what a 2-functor $F : \mathcal{B} \rightarrow \text{Fun}(\mathcal{A}, \mathcal{C})$ is.
- ▶ One obtains “quasi-functor of two variables” $H : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$ defined by relations among $F(B)(A)$, $A \in \mathcal{A}$, $B \in \mathcal{B}$, and concludes the relations holding in $\mathcal{A} \otimes \mathcal{B}$.

Gray tensor product on 2-Cat by generators and relations

One looks for a 2-category $\mathcal{A} \otimes \mathcal{B}$ s.t.:

$$2\text{-Cat}(\mathcal{A} \otimes \mathcal{B}, \mathcal{C}) \cong 2\text{-Cat}(\mathcal{B}, \text{Fun}(\mathcal{A}, \mathcal{C})).$$

- ▶ A description of $\mathcal{A} \otimes \mathcal{B}$ is obtained by seeing what a 2-functor $F : \mathcal{B} \rightarrow \text{Fun}(\mathcal{A}, \mathcal{C})$ is.
- ▶ One obtains “quasi-functor of two variables” $H : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$ defined by relations among $F(B)(A)$, $A \in \mathcal{A}$, $B \in \mathcal{B}$, and concludes the relations holding in $\mathcal{A} \otimes \mathcal{B}$.

The relation that differs from what holds in the Cartesian product is:

$$(f \otimes B')(A \otimes g) \not\Rightarrow (A' \otimes g)(f \otimes B).$$

Gray tensor product on 2-Cat by generators and relations

One looks for a 2-category $\mathcal{A} \otimes \mathcal{B}$ s.t.:

$$2\text{-Cat}(\mathcal{A} \otimes \mathcal{B}, \mathcal{C}) \cong 2\text{-Cat}(\mathcal{B}, \text{Fun}(\mathcal{A}, \mathcal{C})).$$

- ▶ A description of $\mathcal{A} \otimes \mathcal{B}$ is obtained by seeing what a 2-functor $F : \mathcal{B} \rightarrow \text{Fun}(\mathcal{A}, \mathcal{C})$ is.
- ▶ One obtains “quasi-functor of two variables” $H : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$ defined by relations among $F(B)(A)$, $A \in \mathcal{A}$, $B \in \mathcal{B}$, and concludes the relations holding in $\mathcal{A} \otimes \mathcal{B}$.

The relation that differs from what holds in the Cartesian product is:

$$(f \otimes B')(A \otimes g) \not\Rightarrow (A' \otimes g)(f \otimes B).$$

Gray proved that $\mathcal{A} \otimes \mathcal{B}$ yields a monoidal product on 2-Cat.

Gray tensor product for (strict-strict) double categories

In

[G. Böhm: “The Gray Monoidal Product of Double Categories” (2020)]

monoidal structure in $(\mathbf{Db}^{st}_{st}, \otimes)$ is obtained from:

$$\mathbf{Db}^{st}_{st}(\mathbb{A} \otimes \mathbb{B}, \mathbb{C}) \cong \mathbf{Db}^{st}_{st}(\mathbb{A}, [\mathbb{B}, \mathbb{C}]).$$

Gray tensor product for (strict-strict) double categories

In

[G. Böhm: “The Gray Monoidal Product of Double Categories” (2020)]

monoidal structure in $(\mathbf{Db}^{st}_{st}, \otimes)$ is obtained from:

$$\mathbf{Db}^{st}_{st}(\mathbb{A} \otimes \mathbb{B}, \mathbb{C}) \cong \mathbf{Db}^{st}_{st}(\mathbb{A}, [\mathbb{B}, \mathbb{C}]).$$

In

[B. Femić: “Enrichment and internalization in tricategories, the case of tensor categories and alternative notion to intercategories”]:

- ▶ we characterized a double functor $F : \mathbb{A} \rightarrow [\mathbb{B}, \mathbb{C}]$,

Gray tensor product for (strict-strict) double categories

In

[G. Böhm: “The Gray Monoidal Product of Double Categories” (2020)]

monoidal structure in $(\mathbf{Db}^{st}_{st}, \otimes)$ is obtained from:

$$\mathbf{Db}^{st}_{st}(\mathbb{A} \otimes \mathbb{B}, \mathbb{C}) \cong \mathbf{Db}^{st}_{st}(\mathbb{A}, [\mathbb{B}, \mathbb{C}]).$$

In

[B. Femić: “Enrichment and internalization in tricategories, the case of tensor categories and alternative notion to intercategories”]:

- ▶ we characterized a double functor $F : \mathbb{A} \rightarrow [\mathbb{B}, \mathbb{C}]$,
- ▶ we read off the structure of the double cat. $F(\mathbb{A})(\mathbb{B})$

Gray tensor product for (strict-strict) double categories

In

[G. Böhm: “The Gray Monoidal Product of Double Categories” (2020)]

monoidal structure in $(\mathbf{Db}^{st}_{st}, \otimes)$ is obtained from:

$$\mathbf{Db}^{st}_{st}(\mathbb{A} \otimes \mathbb{B}, \mathbb{C}) \cong \mathbf{Db}^{st}_{st}(\mathbb{A}, [\mathbb{B}, \mathbb{C}]).$$

In

[B. Femić: “Enrichment and internalization in tricategories, the case of tensor categories and alternative notion to intercategories”]:

- ▶ we characterized a double functor $F : \mathbb{A} \rightarrow [\mathbb{B}, \mathbb{C}]$,
- ▶ we read off the structure of the double cat. $F(\mathbb{A})(\mathbb{B})$
- ▶ and described $\mathbb{A} \otimes \mathbb{B}$ by relations.

Bifunctor Theorem for (lax-hop) double categories

Towards the Gray tensor product on (strict-lax) double categories

[B. Femić: "Bifunctor Theorem and Gray monoidal structure for double categories with lax double functors"]

We define $[[\mathbb{A}, \mathbb{B}]]$ (as a candidate for inner-hom in DbI_{IX}^{st})

- 0: **lax** double functors
- 1v: vertical **lax** transf.
- 1h: horizontal **oplax** transf.
- modifications

Towards the Gray tensor product on (strict-lax) double categories

[B. Femić: "Bifunctor Theorem and Gray monoidal structure for double categories with lax double functors"]

We define $\llbracket \mathbb{A}, \mathbb{B} \rrbracket$ (as a candidate for inner-hom in DbI_{lx}^{st})

- 0: **lax** double functors
- 1v: vertical **lax** transf.
- 1h: horizontal **oplax** transf.
- modifications

Problem: $\llbracket -, - \rrbracket : (DbI_{lx}^{st})^{op} \times DbI_{ps}^{st} \rightarrow DbI_{lx}^{st}$

Towards the Gray tensor product on (strict-lax) double categories

[B. Femić: "Bifunctor Theorem and Gray monoidal structure for double categories with lax double functors"]

We define $\llbracket A, B \rrbracket$ (as a candidate for inner-hom in DbI_{lx}^{st})

- 0: **lax** double functors
- 1v: vertical **lax** transf.
- 1h: horizontal **oplax** transf.
- modifications

Problem: $\llbracket -, - \rrbracket : (DbI_{lx}^{st})^{op} \times DbI_{ps}^{st} \rightarrow DbI_{lx}^{st}$

$$(F, G) \mapsto \llbracket F, G \rrbracket : \llbracket A', B \rrbracket \rightarrow \llbracket A, B' \rrbracket$$

Towards the Gray tensor product on (strict-lax) double categories

[B. Femić: "Bifunctor Theorem and Gray monoidal structure for double categories with lax double functors"]

We define $\llbracket A, B \rrbracket$ (as a candidate for inner-hom in DbI_{lx}^{st})

- 0: **lax** double functors
- 1v: vertical **lax** transf.
- 1h: horizontal **oplax** transf.
- modifications

Problem: $\llbracket -, - \rrbracket : (DbI_{lx}^{st})^{op} \times DbI_{ps}^{st} \rightarrow DbI_{lx}^{st}$

$$(F, G) \mapsto \llbracket F, G \rrbracket : \llbracket A', B \rrbracket \rightarrow \llbracket A, B' \rrbracket$$

- We characterized a lax double functor $F : A \rightarrow \llbracket B, C \rrbracket$,

Towards the Gray tensor product on (strict-lax) double categories

[B. Femić: "Bifunctor Theorem and Gray monoidal structure for double categories with lax double functors"]

We define $\llbracket \mathbb{A}, \mathbb{B} \rrbracket$ (as a candidate for inner-hom in $\mathbf{Dbl}_{\text{lx}}^{\text{st}}$)

- 0: **lax** double functors
- 1v: vertical **lax** transf.
- 1h: horizontal **oplax** transf.
- modifications

Problem: $\llbracket -, - \rrbracket : (\mathbf{Dbl}_{\text{lx}}^{\text{st}})^{\text{op}} \times \mathbf{Dbl}_{\text{ps}}^{\text{st}} \rightarrow \mathbf{Dbl}_{\text{lx}}^{\text{st}}$

$$(F, G) \mapsto \llbracket F, G \rrbracket : \llbracket \mathbb{A}', \mathbb{B} \rrbracket \rightarrow \llbracket \mathbb{A}, \mathbb{B}' \rrbracket$$

- ▶ We characterized a lax double functor $F : \mathbb{A} \rightarrow \llbracket \mathbb{B}, \mathbb{C} \rrbracket$,
- ▶ got to the notion of **lax double quasi-functor** $H : \mathbb{A} \times \mathbb{B} \rightarrow \mathbb{C}$,

Towards the Gray tensor product on (strict-lax) double categories

[B. Femić: "Bifunctor Theorem and Gray monoidal structure for double categories with lax double functors"]

We define $\llbracket \mathbb{A}, \mathbb{B} \rrbracket$ (as a candidate for inner-hom in DbI_{lx}^{st})

- 0: **lax** double functors
- 1v: vertical **lax** transf.
- 1h: horizontal **oplax** transf.
- modifications

Problem: $\llbracket -, - \rrbracket : (DbI_{lx}^{st})^{op} \times DbI_{ps}^{st} \rightarrow DbI_{lx}^{st}$

$$(F, G) \mapsto \llbracket F, G \rrbracket : \llbracket \mathbb{A}', \mathbb{B} \rrbracket \rightarrow \llbracket \mathbb{A}, \mathbb{B}' \rrbracket$$

- ▶ We characterized a lax double functor $F : \mathbb{A} \rightarrow \llbracket \mathbb{B}, \mathbb{C} \rrbracket$,
- ▶ got to the notion of **lax double quasi-functor** $H : \mathbb{A} \times \mathbb{B} \rightarrow \mathbb{C}$,
- ▶ and concluded the relations holding in $\mathbb{A} \otimes \mathbb{B}$.

Some results on quasi-functors and $\mathbb{A} \otimes \mathbb{B}$

The 2-cat. of **lax double quasi-functors**: $q\text{-Lax}_{hop}(\mathbb{A} \times \mathbb{B}, \mathbb{C})$.

Some results on quasi-functors and $\mathbb{A} \otimes \mathbb{B}$

The 2-cat. of **lax double quasi-functors**: $q\text{-Lax}_{hop}(\mathbb{A} \times \mathbb{B}, \mathbb{C})$.

- Isom. of 2-cats: $q\text{-Lax}_{hop}(\mathbb{A} \times \mathbb{B}, \mathbb{C}) \stackrel{*}{\cong} \text{Lax}_{hop}(\mathbb{A}, [\mathbb{B}, \mathbb{C}])$.

Some results on quasi-functors and $\mathbb{A} \otimes \mathbb{B}$

The 2-cat. of **lax double quasi-functors**: $q\text{-Lax}_{hop}(\mathbb{A} \times \mathbb{B}, \mathbb{C})$.

- Isom. of 2-cats: $q\text{-Lax}_{hop}(\mathbb{A} \times \mathbb{B}, \mathbb{C}) \stackrel{*}{\cong} \text{Lax}_{hop}(\mathbb{A}, [\mathbb{B}, \mathbb{C}])$.
- Isom. of 2-cats: $q\text{-Lax}_{hop}(\mathbb{A} \times \mathbb{B}, \mathbb{C}) \cong \text{Lax}_{hop}(\mathbb{A} \otimes \mathbb{B}, \mathbb{C})$.

Some results on quasi-functors and $A \otimes B$

The 2-cat. of **lax double quasi-functors**: $q\text{-Lax}_{hop}(A \times B, C)$.

- Isom. of 2-cats: $q\text{-Lax}_{hop}(A \times B, C) \cong^* \text{Lax}_{hop}(A, [B, C])$.
- Isom. of 2-cats: $q\text{-Lax}_{hop}(A \times B, C) \cong \text{Lax}_{hop}(A \otimes B, C)$.

Consequently: $\text{Lax}_{hop}(A \otimes B, C) \cong \text{Lax}_{hop}(A, [B, C])$.

Some results on quasi-functors and $A \otimes B$

The 2-cat. of **lax double quasi-functors**: $q\text{-Lax}_{hop}(A \times B, C)$.

- Isom. of 2-cats: $q\text{-Lax}_{hop}(A \times B, C) \cong^* \text{Lax}_{hop}(A, [B, C])$.
- Isom. of 2-cats: $q\text{-Lax}_{hop}(A \times B, C) \cong \text{Lax}_{hop}(A \otimes B, C)$.

Consequently: $\text{Lax}_{hop}(A \otimes B, C) \cong \text{Lax}_{hop}(A, [B, C])$.

► Hence, there is an isomorphism of sets:

$$DbI_{lx}^{st}(A \otimes B, C) \cong DbI_{lx}^{st}(A, [B, C]).$$

Bifunctor Theorem: 1- and 2-categories

[MacLane:]

Given functors $L_C: \mathcal{B} \rightarrow \mathcal{D}$ and $M_B: \mathcal{C} \rightarrow \mathcal{D}$ so that
 $L_C(B) = M_B(C), \forall B \in \mathcal{B}, C \in \mathcal{C}.$

Bifunctor Theorem: 1- and 2-categories

[MacLane:]

Given functors $L_C: \mathcal{B} \rightarrow \mathcal{D}$ and $M_B: \mathcal{C} \rightarrow \mathcal{D}$ so that

$$L_C(B) = M_B(C), \forall B \in \mathcal{B}, C \in \mathcal{C}.$$

Then \exists a bifunctor $P: \mathcal{B} \times \mathcal{C} \rightarrow \mathcal{D}$ / $L_C = P(-, C)$ and $M_B = P(B, -)$

$$\iff L_{C'}(f)M_B(g) = M_{B'}(g)L_C(f), \forall f: B \rightarrow B' \text{ in } \mathcal{B}, g: C \rightarrow C' \text{ in } \mathcal{C}.$$

Bifunctor Theorem: 1- and 2-categories

[MacLane:]

Given functors $L_C: \mathcal{B} \rightarrow \mathcal{D}$ and $M_B: \mathcal{C} \rightarrow \mathcal{D}$ so that

$$L_C(B) = M_B(C), \forall B \in \mathcal{B}, C \in \mathcal{C}.$$

Then \exists a bifunctor $P: \mathcal{B} \times \mathcal{C} \rightarrow \mathcal{D}$ / $L_C = P(-, C)$ and $M_B = P(B, -)$

$$\iff L_{C'}(f)M_B(g) = M_{B'}(g)L_C(f), \forall f: B \rightarrow B' \text{ in } \mathcal{B}, g: C \rightarrow C' \text{ in } \mathcal{C}.$$

In this case, $P(B, C) = L_C(B) = M_B(C)$ and $P(f, g) = M_{B'}(g)L_C(f)$.

Bifunctor Theorem: 1- and 2-categories

[MacLane:]

Given functors $L_C: \mathcal{B} \rightarrow \mathcal{D}$ and $M_B: \mathcal{C} \rightarrow \mathcal{D}$ so that

$$L_C(B) = M_B(C), \forall B \in \mathcal{B}, C \in \mathcal{C}.$$

Then \exists a bifunctor $P: \mathcal{B} \times \mathcal{C} \rightarrow \mathcal{D}$ / $L_C = P(-, C)$ and $M_B = P(B, -)$

$$\iff L_{C'}(f)M_B(g) = M_{B'}(g)L_C(f), \forall f: B \rightarrow B' \text{ in } \mathcal{B}, g: C \rightarrow C' \text{ in } \mathcal{C}.$$

In this case, $P(B, C) = L_C(B) = M_B(C)$ and $P(f, g) = M_{B'}(g)L_C(f)$.

[P.F. Faul, G. Manuell, J. Siqueira: “2-Dimensional Bifunctor Theorems and Distributive laws”, (2021)]

Let $\sigma \in \text{Dist}(\mathcal{B}, \mathcal{C}, \mathcal{D})$ be a distributive law between families of lax functors $L_C: \mathcal{B} \rightarrow \mathcal{D}$ and $M_B: \mathcal{C} \rightarrow \mathcal{D} \forall B \in \mathcal{B}, C \in \mathcal{C}$.

Bifunctor Theorem: 1- and 2-categories

[MacLane:]

Given functors $L_C: \mathcal{B} \rightarrow \mathcal{D}$ and $M_B: \mathcal{C} \rightarrow \mathcal{D}$ so that

$$L_C(B) = M_B(C), \forall B \in \mathcal{B}, C \in \mathcal{C}.$$

Then \exists a bifunctor $P: \mathcal{B} \times \mathcal{C} \rightarrow \mathcal{D}$ / $L_C = P(-, C)$ and $M_B = P(B, -)$

$$\iff L_{C'}(f)M_B(g) = M_{B'}(g)L_C(f), \forall f: B \rightarrow B' \text{ in } \mathcal{B}, g: C \rightarrow C' \text{ in } \mathcal{C}.$$

In this case, $P(B, C) = L_C(B) = M_B(C)$ and $P(f, g) = M_{B'}(g)L_C(f)$.

[P.F. Faul, G. Manuell, J. Siqueira: “2-Dimensional Bifunctor Theorems and Distributive laws”, (2021)]

Let $\sigma \in \text{Dist}(\mathcal{B}, \mathcal{C}, \mathcal{D})$ be a distributive law between families of lax functors

$$L_C: \mathcal{B} \rightarrow \mathcal{D} \text{ and } M_B: \mathcal{C} \rightarrow \mathcal{D} \quad \forall B \in \mathcal{B}, C \in \mathcal{C}.$$

Then there is a lax bifunctor $P: \mathcal{B} \times \mathcal{C} \rightarrow \mathcal{D}$ with

$$P(B, C) = L_C(B) = M_B(C), \quad P(f, g) = M_{B'}(g)L_C(f) \text{ and}$$

$$P(\alpha, \beta) = [L_C(\alpha) | M_{B'}(\beta)] \text{ on 0-, 1- and 2-cells, resp.,}$$

Bifunctor Theorem: 1- and 2-categories

[MacLane:]

Given functors $L_C: \mathcal{B} \rightarrow \mathcal{D}$ and $M_B: \mathcal{C} \rightarrow \mathcal{D}$ so that

$$L_C(B) = M_B(C), \forall B \in \mathcal{B}, C \in \mathcal{C}.$$

Then \exists a bifunctor $P: \mathcal{B} \times \mathcal{C} \rightarrow \mathcal{D}$ / $L_C = P(-, C)$ and $M_B = P(B, -)$

$$\iff L_{C'}(f)M_B(g) = M_{B'}(g)L_C(f), \forall f: B \rightarrow B' \text{ in } \mathcal{B}, g: C \rightarrow C' \text{ in } \mathcal{C}.$$

In this case, $P(B, C) = L_C(B) = M_B(C)$ and $P(f, g) = M_{B'}(g)L_C(f)$.

[P.F. Faul, G. Manuell, J. Siqueira: “2-Dimensional Bifunctor Theorems and Distributive laws”, (2021)]

Let $\sigma \in \text{Dist}(\mathcal{B}, \mathcal{C}, \mathcal{D})$ be a distributive law between families of lax functors

$$L_C: \mathcal{B} \rightarrow \mathcal{D} \text{ and } M_B: \mathcal{C} \rightarrow \mathcal{D} \quad \forall B \in \mathcal{B}, C \in \mathcal{C}.$$

Then there is a lax bifunctor $P: \mathcal{B} \times \mathcal{C} \rightarrow \mathcal{D}$ with

$$P(B, C) = L_C(B) = M_B(C), \quad P(f, g) = M_{B'}(g)L_C(f) \text{ and}$$

$$P(\alpha, \beta) = [L_C(\alpha) | M_{B'}(\beta)] \text{ on 0-, 1- and 2-cells, resp.,}$$

with given compositor (in terms of σ) and unitor.

Bifunctor Theorem: 1- and 2-categories

[MacLane:]

Given functors $L_C: \mathcal{B} \rightarrow \mathcal{D}$ and $M_B: \mathcal{C} \rightarrow \mathcal{D}$ so that

$$L_C(B) = M_B(C), \forall B \in \mathcal{B}, C \in \mathcal{C}.$$

Then \exists a bifunctor $P: \mathcal{B} \times \mathcal{C} \rightarrow \mathcal{D}$ / $L_C = P(-, C)$ and $M_B = P(B, -)$

$$\iff L_{C'}(f)M_B(g) = M_{B'}(g)L_C(f), \forall f: B \rightarrow B' \text{ in } \mathcal{B}, g: C \rightarrow C' \text{ in } \mathcal{C}.$$

In this case, $P(B, C) = L_C(B) = M_B(C)$ and $P(f, g) = M_{B'}(g)L_C(f)$.

[P.F. Faul, G. Manuell, J. Siqueira: “2-Dimensional Bifunctor Theorems and Distributive laws”, (2021)]

Let $\sigma \in \text{Dist}(\mathcal{B}, \mathcal{C}, \mathcal{D})$ be a distributive law between families of lax functors $L_C: \mathcal{B} \rightarrow \mathcal{D}$ and $M_B: \mathcal{C} \rightarrow \mathcal{D} \forall B \in \mathcal{B}, C \in \mathcal{C}$.

Then there is a lax bifunctor $P: \mathcal{B} \times \mathcal{C} \rightarrow \mathcal{D}$ with

$$P(B, C) = L_C(B) = M_B(C), \quad P(f, g) = M_{B'}(g)L_C(f) \text{ and}$$

$$P(\alpha, \beta) = [L_C(\alpha) | M_{B'}(\beta)] \text{ on 0-, 1- and 2-cells, resp.,}$$

with given compositor (in terms of σ) and unitor.

► 2-functor $K: \text{Dist}(\mathcal{B}, \mathcal{C}, \mathcal{D}) \rightarrow \text{Lax}_{op}(\mathcal{B} \times \mathcal{C}, \mathcal{D})$

Bifunctor Theorem: 1- and 2-categories

[MacLane:]

Given functors $L_C: \mathcal{B} \rightarrow \mathcal{D}$ and $M_B: \mathcal{C} \rightarrow \mathcal{D}$ so that

$$L_C(B) = M_B(C), \forall B \in \mathcal{B}, C \in \mathcal{C}.$$

Then \exists a bifunctor $P: \mathcal{B} \times \mathcal{C} \rightarrow \mathcal{D}$ / $L_C = P(-, C)$ and $M_B = P(B, -)$

$$\iff L_{C'}(f)M_B(g) = M_{B'}(g)L_C(f), \forall f: B \rightarrow B' \text{ in } \mathcal{B}, g: C \rightarrow C' \text{ in } \mathcal{C}.$$

In this case, $P(B, C) = L_C(B) = M_B(C)$ and $P(f, g) = M_{B'}(g)L_C(f)$.

[P.F. Faul, G. Manuell, J. Siqueira: “2-Dimensional Bifunctor Theorems and Distributive laws”, (2021)]

Let $\sigma \in \text{Dist}(\mathcal{B}, \mathcal{C}, \mathcal{D})$ be a distributive law between families of lax functors

$$L_C: \mathcal{B} \rightarrow \mathcal{D} \text{ and } M_B: \mathcal{C} \rightarrow \mathcal{D} \quad \forall B \in \mathcal{B}, C \in \mathcal{C}.$$

Then there is a lax bifunctor $P: \mathcal{B} \times \mathcal{C} \rightarrow \mathcal{D}$ with

$$P(B, C) = L_C(B) = M_B(C), \quad P(f, g) = M_{B'}(g)L_C(f) \text{ and}$$

$$P(\alpha, \beta) = [L_C(\alpha) | M_{B'}(\beta)] \text{ on 0-, 1- and 2-cells, resp.,}$$

with given compositor (in terms of σ) and unitor.

- ▶ 2-functor $K: \text{Dist}(\mathcal{B}, \mathcal{C}, \mathcal{D}) \rightarrow \text{Lax}_{op}(\mathcal{B} \times \mathcal{C}, \mathcal{D})$
- ▶ K restricts to a 2-equivalence.

Bifunctor Theorem: double categories

$$\mathcal{F} : q\text{-}\mathbf{Lax}_{hop}^{\text{ns}}(\mathbb{A} \times \mathbb{B}, \mathbb{C}) \rightarrow \mathbf{Lax}_{hop}(\mathbb{A} \times \mathbb{B}, \mathbb{C})$$
$$H \mapsto (P, \gamma, \iota)$$

Bifunctor Theorem: double categories

$$\mathcal{F} : q\text{-Lax}_{hop}^{ns}(\mathbb{A} \times \mathbb{B}, \mathbb{C}) \rightarrow \text{Lax}_{hop}(\mathbb{A} \times \mathbb{B}, \mathbb{C})$$

$$H \mapsto (P, \gamma, \iota)$$

full sub-2-categories:

$q\text{-Lax}_{hop}^{ns-u}(\mathbb{A} \times \mathbb{B}, \mathbb{C})$: *unital* lax double quasi-functors (meaning that $(-, A)$

and $(B, -)$ are unital, i.e. ι^A, ι^B invertible)

$\text{Lax}_{hop}^{u-d}(\mathbb{A} \times \mathbb{B}, \mathbb{C})$: *unital and decomposable* lax double functors

(ι and $\gamma_{(1_A, g), (f, 1_{B'})}$ are invertible).

Bifunctor Theorem: double categories

$$\mathcal{F} : q\text{-Lax}_{hop}^{ns}(\mathbb{A} \times \mathbb{B}, \mathbb{C}) \rightarrow \text{Lax}_{hop}(\mathbb{A} \times \mathbb{B}, \mathbb{C})$$

$$H \mapsto (P, \gamma, \iota)$$

full sub-2-categories:

$q\text{-Lax}_{hop}^{ns-u}(\mathbb{A} \times \mathbb{B}, \mathbb{C})$: *unital* lax double quasi-functors (meaning that $(-, A)$

and $(B, -)$ are unital, i.e. ι^A, ι^B invertible)

$\text{Lax}_{hop}^{u-d}(\mathbb{A} \times \mathbb{B}, \mathbb{C})$: *unital and decomposable* lax double functors

(ι and $\gamma_{(1_A, g), (f, 1_{B'})}$ are invertible).

\mathcal{F} restricts to 2-equivalences:

$$\mathcal{F}' : q\text{-Lax}_{hop}^{ns-u}(\mathbb{A} \times \mathbb{B}, \mathbb{C}) \xrightarrow{\cong} \text{Lax}_{hop}^{u-d}(\mathbb{A} \times \mathbb{B}, \mathbb{C}).$$

Bifunctor Theorem: double categories

$$\mathcal{F} : q\text{-Lax}_{hop}^{ns}(\mathbb{A} \times \mathbb{B}, \mathbb{C}) \rightarrow \text{Lax}_{hop}(\mathbb{A} \times \mathbb{B}, \mathbb{C})$$

$$H \mapsto (P, \gamma, \iota)$$

full sub-2-categories:

$q\text{-Lax}_{hop}^{ns-u}(\mathbb{A} \times \mathbb{B}, \mathbb{C})$: *unital* lax double quasi-functors (meaning that $(-, A)$

and $(B, -)$ are unital, i.e. ι^A, ι^B invertible)

$\text{Lax}_{hop}^{u-d}(\mathbb{A} \times \mathbb{B}, \mathbb{C})$: *unital and decomposable* lax double functors

(ι and $\gamma_{(1_A, g), (f, 1_{B'})}$ are invertible).

\mathcal{F} restricts to 2-equivalences:

$$\mathcal{F}' : q\text{-Lax}_{hop}^{ns-u}(\mathbb{A} \times \mathbb{B}, \mathbb{C}) \xrightarrow{\cong} \text{Lax}_{hop}^{u-d}(\mathbb{A} \times \mathbb{B}, \mathbb{C}).$$

$$\mathcal{F}'' : q\text{-Ps}_{hop}^{ns}(\mathbb{A} \times \mathbb{B}, \mathbb{C}) \xrightarrow{\cong} \text{Ps}_{hop}(\mathbb{A} \times \mathbb{B}, \mathbb{C}).$$

“(Un)currying” 2-functor

- $q\text{-Lax}_{hop}^{ns}(\mathbb{A} \times \mathbb{B}, \mathbb{C}) \stackrel{*}{\cong} \text{Lax}_{hop}(\mathbb{A}, [\mathbb{B}, \mathbb{C}]^{ns});$

“(Un)currying” 2-functor

- $q\text{-Lax}_{hop}^{ns}(\mathbb{A} \times \mathbb{B}, \mathbb{C}) \overset{*}{\cong} \text{Lax}_{hop}(\mathbb{A}, [\mathbb{B}, \mathbb{C}]^{ns})$;
composing with \mathcal{F} yields an **uncurrying** 2-functor:

$$\text{Lax}_{hop}(\mathbb{A}, [\mathbb{B}, \mathbb{C}]^{ns}) \rightarrow \text{Lax}_{hop}(\mathbb{A} \times \mathbb{B}, \mathbb{C})$$

“(Un)currying” 2-functor

- $q\text{-Lax}_{hop}^{ns}(\mathbb{A} \times \mathbb{B}, \mathbb{C}) \overset{*}{\cong} \text{Lax}_{hop}(\mathbb{A}, [\mathbb{B}, \mathbb{C}]^{ns});$
composing with \mathcal{F} yields an **uncurrying** 2-functor:

$$\text{Lax}_{hop}(\mathbb{A}, [\mathbb{B}, \mathbb{C}]^{ns}) \rightarrow \text{Lax}_{hop}(\mathbb{A} \times \mathbb{B}, \mathbb{C})$$

- $q\text{-Lax}_{hop}^{ns-u}(\mathbb{A} \times \mathbb{B}, \mathbb{C}) \cong \text{Lax}_{hop}^u(\mathbb{A}, [\mathbb{B}, \mathbb{C}]^{ns-u});$

“(Un)currying” 2-functor

- $q\text{-Lax}_{hop}^{ns}(\mathbb{A} \times \mathbb{B}, \mathbb{C}) \cong \text{Lax}_{hop}^*(\mathbb{A}, [\mathbb{B}, \mathbb{C}]^{ns});$
composing with \mathcal{F} yields an **uncurrying** 2-functor:

$$\text{Lax}_{hop}(\mathbb{A}, [\mathbb{B}, \mathbb{C}]^{ns}) \rightarrow \text{Lax}_{hop}(\mathbb{A} \times \mathbb{B}, \mathbb{C})$$

- $q\text{-Lax}_{hop}^{ns-u}(\mathbb{A} \times \mathbb{B}, \mathbb{C}) \cong \text{Lax}_{hop}^u(\mathbb{A}, [\mathbb{B}, \mathbb{C}]^{ns-u});$
composing with \mathcal{F}' one gets a **currying** 2-functor which is a 2-equivalence:

$$\text{Lax}_{hop}^{u-d}(\mathbb{A} \times \mathbb{B}, \mathbb{C}) \simeq \text{Lax}_{hop}^u(\mathbb{A}, [\mathbb{B}, \mathbb{C}]^{ns-u})$$

Application to monads in double categories

The following are straightforward:

- ▶ A lax double functor $* \rightarrow \mathbb{D}$ is a monad in \mathbb{D} .

Application to monads in double categories

The followong are straightforward:

- ▶ A lax double functor $* \rightarrow \mathbb{D}$ is a monad in \mathbb{D} .
- ▶ There are 2-category isomorphisms:

$$\mathbf{Lax}_{hop}(*, \mathbb{D}) \cong \mathbf{Mnd}(\mathcal{H}(\mathbb{D}))$$

Application to monads in double categories

The following are straightforward:

- ▶ A lax double functor $* \rightarrow \mathbb{D}$ is a monad in \mathbb{D} .
- ▶ There are 2-category isomorphisms:

$$\text{Lax}_{hop}(*, \mathbb{D}) \cong \text{Mnd}(\mathcal{H}(\mathbb{D}))$$

$$q\text{-Lax}_{hop}(* \times *, \mathbb{D}) \cong \text{Mnd}(\text{Mnd}(\mathcal{H}(\mathbb{D}))).$$

Application to monads in double categories

The following are straightforward:

- ▶ A lax double functor $* \rightarrow \mathbb{D}$ is a monad in \mathbb{D} .
- ▶ There are 2-category isomorphisms:

$$\mathbf{Lax}_{hop}(*, \mathbb{D}) \cong \mathbf{Mnd}(\mathcal{H}(\mathbb{D}))$$

$$q\text{-}\mathbf{Lax}_{hop}(* \times *, \mathbb{D}) \cong \mathbf{Mnd}(\mathbf{Mnd}(\mathcal{H}(\mathbb{D}))).$$

Bifunctor Theorem as a generalization of Beck's result on the composition of monads:

$$\begin{array}{ccc}
 q\text{-}\mathbf{Lax}_{hop}(* \times *, \mathbb{D}) & \xrightarrow{\mathcal{F}} & \mathbf{Lax}_{hop}(*, \mathbb{D}) \\
 \cong \downarrow & & \downarrow \cong \\
 \mathbf{Mnd}(\mathbf{Mnd}(\mathcal{H}(\mathbb{D}))) & \xrightarrow{\text{Comp}(\mathcal{H}(\mathbb{D}))} & \mathbf{Mnd}(\mathcal{H}(\mathbb{D}))
 \end{array}$$