

Bryce Monaco  
CS 455  
February 22, 2018

## Homework 2

1.

a. Computed Sigma Norms:

i. Sigma Norm of  $[3 \ 5] - [7 \ 5] = 6.1245$

ii. Sigma Norm of  $[2 \ 8] - [4 \ 5] = 5.1658$

b. Sigma Norm Function (Full Testing Code Attached):

```
function s = snorm(x, e)
s = (1/e) * (sqrt(1 + (e * (norm (x))^2)) - 1);
end
```

2.

a. Sigma Norm vs Euclidean Norm (Assigned Values):

Input	Sigma Norm	Euclidean
$[3 \ 5] - [7 \ 5] = [-4 \ 0]$	6.1245	4
$[2 \ 8] - [4 \ 5] = [-2 \ 3]$	5.1658	3.6056

b. Sigma Norm vs Euclidean Norm (Test Values):

Input	Sigma Norm	Euclidean
2	1.8322	2
$[0 \ 0]$	0	0
$[3 \ 5 \ 7]$	20.4959	9.1104

## Homework Code:

```
%Constants
e = 0.1; %Epsilon of Sigma Norm

%Qi and Qj for the first pair of vectors
a1 = [3 5];
a2 = [7 5];

%Qi and Qj for the second pair of vectors
b1 = [2 8];
b2 = [4 5];

%i,j are the sigma norms of their respective vectors
i = snorm (a2 - a1, e)
j = snorm (b2 - b1, e)

%k,l are the euclidean norms of their respective vectors
k = norm (a2 - a1)
l = norm (b2 - b1)

function s = snorm(x, e)
s = (1/e) * (sqrt(1 + (e * (norm (x))^2)) - 1);
end
```