

Mathematical Analysis Zorich Comprehensions

Buce-Ithon

15th August, 2025

Contents

1	Multiple Integrals Short Sheet (Calculus)	4
1.1	Development of 1-dim Reimann Integral-Line Integral	4
1.1.1	Introdution (ds)	4
1.1.2	I-Line Integral	4
1.1.3	II-Line Integral	4
1.1.4	Calculate area using II-Line Integral	5
1.1.5	Conditions for curve integrals to be independent of the path	5
1.2	2-dim Multiple Integral	6
1.2.1	Definition	6
1.2.2	Calculate: 2-dim Multiple Integral $\rightarrow$ Iterated Integral . .	6
1.2.3	Green's Formula: 2-dim Multiple Integral $\longleftrightarrow$ II-Line In- tegral	7
1.2.4	Variable Substitution in Double Integrals	7
1.3	Surface Integral	9
1.3.1	Definition and basics	9
2	Multiple Integrals	10
2.1	The Riemann Integral over an n -Dimensional Interval	10
2.1.1	Comprehensions	10
2.1.2	Problems Sets	11



Figure 1: One week, one Zorich

1 Multiple Integrals Short Sheet (Calculus)

1.1 Development of 1-dim Reimann Integral-Line Integral

1.1.1 Intrduction (ds)

Introduction

$$I = \int_C f(x, y) ds, \quad C : r = r(\theta), \quad \theta \in [\theta_1, \theta_2]$$

$$C : \begin{cases} x = r(\theta) \cos \theta \\ y = r(\theta) \sin \theta \end{cases}, \quad f(x, y) = f(r \cos \theta, r \sin \theta)$$

$$ds = \sqrt{x'(\theta)^2 + y'(\theta)^2} d\theta = \sqrt{r'^2 + r^2} d\theta$$

$$I = \int_{\theta_1}^{\theta_2} f(r \cos \theta, r \sin \theta) \sqrt{r'^2 + r^2} d\theta$$

1.1.2 I-Line Integral


Definition

$$I = \int_C f(x, y) ds = \int_{D_t} f(\varphi(t), \psi(t)) \sqrt{\varphi'(t)^2 + \psi'(t)^2} dt$$

A mean value property

$$\rho(x, y) = f(x, y), \quad \int_{C(A, B)} f(x, y) ds = M, \quad \int_{C(C, D)} f(x, y) ds = m$$

$$\frac{\int_{C(C, D)} f(x, y) ds}{\int_{C(A, B)} f(x, y) ds} = \frac{\int_{C(C, D)} C ds}{\int_{C(A, B)} C ds} = \frac{S_{CD}}{S_{AB}}$$



1.1.3 II-Line Integral

Definition

$$I = \int_C f(x, y) dx = [x = \varphi(t), y = \psi(t)] = \int_{D_t} f(\varphi(t), \psi(t)) \varphi'(t) dt$$


$$I = \int_C f(x, y) dy = [x = \varphi(t), y = \psi(t)] = \int_{D_t} f(\varphi(t), \psi(t)) \psi'(t) dt$$

$$I = \int_C P dx + Q dy = [x = \varphi(t), y = \psi(t)] = \int_{D_t} [P\varphi' + Q\psi'] dt$$

1.1.4 Calculate area using II-Line Integral

Calculate

$D = - \int_C y dx = \int_C x dy = \frac{1}{2} \int_C x dy - y dx$, where C is a simple closed curve with positive direction.



1.1.5 Conditions for curve integrals to be independent of the path

Theorem

In simply connected space D and given a path $C : A \rightarrow B$, II-line integral $\int_C P dx + Q dy$ is dependent of the integration path $C : A \rightarrow B$ i.f.f. $\frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x}$.

Proof idea

Necessity " $\Rightarrow$ ":

By studying the differentiation method (partial differentiation with respect to x and y) of $F(x, y) = \int_{(x_0, y_0)}^{(x, y)} P dx + Q dy$ and the primitive function of $P dx + Q dy$ (regarding it as the total differential of some function) (i.e., $\exists \Phi$ such that $P = \frac{\partial \Phi}{\partial x}$, $Q = \frac{\partial \Phi}{\partial y}$ and $\int_C P dx + Q dy = \Phi(x, y)|_{(x_A, y_A)}^{(x_B, y_B)}$), we can conclude that the necessary and sufficient condition for $\int_C P dx + Q dy$ to be path-independent is that $P dx + Q dy (= d\Phi(x))$ is an exact differential. The latter implies $\frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x}$ (necessity).

Sufficiency " $\Leftarrow$ ":

Fix $y \in [c, d]$, integrate $P = \frac{\partial \Phi}{\partial x}$ on $[x_0, x] \subset D$, then $\Phi(x, y) = \int_{x_0}^x P dx + \Phi(x_0, y)$. Also, fix $x = x_0$, integrate $Q = \frac{\partial \Phi}{\partial y}$ on $[y_0, y] \subset D$, then $\Phi(x_0, y) = \int_{y_0}^y Q(x_0, y) dy + \Phi(x_0, y_0)$.


Then $\Phi(x, y) = \int_{x_0}^x P(x, y) dx + \int_{y_0}^y Q(x_0, y) dy + \Phi(x_0, y_0)$ or $\Phi(x, y) = \int_{y_0}^y Q(x, y) dy + \int_{x_0}^x P(x, y_0) dx + \Phi(x_0, y_0)$, which satisfies $\frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x}$.

1.2 2-dim Multiple Integral

1.2.1 Definition

Definition

$$\iint_P f(x, y) dP = \lim_{\lambda = \max_i \{P_i\} \rightarrow 0} \sum_i f(\xi_i, \eta_i) \Delta P_i = V$$



1.2.2 Calculate: 2-dim Multiple Integral $\rightarrow$ Iterated Integral

2-dim Multiple Integral $\rightarrow$ Iterated Integral

Rectangle $P = [a, b] \times [c, d]$:

If $\iint_P f(x, y) dP$ exists and $I_1 = \int_a^b f(x, y) dx$ exists $\forall y \in [c, d]$, then $\iint_P f(x, y) dP = \int_c^d dy \int_a^b f(x, y) dx$.

If $\iint_P f(x, y) dP$ exists and $I_2 = \int_c^d f(x, y) dy$ exists $\forall x \in [a, b]$, then $\iint_P f(x, y) dP = \int_a^b dx \int_c^d f(x, y) dy$.


$\Rightarrow$ If I_1 and I_2 exist, then $\iint_P f(x, y) dP = \int_a^b dx \int_c^d f(x, y) dy = \int_c^d dy \int_a^b f(x, y) dx$.

Curved trapezoid $P = [a, b] \times [y_0(x), Y(x)]$ or $P = [x_0(y), X(y)] \times [c, d]$:

If $\iint_P f(x, y) dP$ exists and $I_2 = \int_{y_0(x)}^{Y(x)} f(x, y) dy$ exists $\forall x \in [a, b]$, then $\iint_P f(x, y) dP = \int_a^b dx \int_{y_0(x)}^{Y(x)} f(x, y) dy$.

If $\iint_P f(x, y) dP$ exists and $I_1 = \int_{x_0(y)}^{X(y)} f(x, y) dx$ exists $\forall y \in [c, d]$, then $\iint_P f(x, y) dP = \int_c^d dy \int_{x_0(y)}^{X(y)} f(x, y) dx$.

$\Rightarrow$ If I_1 and I_2 exist, then $\iint_P f(x, y) dP = \int_a^b dx \int_{y_0(x)}^{Y(x)} f(x, y) dy = \int_c^d dy \int_{x_0(y)}^{X(y)} f(x, y) dx$.



1.2.3 Green's Formula: 2-dim Multiple Integral $\longleftrightarrow$ II-Line Integral

Green's Formula

$$\iint_D \left(\frac{\partial P}{\partial x} - \frac{\partial Q}{\partial y} \right) dx dy = \oint_{C=\partial D} P dy + Q dx$$

or

$$\iint_D \left(\frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} \right) dx dy = \oint_{C=\partial D} P dy - Q dx$$


where D is a simply connected domain in $\mathbb{R}^2$ with boundary $C = \partial D$ and P, Q are continuously differentiable functions on D .

Proof idea

$P, Q, \frac{\partial P}{\partial x}, \frac{\partial Q}{\partial y}$ continuously on D , $\partial D \triangleq C$.

$$\iint_D \frac{\partial P}{\partial x} dx dy = \int_c^d dy \int_{x_0(y)}^{X(y)} \frac{\partial P}{\partial x} dx = \int_c^d [P(X(y), y) - P(x_0(y), y)] dy = \oint_{\partial D} P(x, y) dy$$

$$\iint_D \frac{\partial Q}{\partial y} dx dy = \int_a^b dx \int_{y_0(x)}^{Y(x)} \frac{\partial Q}{\partial y} dy = \int_a^b [Q(x, Y(x)) - Q(x, y_0(x))] dx = - \oint_{\partial D} Q(x, y) dx$$



1.2.4 Variable Substitution in Double Integrals

Intro


$$D = \oint_{S=\partial D} x dy = \left[\begin{matrix} x = x(\xi, \eta) \\ y = y(\xi, \eta) \end{matrix} \right] = \oint_{\Sigma=\partial \Delta} x(\xi, \eta) \frac{\partial y}{\partial \xi} d\xi + x(\xi, \eta) \frac{\partial y}{\partial \eta} d\eta = \iint_{\Delta} \frac{D(x, y)}{D(\xi, \eta)} d\xi d\eta$$

Analysis

As we could see, the essential of $\begin{cases} x = x(\xi, \eta) \\ y = y(\xi, \eta) \end{cases}$ actually is coordinate transform between the x, y -plane and the ξ, η -plane.

$$D = \oint_S x dy = \left[\begin{matrix} x = x(\xi, \eta) \\ y = y(\xi, \eta) \end{matrix}, \begin{matrix} \xi = \xi(t) \\ \eta = \eta(t) \end{matrix} \right] = \int_{\alpha}^{\beta} x(\xi(t), \eta(t)) \left(\frac{\partial y}{\partial \xi} \xi'(t) + \frac{\partial y}{\partial \eta} \eta'(t) \right) dt = \oint_{\Sigma} x(\xi, \eta) \frac{\partial y}{\partial \xi} d\xi + x(\xi, \eta) \frac{\partial y}{\partial \eta} d\eta, \text{ let } x(\xi, \eta) \frac{\partial y}{\partial \xi} \triangleq P(x, y), x(\xi, \eta) \frac{\partial y}{\partial \eta} \triangleq Q(x, y), \text{ then } D = \oint_{\Sigma} P d\xi + Q d\eta = \left(\frac{\partial P}{\partial \eta} - \frac{\partial Q}{\partial \xi}, \text{ using Green's formula} \right) = \iint_{\Delta} \left(\frac{\partial Q}{\partial \xi} - \frac{\partial P}{\partial \eta} \right) d\xi d\eta = \iint_{\Delta} \frac{D(x, y)}{D(\xi, \eta)} d\xi d\eta.$$

Figures for aforementioned analysis



Idea of Analysis:


Using $\begin{cases} x = x(\xi, \eta) = x(\xi(t), \eta(t)) \\ y = y(\xi, \eta) = y(\xi(t), \eta(t)) \end{cases}$ (coordinate transform with parameterization), transform the II-line integration $\oint_S xdy$ (or $-\oint_S ydx$, or $\frac{1}{2} \oint_S xdy - ydx$) on x, y -plane to the II-line integral $\oint_{\Sigma} P(\xi, \eta)d\xi + Q(\xi, \eta)d\eta$, then using Green's formula to transform it to double integral $\iint_{\Delta} (\frac{\partial Q}{\partial \xi} - \frac{\partial P}{\partial \eta})d\xi d\eta$ on ξ, η -plane, following is how we calculate the double integral (often transform it to iterated integral $\int_{D_{\xi}} d\xi \int_{\eta_0(\xi)}^{\eta(\xi)} (\frac{\partial Q}{\partial \xi} - \frac{\partial P}{\partial \eta})d\xi$, or others).

/Shortly, calculate the area of D on x, y plane: coordinate transform with parameterization $\rightarrow$ Green's formula $\rightarrow$ Double integration calculate/

Comparison:

Recaping the previous $D = \frac{1}{2} \oint_S xdy - ydx$, the original calculation method for this II-line integral is that find thier parametric equation $\begin{cases} x = x(t) \\ y = y(t) \end{cases}, t \in [a, b]$ for different (closed) curve S , then substitute into the calculation. From here, we could see that the essence of II-line integral is definite integral and has chirality.

However, "coordinate transform + Green's formula" method: $D = \iint_{\Delta} \frac{D(x, y)}{D(\xi, \eta)} d\xi d\eta$ as long as we master the coordinate transformation of a type of shape equation, we can easily calculate the results. Although there is a parameterization in the derivation, it is not reflected in the result (it can be seen from the process that if the parameterization is continued, it will be the result of the II-line integral in ξ, η -plane). In essence, the area is calculated as the volume with the same value, similiar with $\frac{1}{2} \oint_S xdy - ydx = \iint_D 1dxdy$ or $\oint_S xdy = \iint_D 1dxdy$ (If you unfold the left and right sides under the curved trapezoid, you can find the consistency.), it's just that the physical meanings of the two equations are different.



/Note: The choice of which method should be based on which is more convenient, parametric equations or coordinate transformations./

Variable Substitution in Double Integral


$$\iint_D f(x, y) dx dy = \begin{cases} x = x(\xi, \eta) \\ y = y(\xi, \eta) \end{cases} = \iint_{\Delta} f(x(\xi, \eta), y(\xi, \eta)) \left| \frac{D(x, y)}{D(\xi, \eta)} \right| d\xi d\eta$$

1.3 Surface Integral

1.3.1 Definition and basics

Definition

$S = \lim_{\lambda \rightarrow 0} \sum_i T_i$, where T_i is the projection area of S_i on the tangent plane at point $M_i \in S_i$.



Calculate Surface Integral (Omit matrix-vector operations)


$$S = \lim_{\lambda \rightarrow 0} \sum_i T_i = \iint_{\Delta} \sqrt{A^2 + B^2 + C^2} du dv = \iint_{\Delta} \sqrt{EG - F^2} du dv$$

where $A = \begin{vmatrix} y'_u & z'_u \\ y'_v & z'_v \end{vmatrix}$, $B = \begin{vmatrix} z'_u & x'_u \\ z'_v & x'_v \end{vmatrix}$, $C = \begin{vmatrix} x'_u & y'_u \\ x'_v & y'_v \end{vmatrix}$, $E = \sum_{x,y,z} x'^2_u$, $F = \sum_{x,y,z} x'_u x'_v$, $G = \sum_{x,y,z} x'^2_v$.

Procedure:

$\forall M_1 \in S'$, the coordinates of its projection M'_1 on T' are $\begin{cases} \xi = \xi(u, v) \\ \eta = \eta(u, v) \\ \zeta = \zeta(u, v) \end{cases}$,

then $S'_T = \iint_{\Delta} \left| \frac{D(\xi, \eta)}{D(u, v)} \right| du dv$, then the results could appear after a series of matrix-vector operations.



2 Multiple Integrals

2.1 The Riemann Integral over an n -Dimensional Interval

2.1.1 Comprehensions

First, the definition of Riemann integral over n -dimension interval is simimiar (almost same) with 1-dimension condition ($\int_I f(x)dx := \lim_{\lambda(P) \rightarrow 0} \sigma(f, P, \xi)$),

$$\int_I f(x^1, \dots, x^n) dx^1 \dots dx^n \text{ or } \int \dots \int_I f(x^1, \dots, x^n) dx^1 \dots dx^n$$

Then, introduce the necessity condition of it ($f \in \mathcal{R}(I) \implies f$ is bounded on I). After that, book gives some properties (Lemmas) about the set of measure zero (same with 1-dimension condition), and developes the Cantor Theorem (about uniform continuity of continuous functions on compact set). At last, we're given 2 criterions (necessity and sufficiency) of the Riemann integral (Darboux Criterion, Lebesgue Criterion).

Example. Let $f : I \rightarrow \mathbb{R}$ is a continuous real-value function on $I \subset \mathbb{R}^{n-1}$, then it's graph is a set measure zero in n -dimension.

Lemma 2.1. /Development of Cantor Theorem/ If the relation $\omega(f; x) < w_0$ holds at each point of a compact set K for the function $f : K \rightarrow \mathbb{R}$, then for every $\varepsilon > 0$ there exists $\delta > 0$ such that $\omega(f; U_\delta(x)) < w_0 + \varepsilon$ for each point $x \in K$.

Theorem 2.1. (Lebesgue's criterion). $f \in \mathcal{R}(I) \Leftrightarrow (f \text{ is bounded on } I) \wedge (f \text{ is continuous almost everywhere on } I)$.

Theorem 2.2. (Darboux Criterion) $f \in (I) \Leftrightarrow (f \text{ is bounded on } I) \wedge (\underline{\mathcal{J}} = \overline{\mathcal{J}})$, where $\underline{\mathcal{J}} = \sup_P s(f, P)$, $\overline{\mathcal{J}} = \inf_P S(f, P)$.

2.1.2 Problems Sets

Problem 1. a) Show that a set of measure zero has no interior points.

- b) Show that not having interior points by no means guarantees that a set is of measure zero.
- c) Construct a set having measure zero whose closure is the entire space $\mathbb{R}^n$.
- d) A set $E \subset I$ is said to have content zero if for every $\varepsilon > 0$ it can be covered by a finite system of intervals $I_1, \dots, I_k$ such that $\sum_{i=1}^k |I_i| < \varepsilon$. Is every bounded set of measure zero a set of content zero?
- e) Show that if a set $E \subset \mathbb{R}^n$ is the direct product $\mathbb{R} \times e$ of the line $\mathbb{R}$ and a set $e \subset \mathbb{R}^{n-1}$ of $(n-1)$ -dimensional measure zero, then E is a set of n -dimensional measure zero.

Proof. a) [Proof by contradiction] If E has a point x_0 as an interior point, then there exists a ball $B(x_0, r)$ such that $B(x_0, r) \subset E$. Since the ball has positive measure, it follows that E cannot have measure zero.

b) [Counterexample] Consider the irrational numbers $\mathbb{R} \setminus \mathbb{Q}$, which have no interior points in $\mathbb{R}$ but are not of measure zero.

c) /inspired by b)/ Consider the set $E = \mathbb{Q}^n$, the set of all points in $\mathbb{R}^n$ with rational coordinates. The closure of E is $\mathbb{R}^n$, and E has measure zero since it can be covered by countably many intervals whose total length can be made arbitrarily small.

d) [Principle] No, since the Lebesgue measure zero set can be covered by at most countable intervals (which means measure $\mu = o(1) \cdot \mathcal{O}(1)$), while a set of content zero can be covered by a finite number of intervals (which means the content (also a kind of measure) sometimes is $\mathcal{O}(1) \cdot \mathcal{O}(1)$). [Example] The $\mathbb{Q} \cap [0, 1]$ and Cantor set are sets of measure zero but none of content zero.

e) From the construction of the product measure λ_n (Lebesgue measure of $\mathbb{R}^n$) /Lemma1/ if $A \in \mathcal{B}(\mathbb{R}^k)$ and $B \in \mathcal{B}(\mathbb{R}^{n-k})$, ($0 < k < n$), then $\lambda_n(A \times B) = \lambda_k(A)\lambda_{n-k}(B)$. So here (using also that /Lemma2/ if $E_m \uparrow E$, $\lim_{m \rightarrow +\infty} \lambda_n(E_m) = \lambda_n(E)$) $\lambda_n(E) = \lambda_n(\mathbb{R} \times e) = \lim_{m \rightarrow +\infty} \lambda_n((-m, m) \times e) = \lim_{m \rightarrow +\infty} \lambda_1((-m, m)) \times \lambda_{n-1}(e) = \lim_{m \rightarrow +\infty} \lambda_1((-m, m)) \times 0 = 0$.

Notes. 1. The proof of e) /Lemma2/ could found in *Real and Complex Analysis* by Walter Rudin, Theorem 1.19(Page 16), based on the (positive) measure definition (countable additive of disjoint countable measurable sets).

- Problem 2.** a) Construct the analogue of the Dirichlet function in $\mathbb{R}^n$ and show that a bounded function $f : I \rightarrow \mathbb{R}$ equal to zero at almost every point of the interval I may still fail to belong to $\mathcal{R}(I)$.
- b) Show that if $f \in \mathcal{R}(I)$ and $f(x) = 0$, at almost all points of the interval I , then $\int_I f(x) dx = 0$.

Proof. a) [Counterexample] Consider $\mathcal{D}^n = \begin{cases} 1 & \text{if } x \in \mathbb{Q}^n \\ 0 & \text{if } x \in \mathbb{R}^n \setminus \mathbb{Q}^n \end{cases}$, the discontinuous set is $\mathcal{R}^n$.

b) /by definition/ $\int_I f(x) dx = \lim_{\lambda(P) \rightarrow 0} \sum_{\xi_i \in P} f(\xi_i) \Delta(x_i) = \lim_{\lambda(P) \rightarrow 0} \sum_{\xi_i \in A \cap P} f(\xi_i) \Delta(x_i) + \lim_{\lambda(P) \rightarrow 0} \sum_{\xi_j \in B \cap P} f(\xi_j) \Delta(x_i) = \mathcal{O}(1) \cdot o(1) + o(1) \cdot \mathcal{O}(1)$, where $A = \{\cup [x_k, x_{k+1}] \subset P : \forall k, \exists \xi \in [x_k, x_{k+1}], \text{ s.t. } f(\xi) \neq 0\}$ and $B = I \setminus A$.

Problem 3. There is a small difference between our earlier definition of the Riemann integral on a closed interval $I \subset \mathbb{R}$ and Definition 7 for the integral over an interval of arbitrary dimension. This difference involves the definition of a partition and the measure of an interval of the partition. Clarify this nuance for yourself and verify that

$$\int_a^b f(x) dx = \begin{cases} \int_I f(x) dx & \text{if } a < b \\ - \int_I f(x) dx & \text{if } a > b \end{cases}$$

where I is the interval on the real line $\mathbb{R}$ with endpoints a and b .

Proof. Just understand it, it's not hard.

- Problem 4.** a) Prove that a real-valued function $f : I \rightarrow \mathbb{R}$ defined on an interval $I \subset \mathbb{R}^n$ is integrable over that interval if and only if for every $\varepsilon > 0$ there exists a partition P of I such that $S(f; P) - s(f; P) < \varepsilon$.
- b) Using the result of a) and assuming that we are dealing with a real-valued function $f : I \rightarrow \mathbb{R}$, one can simplify slightly the proof of the sufficiency of the Lebesgue criterion. Try to carry out this simplification by yourself.

Proof. a) /by The Darboux Criterion/ $f \in (I) \Leftrightarrow (f \text{ is bounded on } I) \wedge (\underline{\mathcal{J}} = \overline{\mathcal{J}})$, where $\underline{\mathcal{J}} = \sup_P s(f, P)$, $\overline{\mathcal{J}} = \inf_P S(f, P)$. Notice that $\forall \varepsilon > 0, \exists P, \text{ s.t. } S(f, P) - s(f, P) < \varepsilon \Leftrightarrow \underline{\mathcal{J}} = \overline{\mathcal{J}}$ (proof is similar with 1-dimension condition), which is what we need to prove.

b) [Segmented estimation] Sufficiency: procedure is same with book, a) just simplify the last step.