

Unit 3: Foundations for inference

3. Decision errors, significance levels, sample size & power

Sta 101 - Spring 2015

Duke University, Department of Statistical Science

February 16, 2015

1. Housekeeping

2. Main ideas

1. Hypothesis tests and confidence intervals at equivalent significance/confidence levels should agree
2. Results that are statistically significant are not necessarily practically significant
3. Calculate the sample size a priori to achieve desired margin of error
4. Hypothesis tests are prone to decision errors
5. Power depends on the effect size, α , n , and s

3. Summary

- ▶ PA3 due tonight
- ▶ Office hours today: ask questions after class
- ▶ Midterm review: online, you should have received an email from WebEx with instructions
- ▶ Office hours tomorrow: online/on campus depending on weather

1. Housekeeping

2. Main ideas

1. Hypothesis tests and confidence intervals at equivalent significance/confidence levels should agree
2. Results that are statistically significant are not necessarily practically significant
3. Calculate the sample size a priori to achieve desired margin of error
4. Hypothesis tests are prone to decision errors
5. Power depends on the effect size, α , n , and s

3. Summary

1. Housekeeping


2. Main ideas

1. Hypothesis tests and confidence intervals at equivalent significance/confidence levels should agree
2. Results that are statistically significant are not necessarily practically significant
3. Calculate the sample size a priori to achieve desired margin of error
4. Hypothesis tests are prone to decision errors
5. Power depends on the effect size, α , n , and s

3. Summary

1. Hypothesis tests and confidence intervals at equivalent significance/confidence levels should agree


Two sided



95% confidence level
is equivalent to
two sided HT with $\alpha = 0.05$


1. Hypothesis tests and confidence intervals at equivalent significance/confidence levels should agree

Two sided



95% confidence level
is equivalent to
two sided HT with $\alpha = 0.05$

One sided



95% confidence level
is equivalent to
one sided HT with $\alpha = 0.025$

Clicker question

What is the significance level of a two-sided hypothesis test that is equivalent to a 90% confidence interval? *Hint: Draw a picture and mark the confidence level in the center.*

- (a) 0.001
- (b) 0.01
- (c) 0.025
- (d) 0.05
- (e) 0.10

Clicker question

What is the significance level of a two-sided hypothesis test that is equivalent to a 90% confidence interval? *Hint: Draw a picture and mark the confidence level in the center.*

- (a) 0.001
- (b) 0.01
- (c) 0.025
- (d) 0.05
- (e) **0.10**

Clicker question

What is the significance level of a one-sided hypothesis test that is equivalent to a 90% confidence interval? *Hint: Draw a picture and mark the confidence level in the center.*

- (a) 0.001
- (b) 0.01
- (c) 0.025
- (d) 0.05
- (e) 0.10

Clicker question

What is the significance level of a one-sided hypothesis test that is equivalent to a 90% confidence interval? *Hint: Draw a picture and mark the confidence level in the center.*

- (a) 0.001
- (b) 0.01
- (c) 0.025
- (d) **0.05**
- (e) 0.10

Clicker question

What is the confidence level of a confidence interval that is equivalent to a two-sided hypothesis test with $\alpha = 0.01$. *Hint: Draw a picture and mark the confidence level in the center.*

- (a) 0.80
- (b) 0.90
- (c) 0.95
- (d) 0.98
- (e) 0.99

Clicker question

What is the confidence level of a confidence interval that is equivalent to a two-sided hypothesis test with $\alpha = 0.01$. *Hint: Draw a picture and mark the confidence level in the center.*

- (a) 0.80
- (b) 0.90
- (c) 0.95
- (d) 0.98
- (e) **0.99**

Clicker question

What is the confidence level of a confidence interval that is equivalent to a one-sided hypothesis test with $\alpha = 0.01$. *Hint: Draw a picture and mark the confidence level in the center.*

- (a) 0.80
- (b) 0.90
- (c) 0.95
- (d) 0.98
- (e) 0.99

Clicker question

What is the confidence level of a confidence interval that is equivalent to a one-sided hypothesis test with $\alpha = 0.01$. *Hint: Draw a picture and mark the confidence level in the center.*

- (a) 0.80
- (b) 0.90
- (c) 0.95
- (d) **0.98**
- (e) 0.99

Clicker question

A 95% confidence interval for the average normal body temperature of humans is found to be (98.1 F, 98.4 F). Which of the following is true?

- (a) The hypothesis $H_A : \mu = 98.2$ would be rejected at $\alpha = 0.05$ in favor of $H_A : \mu \neq 98.2$.
- (b) The hypothesis $H_A : \mu = 98.2$ would be rejected at $\alpha = 0.025$ in favor of $H_A : \mu > 98.2$.
- (c) The hypothesis $H_A : \mu = 98$ would be rejected using a 90% confidence interval.
- (d) The hypothesis $H_A : \mu = 98.2$ would not be rejected using a 99% confidence interval.

Clicker question

A 95% confidence interval for the average normal body temperature of humans is found to be (98.1 F, 98.4 F). Which of the following is true?

- (a) The hypothesis $H_A : \mu = 98.2$ would be rejected at $\alpha = 0.05$ in favor of $H_A : \mu \neq 98.2$.
- (b) The hypothesis $H_A : \mu = 98.2$ would be rejected at $\alpha = 0.025$ in favor of $H_A : \mu > 98.2$.
- (c) *The hypothesis $H_A : \mu = 98$ would be rejected using a 90% confidence interval.*
- (d) The hypothesis $H_A : \mu = 98.2$ would not be rejected using a 99% confidence interval.

1. Housekeeping

2. Main ideas

1. Hypothesis tests and confidence intervals at equivalent significance/confidence levels should agree
- 2. Results that are statistically significant are not necessarily practically significant**
3. Calculate the sample size a priori to achieve desired margin of error
4. Hypothesis tests are prone to decision errors
5. Power depends on the effect size, α , n , and s

3. Summary

2. Results that are statistically significant are not necessarily practically significant

Clicker question

All else held equal, will p-value be lower if $n = 100$ or $n = 10,000$?

- (a) $n = 100$
- (b) $n = 10,000$

2. Results that are statistically significant are not necessarily practically significant

Clicker question

All else held equal, will p-value be lower if $n = 100$ or $n = 10,000$?

- (a) $n = 100$
- (b) $n = 10,000$

2. Results that are statistically significant are not necessarily practically significant

Clicker question

All else held equal, will p-value be lower if $n = 100$ or $n = 10,000$?

(a) $n = 100$

(b) $n = 10,000$

Suppose $\bar{x} = 5$, $s = 2$, $H_0 : \mu = 4.5$, and $H_A : \mu \geq 4.5$.

2. Results that are statistically significant are not necessarily practically significant

Clicker question

All else held equal, will p-value be lower if $n = 100$ or $n = 10,000$?

(a) $n = 100$

(b) $n = 10,000$

Suppose $\bar{x} = 5$, $s = 2$, $H_0 : \mu = 4.5$, and $H_A : \mu \geq 4.5$.

$$Z_{n=100} = \frac{5 - 4.5}{\frac{2}{\sqrt{100}}}$$

2. Results that are statistically significant are not necessarily practically significant

Clicker question

All else held equal, will p-value be lower if $n = 100$ or $n = 10,000$?

(a) $n = 100$

(b) $n = 10,000$

Suppose $\bar{x} = 5$, $s = 2$, $H_0 : \mu = 4.5$, and $H_A : \mu \geq 4.5$.

$$Z_{n=100} = \frac{5 - 4.5}{\frac{2}{\sqrt{100}}} = \frac{5 - 4.5}{\frac{2}{10}} = \frac{0.5}{0.2} = 2.5, \quad p\text{-value} = 0.0062$$

2. Results that are statistically significant are not necessarily practically significant

Clicker question

All else held equal, will p-value be lower if $n = 100$ or $n = 10,000$?

(a) $n = 100$

(b) $n = 10,000$

Suppose $\bar{x} = 5$, $s = 2$, $H_0 : \mu = 4.5$, and $H_A : \mu \geq 4.5$.

$$Z_{n=100} = \frac{5 - 4.5}{\frac{2}{\sqrt{100}}} = \frac{5 - 4.5}{\frac{2}{10}} = \frac{0.5}{0.2} = 2.5, \quad p\text{-value} = 0.0062$$

$$Z_{n=10000} = \frac{5 - 4.5}{\frac{2}{\sqrt{10000}}}$$

2. Results that are statistically significant are not necessarily practically significant

Clicker question

All else held equal, will p-value be lower if $n = 100$ or $n = 10,000$?

(a) $n = 100$

(b) $n = 10,000$

Suppose $\bar{x} = 5$, $s = 2$, $H_0 : \mu = 4.5$, and $H_A : \mu \geq 4.5$.

$$Z_{n=100} = \frac{5 - 4.5}{\frac{2}{\sqrt{100}}} = \frac{5 - 4.5}{\frac{2}{10}} = \frac{0.5}{0.2} = 2.5, \quad p\text{-value} = 0.0062$$

$$Z_{n=10000} = \frac{5 - 4.5}{\frac{2}{\sqrt{10000}}} = \frac{5 - 4.5}{\frac{2}{100}} = \frac{0.5}{0.02} = 25, \quad p\text{-value} \approx 0$$

2. Results that are statistically significant are not necessarily practically significant

Clicker question

All else held equal, will p-value be lower if $n = 100$ or $n = 10,000$?

(a) $n = 100$

(b) $n = 10,000$

Suppose $\bar{x} = 5$, $s = 2$, $H_0 : \mu = 4.5$, and $H_A : \mu \geq 4.5$.

$$Z_{n=100} = \frac{5 - 4.5}{\frac{2}{\sqrt{100}}} = \frac{5 - 4.5}{\frac{2}{10}} = \frac{0.5}{0.2} = 2.5, \quad p\text{-value} = 0.0062$$

$$Z_{n=10000} = \frac{5 - 4.5}{\frac{2}{\sqrt{10000}}} = \frac{5 - 4.5}{\frac{2}{100}} = \frac{0.5}{0.02} = 25, \quad p\text{-value} \approx 0$$

As n increases - $SE \downarrow$, $Z \uparrow$, $p\text{-value} \downarrow$

1. Housekeeping

2. Main ideas

1. Hypothesis tests and confidence intervals at equivalent significance/confidence levels should agree
2. Results that are statistically significant are not necessarily practically significant
3. Calculate the sample size a priori to achieve desired margin of error
4. Hypothesis tests are prone to decision errors
5. Power depends on the effect size, α , n , and s

3. Summary

3. Calculate the sample size *a priori* to achieve desired margin of error

Application exercise: 3.3 Sample size

See course website for details.

1. Housekeeping

2. Main ideas

1. Hypothesis tests and confidence intervals at equivalent significance/confidence levels should agree
2. Results that are statistically significant are not necessarily practically significant
3. Calculate the sample size a priori to achieve desired margin of error
- 4. Hypothesis tests are prone to decision errors**
5. Power depends on the effect size, α , n , and s

3. Summary

4. Hypothesis tests are prone to decision errors

		Decision	
		fail to reject H_0	reject H_0
Truth	H_0 true		
	H_A true		

4. Hypothesis tests are prone to decision errors

		Decision	
		fail to reject H_0	reject H_0
Truth	H_0 true		
	H_A true		

4. Hypothesis tests are prone to decision errors

		Decision	
		fail to reject H_0	reject H_0
Truth	H_0 true		
	H_A true		

4. Hypothesis tests are prone to decision errors

		Decision	
		fail to reject H_0	reject H_0
Truth	H_0 true	✓	Type 1 Error, α
	H_A true		

- A *Type 1 Error* is rejecting the null hypothesis when H_0 is true: α
- For those cases where H_0 is actually true, we do not want to incorrectly reject it more than 5% of those times
 - Increasing α increases the Type 1 error rate, hence we prefer to small values of α

4. Hypothesis tests are prone to decision errors

	Decision	
	fail to reject H_0	reject H_0
Truth	H_0 true	✓ <i>Type 1 Error, α</i>
	H_A true	<i>Type 2 Error, β</i>

- ▶ A *Type 1 Error* is rejecting the null hypothesis when H_0 is true: α
 - For those cases where H_0 is actually true, we do not want to incorrectly reject it more than 5% of those times
 - Increasing α increases the Type 1 error rate, hence we prefer to small values of α
- ▶ A *Type 2 Error* is failing to reject the null hypothesis when H_A is true: β

4. Hypothesis tests are prone to decision errors

Truth	Decision	
	fail to reject H_0	reject H_0
	H_0 true	H_A true
	✓	Type 1 Error, α
	Type 2 Error, β	Power, $1 - \beta$

- ▶ A **Type 1 Error** is rejecting the null hypothesis when H_0 is true: α
 - For those cases where H_0 is actually true, we do not want to incorrectly reject it more than 5% of those times
 - Increasing α increases the Type 1 error rate, hence we prefer to small values of α
- ▶ A **Type 2 Error** is failing to reject the null hypothesis when H_A is true: β
- ▶ **Power** is the probability of correctly rejecting H_0 , and hence the complement of the probability of a Type 2 Error: $1 - \beta$

1. Housekeeping

2. Main ideas

1. Hypothesis tests and confidence intervals at equivalent significance/confidence levels should agree
2. Results that are statistically significant are not necessarily practically significant
3. Calculate the sample size a priori to achieve desired margin of error
4. Hypothesis tests are prone to decision errors
5. Power depends on the effect size, α , n , and s

3. Summary

Power can be increased (and hence Type 2 error rate can be decreased) by

Power can be increased (and hence Type 2 error rate can be decreased) by

- ▶ increasing the sample size

Power can be increased (and hence Type 2 error rate can be decreased) by

- ▶ increasing the sample size
- ▶ decreasing the standard deviation of the sample (difficult to ensure but cautious measurement process and limiting the population so that it is more homogenous may help)

Power can be increased (and hence Type 2 error rate can be decreased) by

- ▶ increasing the sample size
- ▶ decreasing the standard deviation of the sample (difficult to ensure but cautious measurement process and limiting the population so that it is more homogenous may help)
- ▶ increasing α

Power can be increased (and hence Type 2 error rate can be decreased) by

- ▶ increasing the sample size
- ▶ decreasing the standard deviation of the sample (difficult to ensure but cautious measurement process and limiting the population so that it is more homogenous may help)
- ▶ increasing α
- ▶ increasing the *effect size*

1. Housekeeping

2. Main ideas

1. Hypothesis tests and confidence intervals at equivalent significance/confidence levels should agree
2. Results that are statistically significant are not necessarily practically significant
3. Calculate the sample size a priori to achieve desired margin of error
4. Hypothesis tests are prone to decision errors
5. Power depends on the effect size, α , n , and s

3. Summary

1. Hypothesis tests and confidence intervals at equivalent significance/confidence levels should agree
2. Results that are statistically significant are not necessarily practically significant
3. Calculate the sample size a priori to achieve desired margin of error
4. Hypothesis tests are prone to decision errors
5. Power depends on the effect size, α , n , and s