

Unit 3: Foundations for inference

2. Confidence intervals and hypothesis tests

Sta 101 - Spring 2015

Duke University, Department of Statistical Science

February 11, 2015

1. Housekeeping

2. Main ideas

1. Statistical inference methods based on the CLT depend on the same conditions as the CLT
2. Use confidence intervals to estimate population parameters
3. Critical value depends on the confidence level
4. Use hypothesis tests to make decisions about population parameters

3. Summary

- ▶ No new PS this week, but review questions provided on new material that will be on the exam next week
- ▶ Sample midterm + review questions posted on course website
- ▶ Additional practice quizzes on Coursera

Duke UNIVERSITY Data Analysis and Statistical Inference
by Dr. Mine Çetinkaya-Rundel

Quizzes

▼ **Unit 1**

Unit 1 Quiz - Introduction to Data

[Attempt Quiz](#)

Hard Deadline Mon 15 Sep 2014 10:00 AM PDT
If you submit any time after the hard deadline, you will not receive credit.

Effective Score N/A
Each time that you attempt it, we'll record a score based on your performance and any penalties due to late submit highest score of all the allowed attempts made before the hard deadline.

of Attempts 0 / 3

▼ **Unit 2**

Unit 2 Quiz - Probability and Distributions

- ▶ PA3 will be posted on Monday after class and will be due that evening, so budget your time accordingly for that short turnaround!

1. Housekeeping

2. Main ideas

1. Statistical inference methods based on the CLT depend on the same conditions as the CLT
2. Use confidence intervals to estimate population parameters
3. Critical value depends on the confidence level
4. Use hypothesis tests to make decisions about population parameters

3. Summary

1. Housekeeping

2. Main ideas

1. Statistical inference methods based on the CLT depend on the same conditions as the CLT
2. Use confidence intervals to estimate population parameters
3. Critical value depends on the confidence level
4. Use hypothesis tests to make decisions about population parameters

3. Summary

1. Statistical inference methods based on the CLT depend on the same conditions as the CLT

Always check these in context of the data and the research question!

1. *Independence*: Sampled observations must be independent.

This is difficult to verify, but is more likely if

- random sampling/assignment is used, and,
- if sampling without replacement, $n < 10\%$ of the population.

2. *Sample size/skew*: Either the population distribution is normal

or

$n > 30$ and the population distribution is not extremely skewed (the more skewed the distribution, the higher n necessary for the CLT to apply).

This is also difficult to verify for the population, but we can 2

1. Housekeeping

2. Main ideas

1. Statistical inference methods based on the CLT depend on the same conditions as the CLT
- 2. Use confidence intervals to estimate population parameters**
3. Critical value depends on the confidence level
4. Use hypothesis tests to make decisions about population parameters

3. Summary

2. Use confidence intervals to estimate population parameters

CI: point estimate \pm margin of error

2. Use confidence intervals to estimate population parameters

$$CI: \textit{point estimate} \pm \textit{margin of error}$$

If the parameter of interest is the population mean, and the point estimate is the sample mean,

$$\bar{x} \pm Z^* \frac{s}{\sqrt{n}}$$

Application exercise: 3.1 Confidence interval for a single mean

See course website for details.

1. Housekeeping

2. Main ideas

1. Statistical inference methods based on the CLT depend on the same conditions as the CLT
2. Use confidence intervals to estimate population parameters
- 3. Critical value depends on the confidence level**
4. Use hypothesis tests to make decisions about population parameters

3. Summary

3. Critical value depends on the confidence level

Clicker question

What is the critical value (Z^*) for a confidence interval at the 91% confidence level?


- (a) $Z^* = 1.34$
- (b) $Z^* = 1.65$
- (c) $Z^* = 1.70$
- (d) $Z^* = 1.96$
- (e) $Z^* = 2.33$

3. Critical value depends on the confidence level

Clicker question

What is the critical value (Z^*) for a confidence interval at the 91% confidence level?

- (a) $Z^* = 1.34$
- (b) $Z^* = 1.65$
- (c) $Z^* = 1.70$
- (d) $Z^* = 1.96$
- (e) $Z^* = 2.33$



1. The confidence level of a confidence interval is the probability that a given interval contains the true population parameter.

This is incorrect, CIs are part of the frequentist paradigm and as such the population parameter is fixed but unknown. Consequently, the probability any given CI contains the true value must be 0 or 1 (it does or does not).

1. The confidence level of a confidence interval is the probability that a given interval contains the true population parameter.

This is incorrect, CIs are part of the frequentist paradigm and as such the population parameter is fixed but unknown. Consequently, the probability any given CI contains the true value must be 0 or 1 (it does or does not).

2. A narrower confidence interval is always better.

This is incorrect since the width is a function of both the confidence level and the standard error.

1. The confidence level of a confidence interval is the probability that a given interval contains the true population parameter.

This is incorrect, CIs are part of the frequentist paradigm and as such the population parameter is fixed but unknown. Consequently, the probability any given CI contains the true value must be 0 or 1 (it does or does not).

2. A narrower confidence interval is always better.

This is incorrect since the width is a function of both the confidence level and the standard error.

3. A wider interval means less confidence.

This is incorrect since it is possible to make very precise statements with very little confidence.

1. Housekeeping

2. Main ideas

1. Statistical inference methods based on the CLT depend on the same conditions as the CLT
2. Use confidence intervals to estimate population parameters
3. Critical value depends on the confidence level
4. Use hypothesis tests to make decisions about population parameters

3. Summary

4. Use hypothesis tests to make decisions about population parameters

Hypothesis testing framework:

1. Set the hypotheses.
2. Check assumptions and conditions.
3. Calculate a *test statistic* and a p-value.
4. Make a decision, and interpret it in context of the research question.

1. Set the hypotheses

- $H_0 : \mu = \text{null value}$
- $H_A : \mu < \text{or } > \text{or } \neq \text{null value}$

1. Set the hypotheses

- $H_0 : \mu = \text{null value}$
- $H_A : \mu < \text{or } > \text{or } \neq \text{null value}$

2. Check assumptions and conditions

- Independence: random sample/assignment, 10% condition when sampling without replacement
- Sample size / skew: $n \geq 30$ (or larger if sample is skewed), no extreme skew

1. Set the hypotheses
 - $H_0 : \mu = \text{null value}$
 - $H_A : \mu < \text{or } > \text{or } \neq \text{null value}$
2. Check assumptions and conditions
 - Independence: random sample/assignment, 10% condition when sampling without replacement
 - Sample size / skew: $n \geq 30$ (or larger if sample is skewed), no extreme skew
3. Calculate a *test statistic* and a p-value (draw a picture!)

$$Z = \frac{\bar{x} - \mu}{SE}, \text{ where } SE = \frac{s}{\sqrt{n}}$$

1. Set the hypotheses
 - $H_0 : \mu = \text{null value}$
 - $H_A : \mu < \text{or } > \text{or } \neq \text{null value}$
2. Check assumptions and conditions
 - Independence: random sample/assignment, 10% condition when sampling without replacement
 - Sample size / skew: $n \geq 30$ (or larger if sample is skewed), no extreme skew

3. Calculate a *test statistic* and a p-value (draw a picture!)

$$Z = \frac{\bar{x} - \mu}{SE}, \text{ where } SE = \frac{s}{\sqrt{n}}$$

4. Make a decision, and interpret it in context of the research question
 - If p-value $< \alpha$, reject H_0 , data provide evidence for H_A
 - If p-value $> \alpha$, do not reject H_0 , data do not provide evidence for H_A

4. Use hypothesis tests to make decisions about population parameters

Application exercise: 3.2 Hypothesis testing for a single mean

See course website for details.

Clicker question

Which of the following is the correct interpretation of the p-value?

- (a) The probability that average GPA of Duke students has changed since 2001.
- (b) The probability that average GPA of Duke students has not changed since 2001.
- (c) The probability that average GPA of Duke students has not changed since 2001, if in fact a random sample of 63 Duke students this year have an average GPA of 3.58 or higher.
- (d) The probability that a random sample of 63 Duke students have an average GPA of 3.58 or higher, if in fact the average GPA has not changed since 2001.
- (e) The probability that a random sample of 63 Duke students have an average GPA of 3.58 or higher or 3.16 or lower, if in fact the average GPA has not changed since 2001.

Clicker question

Which of the following is the correct interpretation of the p-value?

- (a) The probability that average GPA of Duke students has changed since 2001.
- (b) The probability that average GPA of Duke students has not changed since 2001.
- (c) The probability that average GPA of Duke students has not changed since 2001, if in fact a random sample of 63 Duke students this year have an average GPA of 3.58 or higher.
- (d) The probability that a random sample of 63 Duke students have an average GPA of 3.58 or higher, if in fact the average GPA has not changed since 2001.
- (e) *The probability that a random sample of 63 Duke students have an average GPA of 3.58 or higher or 3.16 or lower, if in fact the average GPA has not changed since 2001.*

1. P-value is the probability that the null hypothesis is true
This is incorrect, p-value is the probability of the observed or more extreme outcome if in fact the null hypothesis is correct. It is the conditional probability of the observed data (or something more extreme), conditioned on the null hypothesis being correct.

1. P-value is the probability that the null hypothesis is true
This is incorrect, p-value is the probability of the observed or more extreme outcome if in fact the null hypothesis is correct. It is the conditional probability of the observed data (or something more extreme), conditioned on the null hypothesis being correct.
2. A high p-value confirms the null hypothesis.
This is incorrect, a high p-value means the data do not provide convincing evidence for the alternative hypothesis and hence that the null hypothesis can't be rejected.

1. P-value is the probability that the null hypothesis is true
This is incorrect, p -value is the probability of the observed or more extreme outcome if in fact the null hypothesis is correct. It is the conditional probability of the observed data (or something more extreme), conditioned on the null hypothesis being correct.
2. A high p -value confirms the null hypothesis.
This is incorrect, a high p -value means the data do not provide convincing evidence for the alternative hypothesis and hence that the null hypothesis can't be rejected.
3. A low p -value confirms the alternative hypothesis.
This is incorrect, a low p -value means the data provide convincing evidence for the alternative hypothesis, but not necessarily that it is confirmed.

1. Housekeeping

2. Main ideas

1. Statistical inference methods based on the CLT depend on the same conditions as the CLT
2. Use confidence intervals to estimate population parameters
3. Critical value depends on the confidence level
4. Use hypothesis tests to make decisions about population parameters

3. Summary

1. Statistical inference methods based on the CLT depend on the same conditions as the CLT
2. Use confidence intervals to estimate population parameters
3. Critical value depends on the confidence level
4. Use hypothesis tests to make decisions about population parameters