

# **Unit 2: Probability and distributions**

## 3. Probability and conditional probability

Sta 101 - Spring 2015

Duke University, Department of Statistical Science

January 26, 2015

## 1. Housekeeping

## 2. Main ideas

1. Disjoint and independent do mean the same thing
2. Application of the addition rule depends on disjointness of events
3. Bayes' theorem works for all types of events

## 3. Summary

- ▶ Please review the regrade policy on the course syllabus:  
*Regrade requests must be made within 3 days of when the assignment is returned, and must be submitted in writing. These will be honored if points were tallied incorrectly, or if you feel your answer is correct but it was marked wrong. No regrade will be made to alter the number of points deducted for a mistake. There will be no grade changes after the final exam.*
- ▶ Any regrade requests should be submitted to me, I will regrade the entire assignment.

## 1. Housekeeping

## 2. Main ideas

1. Disjoint and independent do mean the same thing
2. Application of the addition rule depends on disjointness of events
3. Bayes' theorem works for all types of events

## 3. Summary

## 1. Housekeeping

## 2. Main ideas

1. Disjoint and independent do mean the same thing
2. Application of the addition rule depends on disjointness of events
3. Bayes' theorem works for all types of events

## 3. Summary

## 1. Disjoint and independent do not mean the same thing

- ▶ *Disjoint (mutually exclusive) events* cannot happen at the same time
  - A voter cannot register as a Democrat and a Republican at the same time
  - But s/he might be a Republican and a Moderate at the same time – *non-disjoint events*
  - For disjoint A and B:  $P(A \text{ and } B) = 0$

## 1. Disjoint and independent do not mean the same thing

- ▶ *Disjoint (mutually exclusive) events* cannot happen at the same time
  - A voter cannot register as a Democrat and a Republican at the same time
  - But s/he might be a Republican and a Moderate at the same time – *non-disjoint events*
  - For disjoint A and B:  $P(A \text{ and } B) = 0$
- ▶ If A and B are *independent events*, having information on A does not tell us anything about B (and vice versa)
  - If A and B are independent:
    - $P(A | B) = P(A)$
    - $P(A \text{ and } B) = P(A) \times P(B)$

## 1. Housekeeping

## 2. Main ideas

1. Disjoint and independent do mean the same thing
2. Application of the addition rule depends on disjointness of events
3. Bayes' theorem works for all types of events

## 3. Summary



## 2. Application of the addition rule depends on disjointness of events


- ▶ *General addition rule:*  $P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$
- ▶  $A \text{ or } B = \text{either } A \text{ or } B \text{ or both}$

## 2. Application of the addition rule depends on disjointness of events

- ▶ *General addition rule:*  $P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$
- ▶  $A \text{ or } B$  = either A or B or both

### **disjoint events:**

$$\begin{aligned} &P(A \text{ or } B) \\ &= P(A) + P(B) - P(A \text{ and } B) \\ &= 0.4 + 0.3 - 0 = 0.7 \end{aligned}$$




## 2. Application of the addition rule depends on disjointness of events

- ▶ *General addition rule:*  $P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$
- ▶  $A \text{ or } B$  = either A or B or both


### disjoint events:

$$\begin{aligned}P(A \text{ or } B) \\&= P(A) + P(B) - P(A \text{ and } B) \\&= 0.4 + 0.3 - 0 = 0.7\end{aligned}$$



### non-disjoint events:

$$\begin{aligned}P(A \text{ or } B) \\&= P(A) + P(B) - P(A \text{ and } B) \\&= 0.4 + 0.3 - 0.02 = 0.68\end{aligned}$$



## 1. Housekeeping

## 2. Main ideas

1. Disjoint and independent do mean the same thing
2. Application of the addition rule depends on disjointness of events
3. Bayes' theorem works for all types of events

## 3. Summary

### 3. Bayes' theorem works for all types of events

► *Bayes' theorem:*  $P(A | B) = \frac{P(A \text{ and } B)}{P(B)}$

### 3. Bayes' theorem works for all types of events

- ▶ *Bayes' theorem:*  $P(A | B) = \frac{P(A \text{ and } B)}{P(B)}$
- ▶ ... can be rewritten as:  $P(A \text{ and } B) = P(A | B) \times P(B)$

### 3. Bayes' theorem works for all types of events

- ▶ *Bayes' theorem*:  $P(A | B) = \frac{P(A \text{ and } B)}{P(B)}$
- ▶ ... can be rewritten as:  $P(A \text{ and } B) = P(A | B) \times P(B)$

#### **disjoint events:**

- ▶ We know  $P(A | B) = 0$ ,  
since if B happened A  
could not have happened

### 3. Bayes' theorem works for all types of events

- ▶ *Bayes' theorem:*  $P(A | B) = \frac{P(A \text{ and } B)}{P(B)}$
- ▶ ... can be rewritten as:  $P(A \text{ and } B) = P(A | B) \times P(B)$

#### **disjoint events:**

- ▶ We know  $P(A | B) = 0$ ,  
since if B happened A  
could not have happened
- ▶  $P(A \text{ and } B)$   
 $= P(A | B) \times P(B)$



### 3. Bayes' theorem works for all types of events

- ▶ *Bayes' theorem:*  $P(A | B) = \frac{P(A \text{ and } B)}{P(B)}$
- ▶ ... can be rewritten as:  $P(A \text{ and } B) = P(A | B) \times P(B)$

#### **disjoint events:**

- ▶ We know  $P(A | B) = 0$ ,  
since if B happened A  
could not have happened
- ▶  $P(A \text{ and } B)$   
 $= P(A | B) \times P(B)$   
 $= 0 \times P(B) = 0$

### 3. Bayes' theorem works for all types of events

- ▶ *Bayes' theorem:*  $P(A | B) = \frac{P(A \text{ and } B)}{P(B)}$
- ▶ ... can be rewritten as:  $P(A \text{ and } B) = P(A | B) \times P(B)$

#### disjoint events:

- ▶ We know  $P(A | B) = 0$ ,  
since if B happened A  
could not have happened
- ▶  $P(A \text{ and } B)$   
 $= P(A | B) \times P(B)$   
 $= 0 \times P(B) = 0$

#### independent events:

- ▶ We know  $P(A | B) = P(A)$ ,  
since knowing B doesn't  
tell us anything about A

### 3. Bayes' theorem works for all types of events

- ▶ *Bayes' theorem:*  $P(A | B) = \frac{P(A \text{ and } B)}{P(B)}$
- ▶ ... can be rewritten as:  $P(A \text{ and } B) = P(A | B) \times P(B)$

#### disjoint events:

- ▶ We know  $P(A | B) = 0$ ,  
since if B happened A  
could not have happened
- ▶  $P(A \text{ and } B)$   
 $= P(A | B) \times P(B)$   
 $= 0 \times P(B) = 0$

#### independent events:

- ▶ We know  $P(A | B) = P(A)$ ,  
since knowing B doesn't  
tell us anything about A
- ▶  $P(A \text{ and } B)$   
 $= P(A | B) \times P(B)$

### 3. Bayes' theorem works for all types of events

- ▶ *Bayes' theorem:*  $P(A | B) = \frac{P(A \text{ and } B)}{P(B)}$
- ▶ ... can be rewritten as:  $P(A \text{ and } B) = P(A | B) \times P(B)$

#### disjoint events:

- ▶ We know  $P(A | B) = 0$ ,  
since if B happened A  
could not have happened
- ▶  $P(A \text{ and } B)$   
 $= P(A | B) \times P(B)$   
 $= 0 \times P(B) = 0$

#### independent events:

- ▶ We know  $P(A | B) = P(A)$ ,  
since knowing B doesn't  
tell us anything about A
- ▶  $P(A \text{ and } B)$   
 $= P(A | B) \times P(B)$   
 $= P(A) \times P(B)$

## Application exercise: 2.1 Probability and conditional probability

See the course website for instructions.

## 1. Housekeeping

## 2. Main ideas

1. Disjoint and independent do mean the same thing
2. Application of the addition rule depends on disjointness of events
3. Bayes' theorem works for all types of events

## 3. Summary

1. Disjoint and independent do mean the same thing
2. Application of the addition rule depends on disjointness of events
3. Bayes' theorem works for all types of events