

Unimetry: A Phase-Space Reformulation of Special Relativity

Timur Abizgeldin
Independent researcher, Austria
timurabizgeldin@gmail.com

September 7, 2025

Abstract

We propose a compact reformulation of special relativity in which spacetime units (time and length) are treated as *phase velocities*—directional derivatives of a single underlying parameter, the phase $\vec{\chi} \in \mathbb{C}$. The observable Minkowski interval emerges as a conserved quantity under a change of parameter from the hidden phase coordinate χ to the observer’s proper time τ . In this *unimetry* formalism, familiar relativistic effects—time dilation, Lorentz factor, Doppler shift, and relativistic velocity composition—arise as elementary projections and rotations in a Euclidean phase plane. Hyperbolic features of Lorentz kinematics reappear after a reparameterization of time, yielding the standard relations without altering empirical content. We provide closed-form derivations of the longitudinal/transverse Doppler factors, identify a simple lemma equating the total phase speed to the conserved Minkowski norm, and outline connections to gauge phases, rapidity, and a cosmological time gauge.

Keywords: special relativity; phase; rapidity; Doppler shift; Lorentz factor; phase parameterization.

MSC/PhCS: 83A05; 83-10; 70A05.

1 Introduction

We usually take time and space as primitive. The *phase formalism* introduced here suggests a different viewpoint: time and space are *derived projections* of a single parameter $\vec{\chi} \in \mathbb{C}$ (“phase”). In this picture, relativistic effects such as time dilation and the Doppler shift are geometric consequences of phase-flow rotations.

The proposal does not modify physics; it reorganizes familiar relations in a simpler language. In spirit it is akin to Lagrangian/Hamiltonian re-descriptions of classical mechanics: same empirical content, different coordinates. Throughout, Greek θ will denote the *external* rotation angle associated with relative motion, while ζ denotes an *internal* angle associated with the object’s intrinsic state (mass/density heuristic). We emphasize that no modification of Einstein’s dynamics is proposed; all results are kinematical identities obtained by a change of parameter.

Notation. Tildes, dots and primes indicate derivatives with respect to the phase parameter, proper time, and spatial arclength:

$$\tilde{X} := \frac{dX}{d\chi}, \quad \dot{X} := \frac{dX}{d\tau}, \quad X' := \frac{dX}{dl}.$$

We use c for the speed of light; $\beta := V/c$, $\gamma := 1/\sqrt{1-\beta^2}$, rapidity $\tanh \eta = \beta$. The subscript l in dx_l denotes spatial components, with $l = 1, 2, 3$ a Cartesian index.

2 Time and space as phase derivatives

Let $\vec{\chi} \in \mathbb{C}$ be a variable whose change generates observable time-space effects. We treat the time and space units as directional derivatives (phase velocities) along the real and imaginary directions of a complex basis (\hat{h}, \mathbf{l}) :

$$\hat{h} dx_0 = \frac{\partial \vec{\chi}}{\partial \chi_h} \frac{d\chi_h}{d\chi} d\chi = \tilde{H} d\chi, \quad \mathbf{l} dx_l = \frac{\partial \vec{\chi}}{\partial \chi_l} \frac{d\chi_l}{d\chi} d\chi = \tilde{L} d\chi, \quad l = 1, 2, 3. \quad (2.1)$$

Introduce the phase speed of the SR interval $ds = \tilde{S} d\chi$. The interval conservation takes the form

$$\tilde{S}^2 = \frac{ds^2}{d\chi^2} = \frac{g_{ij} dx^i dx^j}{d\chi^2} = \tilde{H}^2 - \tilde{L}^2, \quad (2.2)$$

equivalently

$$\tilde{H}^2 = \tilde{S}^2 + \tilde{L}^2. \quad (2.3)$$

Writing

$$\tilde{S} = \tilde{H} \cos \theta, \quad \tilde{L} = \tilde{H} \sin \theta, \quad (2.4)$$

where θ is the angle of the phase speed relative to the real axis. Algebraically, (??) is a Euclidean decomposition of a single speed into orthogonal projections; physically, we will see that under reparameterization the *projection* \tilde{S} , not the Euclidean norm \tilde{H} , is the conserved Minkowski quantity.

3 Phase space (*khōra*)

Let the phase vector space (“*khōra*”, after Plato) be \mathbb{C} with orthonormal basis (\hat{h}, \mathbf{l}) . For a phase vector $\vec{\chi} = R e^{\theta \mathbf{l}}$ with $\theta \in [-\pi, \pi]$,

$$\tilde{H} = R, \quad \tilde{S} = R \cos \theta, \quad \tilde{L} = R \sin \theta. \quad (3.1)$$

Choosing coordinates where the projectors onto (\hat{h}, \mathbf{l}) are unit, (??) simplifies to

$$\hat{h} dx_0 = \frac{d\chi_h}{d\chi} d\chi = \tilde{H} d\chi, \quad \mathbf{l} dx_l = \frac{d\chi_l}{d\chi} d\chi = \tilde{L} d\chi. \quad (3.2)$$

The map from phase to observables is an integral transform:

$$x^i(\chi) = x^i(\chi_0) + \int_{\chi_0}^{\chi} \tilde{X}^i(u) du, \quad i = 0, 1, 2, 3, \quad (3.3)$$

where \tilde{X}^i are projections of $d\vec{\chi}/d\chi$ onto (\hat{h}, \mathbf{l}) and $x^i(\chi_0)$ fix initial conditions.

4 Objects

A *fundamental particle* is an *elementary object* with nonzero phase $\vec{\chi} \neq 0$. Composite *objects* are phase configurations; to represent them *in phase space* one may require additional dimensions, except for the *photon*, whose phase is always aligned with the imaginary axis:

$$\mathbf{p} = \frac{d\vec{\chi}}{d\chi_l} = p \mathbf{l} \in \Im. \quad (4.1)$$

Non-photonic phenomena are associated with nonzero real projection and nonzero mass. A complex object can be identified with an *event* or worldline; the photon corresponds to a null-interval point encoding information about the event.

Any object's phase can be rotated to the *zero* (purely real) direction,

$$\vec{\chi}_0 = R \in \Re. \quad (4.2)$$

An object A moving with speed V relative to a rest observer has

$$\vec{\chi}_A = R e^{\theta_A \mathbf{l}}, \quad \sin \theta_A = \frac{V}{c} \equiv \beta. \quad (4.3)$$

4.1 Space as a symmetric phase pair

From (??), a naive zero-angle limit would remove the imaginary projection, contradicting observability. We enforce a nonvanishing spatial projection by pairing opposite-phase tilts:

$$\vec{\chi}^\pm = R e^{\pm \zeta \mathbf{l}}, \quad \vec{\chi}_l := \frac{\vec{\chi}^+ - \vec{\chi}^-}{2} = R \mathbf{l} \sin \zeta, \quad (4.4)$$

where ζ is an *internal angle* (intrinsic to the object; heuristically linked to mass/density). The local decomposition is

$$\vec{\chi}_0 = \vec{\chi}_\tau + \vec{\chi}_l = R \cos \zeta + R \mathbf{l} \sin \zeta, \quad (4.5)$$

with unit components (normalized by R): the real component is $\cos \zeta$ and the imaginary component is $\sin \zeta$.

4.2 Absolute, local, and observed time

Define *absolute* time $t = t(\tilde{H})$ at the zero phase direction; it is the fastest clock and useful for normalization between different phase speeds. Along the local real direction,

$$dx_0 = \frac{d}{d\chi} \Re(\vec{\chi}) d\chi = \frac{\vec{\chi}^+ + \vec{\chi}^-}{2} d\chi = \cos \zeta d\chi =: d\tau. \quad (4.6)$$

Here $d\chi_0 := \cos \zeta d\chi$ is the projection of $d\chi$ onto the local real axis; in Sec. ?? we calibrate $d\tau = (1/\nu_0) d\chi_0$. The observed proper time of A relative to the rest observer is

$$\tilde{H}_A = \Re\left(\frac{d\vec{\chi}_A}{d\vec{\chi}_0}\right) = \cos \theta_A = \sqrt{1 - \sin^2 \theta_A} = \sqrt{1 - \frac{V^2}{c^2}} = \frac{1}{\gamma}. \quad (4.7)$$

4.3 Normalization

Let local time be parameterized by *phase*; introduce a reference frequency ν_0 and set

$$d\tau = \frac{1}{\nu_0} d\chi_0. \quad (4.8)$$

By the chain rule,

$$dx_0 = \tilde{H} d\chi = \frac{dx_0}{d\chi_0} \frac{d\chi_0}{d\tau} d\tau = \tilde{H} \dot{\chi} d\tau =: \dot{H} d\tau, \quad (4.9)$$

where $\nu := d\chi/d\tau$, $\dot{\chi} := \nu/\nu_0$, and $\dot{H} := \tilde{H} \dot{\chi}$. Choosing the calibration $\dot{H} \equiv c$ gives $dx_0 = c d\tau$. Similarly for space,

$$dx_l = \tilde{L} d\chi = \frac{dx_l}{d\chi_0} \frac{d\chi_0}{dl} dl = \tilde{L} \chi' dl =: L' dl, \quad \chi' := \frac{d\chi}{dl}. \quad (4.10)$$

From $dx_0 = dx_l$ for light one gets

$$c = \tilde{L}' \frac{dl}{d\tau}, \quad (4.11)$$

hence with temporal calibration to c the spatial scale becomes unit: $\tilde{L}' = 1$.

4.4 Light and c as a calibration constant

From the normalized forms,

$$\frac{c}{\dot{\chi}} d\chi = \frac{1}{\chi'} d\chi \quad \Rightarrow \quad c = \frac{\dot{\chi}}{\chi'} = \frac{dl}{d\tau}, \quad (4.12)$$

i.e. c is a *calibration constant* tying temporal and spatial measures, independent of local phase variation. Equation (??) also reads

$$c = \left(\frac{d\chi}{d\tau} \right) \left[\frac{dl}{d\chi} \right] \sim (\nu) [\lambda], \quad (4.13)$$

matching frequency and wavelength of a photon, with χ as its phase. For a lightlike trajectory,

$$ds^2 = c^2 \left(\frac{d\chi^2}{\dot{\chi}^2} - \frac{d\chi^2}{\dot{\chi}^2} \right) = 0. \quad (4.14)$$

At unit frequency, $\tau = \chi$: the photon's “proper time” is its phase, and the length of its phase-speed vector equals its wavelength, $\tilde{H}_p = \lambda$. Finally, the kinematic slope in phase coordinates is

$$\frac{dx_l}{dx_0} = \frac{\tilde{L} d\chi}{\tilde{H} d\chi} = \sin \theta = \frac{V}{c} \equiv \beta, \quad (4.15)$$

so $\theta = \pi/2$ implies $V = c$.

4.5 Lorentz factor via reparameterization

A change of direction of the phase speed transforms

$$\tilde{H}^2 = \tilde{S}^2 + \tilde{L}^2 \mapsto \dot{H}^2 = \dot{S}^2 + \dot{L}^2. \quad (4.16)$$

Lemma (parameter-change identity). The transition $\tilde{H} \rightarrow \dot{S}$ is the manifestation of evolving phase speed under the parameter change $\chi \mapsto \tau(\chi)$, with local Jacobian

$$\frac{d\tau}{d\chi} = \cos \zeta(\chi) \cos \theta(\chi) \quad \Rightarrow \quad \mathcal{J}(\zeta, \theta) := \frac{d\chi}{d\tau} = \frac{1}{\cos \zeta \cos \theta}. \quad (4.17)$$

Then

$$\dot{H} = \tilde{H} \mathcal{J}, \quad \dot{L} = \tilde{L} \mathcal{J}. \quad (4.18)$$

In differential form,

$$d \ln \dot{H} = d \ln \mathcal{J} = \tan \zeta d\zeta + \tan \theta d\theta. \quad (4.19)$$

For a *pure boost* ($d\zeta = 0$) one has $d\dot{H} = \dot{H} \tan \theta d\theta$. Absorbing a constant $\cos \zeta$ into the calibration (set $\zeta = 0$ henceforth), we obtain

$$\tilde{H}^2 = \dot{H}^2 - \dot{L}^2 = \sec^2 \theta (\tilde{H}^2 - \tilde{L}^2) = \gamma^2 (\tilde{H}^2 - \tilde{L}^2). \quad (4.20)$$

Corollary. In phase space the Euclidean norm \tilde{H} is conserved; in observed time the Minkowski norm \dot{S} is conserved; they are identical as quantities:

$$\boxed{\tilde{H} = \dot{S}}. \quad (4.21)$$

4.6 Rapidity and the phase angle

By definition,

$$\beta = \frac{V}{c} = \sin \theta, \quad \tanh \eta = \beta, \quad d\eta = \frac{d\beta}{1 - \beta^2}. \quad (4.22)$$

With $d\beta = \cos \theta d\theta$ and $1 - \beta^2 = \cos^2 \theta$,

$$d\eta = \sec \theta d\theta, \quad \eta(\theta) = \int \sec \theta d\theta = \ln |\sec \theta + \tan \theta| = \frac{1}{2} \ln \frac{|1 + \sin \theta|}{|1 - \sin \theta|}. \quad (4.23)$$

Fixing $\eta(0) = 0$,

$$e^{\eta(\theta)} = \sqrt{\frac{1 + \sin \theta}{1 - \sin \theta}}, \quad \gamma = \frac{1}{\sqrt{1 - \beta^2}} = \sec \theta = \cosh \eta. \quad (4.24)$$

Remark (groups). Observables satisfy $\beta = \sin \theta = \tanh \eta$ and $\gamma = \sec \theta = \cosh \eta$. Thus Euclidean rotations in the phase circle ($U(1)$ with angle θ) reproduce the numerical factors of hyperbolic boosts in $SO^+(1, 1)$ (rapidity η) *after* reparameterizing time. We do not claim an isomorphism $U(1) \cong SO(1, 1)$; only the equality of observable combinations under the change of parameter.

4.7 Velocity addition

Rapidity is additive:

$$\eta_{12} = \eta_1 + \eta_2, \quad \beta_{12} = \tanh(\eta_1 + \eta_2) = \frac{\beta_1 + \beta_2}{1 + \beta_1 \beta_2}. \quad (4.25)$$

Equivalently,

$$\gamma_{12} = \gamma_1 \gamma_2 (1 + \beta_1 \beta_2), \quad \gamma_{12} \beta_{12} = \gamma_1 \gamma_2 (\beta_1 + \beta_2). \quad (4.26)$$

4.8 Doppler shift

Define the observed frequency as the phase growth rate in the observer's proper time:

$$\nu := \frac{d\chi}{d\tau}. \quad (4.27)$$

For two successive wavefronts the phase increment is identical, hence

$$\frac{\nu_{\text{obs}}}{\nu_{\text{src}}} = \frac{d\chi/d\tau_{\text{obs}}}{d\chi/d\tau_{\text{src}}} = \frac{d\tau_{\text{src}}}{d\tau_{\text{obs}}}. \quad (4.28)$$

Longitudinal case: during $\gamma d\tau_{\text{src}}$ in the observer frame the source displaces by $\pm V \gamma d\tau_{\text{src}}$ (“+” receding, “−” approaching). Then

$$d\tau_{\text{obs}} = \gamma d\tau_{\text{src}} (1 \pm \beta), \quad \Rightarrow \quad \boxed{\frac{\nu_{\text{obs}}}{\nu_{\text{src}}} = \frac{1}{\gamma(1 \pm \beta)}}. \quad (4.29)$$

Equivalent forms (with $\beta = \sin \theta$, $\gamma = \sec \theta$ and rapidity η):

$$\frac{\nu_{\text{obs}}}{\nu_{\text{src}}} = \sqrt{\frac{1 \mp \beta}{1 \pm \beta}} = \sec \theta (1 \mp \sin \theta) = e^{\mp \eta}. \quad (4.30)$$

Transverse Doppler ($\varphi = 90^\circ$ in the observer's frame):

$$\frac{\nu_{\text{obs}}}{\nu_{\text{src}}} = \frac{1}{\gamma} = \cos \theta. \quad (4.31)$$

General line-of-sight (LOS) angle φ in the observer’s frame:

$$\boxed{\frac{\nu_{\text{obs}}}{\nu_{\text{src}}} = \gamma (1 - \beta \cos \varphi)} \quad (4.32)$$

Wavelength ratios are inverse to frequency ratios.

5 Discussion: links to known structures

Gauge phases. A global shift $\chi \mapsto \chi + \chi_0$ is unobservable. Allowing local reparameterizations $\chi \mapsto \chi + \alpha(x)$ induces a connection when comparing phases at different points. On wavefunctions $\psi \sim e^{i\chi}$ this is the familiar $U(1)$ gauge freedom $\psi \rightarrow e^{i\alpha(x)}\psi$ with $D_\mu = \partial_\mu - iA_\mu$ as the *phase-transport connection*.

Mass and the internal angle. With the decomposition by ζ , mass heuristically correlates with an irreducible real projection: massless objects have $\zeta = \pm\pi/2$ (no proper time; photon subspace), while massive objects have $|\zeta| < \pi/2$ (proper time exists). In the present paper we set $\zeta = 0$ in boost kinematics by calibration; a detailed mass-generation mechanism is left for future work.

Cosmological gauge. A natural global calibration of “absolute” time is the comoving frame with vanishing CMB dipole. This fixes a cosmological time t (FLRW) as a gauge, without affecting local Lorentz invariance; Doppler factors are then operationally referenced to that frame.

6 Conclusion

In unimetry, time and space are integrals of phase velocities; the Minkowski interval appears as a conserved quantity under parameter change. The core relations of SR— γ , rapidity, velocity addition, and Doppler factors—follow from elementary phase-plane geometry with a single rotation angle θ , while hyperbolic structure re-emerges upon reparameterizing time. The formalism is empirically equivalent to standard SR but can clarify causality and composition by treating all effects as projections of a single flow.

Outlook. Future directions include (i) a more explicit group-theoretic embedding, (ii) a rigorous treatment of the internal angle ζ and its relation to mass, and (iii) exploration of curved metrics as spatially varying Jacobians $\mathcal{J}(x)$ in the phase-to-observable map.

References

- [1] A. Einstein. Zur Elektrodynamik bewegter Körper. *Annalen der Physik*, 17:891–921, 1905. (English translation: On the electrodynamics of moving bodies.)
- [2] W. Rindler. *Relativity: Special, General, and Cosmological*. Oxford University Press, 2nd ed., 2006.
- [3] E. F. Taylor and J. A. Wheeler. *Spacetime Physics*. W. H. Freeman, 2nd ed., 1992.

6.1 Natural derivation of $\nu = k$ from the S^3 Hopf torus

We parametrize the 3-sphere $S_R^3 \subset \mathbb{R}^4$ by angles (ξ_0, ξ_1, ξ_2) :

$$\begin{aligned} x_1 &= R \cos \xi_0 \cos \xi_1, & x_2 &= R \cos \xi_0 \sin \xi_1, \\ x_3 &= R \sin \xi_0 \cos \xi_2, & x_4 &= R \sin \xi_0 \sin \xi_2, \end{aligned}$$

which induces the metric

$$ds^2 = R^2 (d\xi_0^2 + \cos^2 \xi_0 d\xi_1^2 + \sin^2 \xi_0 d\xi_2^2).$$

Fixing $\xi_0 = \xi_0^*$ produces a Hopf torus $\mathbb{T}^2 \subset S^3$ with radii

$$R_1 = R \cos \xi_0^*, \quad R_2 = R \sin \xi_0^*,$$

and flat induced metric $ds^2|_{\xi_0^*} = R_1^2 d\xi_1^2 + R_2^2 d\xi_2^2$. On this torus the free (geodesic) Lagrangian is

$$L = \frac{1}{2} (R_1^2 \dot{\xi}_1^2 + R_2^2 \dot{\xi}_2^2),$$

so that ξ_1, ξ_2 are cyclic coordinates with conserved momenta

$$p_1 = R_1^2 \dot{\xi}_1 = \text{const}, \quad p_2 = R_2^2 \dot{\xi}_2 = \text{const}.$$

Introduce arc-length variables along the two circles,

$$\chi = R_1 \xi_1, \quad \zeta = R_2 \xi_2, \quad \dot{\chi} = R_1 \dot{\xi}_1, \quad \dot{\zeta} = R_2 \dot{\xi}_2,$$

and choose the internal-time gauge of unimetry by fixing the speed along the “time” circle:

$$\dot{\zeta} \equiv 1.$$

Then

$$\frac{p_1}{p_2} = \frac{R_1^2 \dot{\xi}_1}{R_2^2 \dot{\xi}_2} = \frac{R_1}{R_2} \frac{\dot{\chi}}{\dot{\zeta}} = \frac{R_1}{R_2} \nu \quad \Rightarrow \quad \boxed{\nu = k, \quad k := \frac{p_1}{p_2} \frac{R_1}{R_2}}.$$

In the isotropic rest calibration $p_1 = p_2$ this reduces to $k = R_1/R_2$. For the Clifford torus $\xi_0^* = \pi/4$ one has $R_1 = R_2$ and hence $k = 1$.

GR outlook. In curved settings (slow spatial variation of $\xi_0(x)$ and nontrivial Hopf connection) the radii $R_{1,2}(x)$ and the ratio $k(x)$ become fields. The local clock rate

$$\nu(x) = k(x)$$

then encodes gravitational time dilation as a purely geometric effect of the torus radii/connection in the unimetry framework.

Notation. We deliberately use (ξ_0, ξ_1, ξ_2) to avoid conflicts with previously used angles (β, η, α) elsewhere in the manuscript.