



Immunotherapy for ovarian cancer: recent advances and perspectives

Emese Zsiros^a, Janos Tanyi^a, Klara Balint^a, and Lana E. Kandalaft^{a,b}

Purpose of review

Epithelial ovarian cancer is the most frequent cause of gynecologic cancer-related mortality in women, and prognosis for patients with recurrent or metastatic disease is extremely poor. Therefore, there is an enormous unmet need for the development of novel therapies in this indication. Although surgery and chemotherapy can improve survival rates, it is necessary to integrate alternative strategies, such as immunotherapy to improve the outcomes for patients with advanced ovarian cancer.

Recent findings

We will discuss the rationale of immunotherapy and some of the mechanisms of immunogenicity in ovarian cancer. We will highlight current results with cancer vaccines, adoptive T-cell therapy and immunomodulatory agents and will summarize the immune effects of selected chemotherapeutic agents, radiotherapy and recent results with combinatorial approaches in this disease setting. We will also discuss recent and potential future therapeutic interventions that might circumvent tumor-mediated immunosuppression.

Summary

Dramatic increase in the number of immunotherapy clinical trials was seen in the past decade with promising results in enhancing antitumor immune response and cancer vaccine efficacy. The future challenge for immunotherapy against ovarian cancer is to use a combinatorial approach to test rational, potentially synergistic immunotherapy combinations that can induce efficient antitumor immunity and prolong patients' survival.

Keywords

immunomodulation, immunotherapy, ovarian cancer

INTRODUCTION

Epithelial ovarian cancer (EOC) is the most common cause of gynecological cancer-associated death among women, with approximately 239 000 new cases to be diagnosed in the world in 2014 [1]. Cytoreductive surgery and platinum-based chemotherapy has remained the mainstay of ovarian cancer treatment and as the disease is discovered at an advanced stage, the prognosis is poor with a 5-year survival rate of 38%. There is a vital need for alternative treatments to increase the response rate and survival [2]. Recent scientific evidence demonstrated that EOC is an immunogenic tumor that can be recognized by the host immune system [3]. Spontaneous antitumor immune response of tumor-reactive T-cells and antibodies can be detected in peripheral blood, tumors and ascites of EOC patients with advanced disease [4,5]. These tumor-reactive T-cells are oligoclonal, recognize autologous tumor-associated antigens (TAAs) *in vitro* and exhibit tumor-specific cytolytic activity *ex vivo* [6]. The frequency of

serological responses to these antigens varies according to tumor type, stage or grade [7]. Our knowledge of TAAs expressed by EOC is still limited; some of the best-studied antigens in ovarian cancer patients are New York-esophageal-1 (NY-ESO-1), p53, human epidermal growth factor receptor 2 (Her2)/neu, survivin, folate receptor- α , sperm surface protein Sp17, WT1, MUC1, melanoma associated antigen-3 (MAGE3) and human telomerase reverse transcriptase (hTERT)

^aOvarian Cancer Research Center, Smilow Translational Research Center, Perelman School of Medicine of University of Pennsylvania Philadelphia, Pennsylvania, USA and ^bDepartment of Oncology, Ludwig Cancer Center, Centre Hospitalier Universitaire Vaudois, Lausanne, Switzerland

Correspondence to Lana E. Kandalaft, PharmD, PhD, MTR, Department of Oncology, Ludwig Cancer Center, Centre Hospitalier Universitaire Vaudois, Lausanne, Switzerland. Tel: +41 (0)21 314 78 23; e-mail: lknd@mail.med.upenn.edu, lana.kandalaft@chuv.ch

Curr Opin Oncol 2014, 26:492–500

DOI:10.1097/CCO.0000000000000111

KEY POINTS

- Ovarian cancer has been validated as a target for immunotherapy.
- Current clinical trials of cancer vaccine, adoptive T-cell transfer or immunomodulatory agents demonstrated that antitumor immunity can be augmented and can improve overall survival.
- New combinatorial approaches comprising of immunotherapy, chemotherapy and radiotherapy need to be tested to enhance therapeutic efficacy.

[8]. Most of these mutations detected in EOC are somatic, they are rarely shared among different tumors and exhibit an extreme degree of heterogeneity with an average of 60 private, nonsynonymous mutations per tumor in ovarian cancer [9], representing a major challenge for the development of a 'one-glove-fits-all' type of immunotherapy targeting all these mutations.

Despite the expression of TAAs by ovarian cancer, spontaneous antitumor immune response has only been demonstrated in approximately 55% of patients with ovarian cancer in the form of intraepithelial tumor-infiltrating lymphocytes (TILs) [10]. Coukos *et al.* reported that patients whose tumors had TILs experienced longer progression-free and overall survival [3,11–13]. On the contrary, immune evasion mechanisms in this patient population correlated with poor survival, such as CD4⁺CD25⁺FoxP3⁺ T regulatory (Treg) cells [14,15], programmed death ligand 1 (PD-L1 or B7-H1), a ligand for the immunosuppressive T-cell receptor programmed death 1 [16] and endothelin B receptor, which suppresses T-cell-endothelial adhesive interactions and T-cell homing to tumor [17,18]. Finally, a recent study demonstrated a strong correlation between low-immune gene signature expression and the development of high-risk tumors [19].

The association of TILs with prolonged survival, as well as the association of immune escape mechanisms with poor survival, suggests that ovarian cancer patients could respond to the same immunotherapy approaches, such as interleukin-2, cytotoxic T lymphocyte-associated protein 4 (CTLA-4) or programmed death-1 antibodies or adoptive transfer of ex-vivo expanded TIL, similarly to patients with other immunogenic tumors, such as melanoma [20]. In this review, we will discuss the mechanisms of immunogenicity in ovarian cancer, will highlight current results with cancer vaccines, adoptive T-cell therapy and immunomodulatory agents and will summarize the immune effects of selected chemotherapeutic agents, radiotherapy and recent results

with combinatorial approaches. We will also discuss recent and potential future therapeutic interventions that might circumvent tumor-mediated immunosuppression.

RATIONALE FOR IMMUNOMODULATORY AGENTS, OPPORTUNITIES AND CLINICAL RESULTS TO DATE

Harnessing the immune system to treat cancer is the major goal of immunotherapy. Although the identification of several tumor-specific antigens in ovarian cancer has provided the foundation for designing successful immunotherapies, it still has to overcome the immune escape of the tumor or cancer immunoevasion. The mechanisms for immunomodulation (therapeutic intervention aimed at modifying the body's immune response) in ovarian cancer include activation of professional antigen-presenting cells (APCs) by engaging costimulatory receptors (such as CD40), activation of effector T lymphocytes by immunostimulatory monoclonal antibodies (mAb) and finally depletion of Treg or immunosuppressive machineries (Fig. 1).

DENDRITIC CELL ACTIVATION

The initiation of an adaptive immune response requires the recognition of the antigen presented on a primary major histocompatibility complex (MHC) II on the surface of APCs by the T-cell antigen receptor, TCR (primary signal). Subsequently, secondary signals' costimulation with CD40 ligand (CD40L) expressed by activated T-cells engaging CD40 expressed by B cells and APCs are necessary for the secretion of cytokines (interleukin-1, interleukin-6, interleukin-8, interleukin-12, tumor necrosis factor- α and macrophage inflammatory protein-1 α) to further enhance, modify and skew the responding CD4⁺ to CD8⁺ effector cells [21**].

CD40-CD40L interactions facilitate dendritic cells to increase expression of CD80 and CD86 and thereby activate T-cells. CD40 antibody showed remarkable therapeutic activity in B-cell lymphomas with 80–100% of mice cured. CD40L antibody treatment was also found to inhibit tumor cell growth in an ovarian carcinoma cell line and freshly obtained ovarian carcinoma cells [22]. Several clinical CD40 antibodies, such as dacetuzumab and lucatumumab, have demonstrated long-term complete remissions in patients with advanced squamous cell cancer [23*] and clinical benefit in patients with melanoma [24].

The effect of CD40 antibodies has been tested in combination with other cancer immunotherapies, in particular interleukin-2 [25]. A fully human mAb




FIGURE 1. Opportunities of immune modulation in ovarian cancer. CTLA-4, cytotoxic T lymphocyte-associated protein 4; PD-L1, programmed death ligand 1; PD1, programmed death 1.

(CP-870,893)-recognizing CD40 has also been used in pancreatic cancer in combination with gemcitabine and induced major clinical remission [22]. Agonistic CD40 antibody was used in combination with toll-like receptor-3 (TLR3) agonist and demonstrated the transformation of dendritic cells from an immunosuppressive to an immunostimulatory cell type in a mouse model [26]; however, none of these therapies have been tested in ovarian carcinoma. These combinations merit further testing in patients with ovarian cancer.

Other dendritic cell activation mechanisms involve controlled pore glass (CpG) oligonucleotides. The immune system recognizes these oligonucleotides through TLR9 receptors that activate B-lymphocytes and dendritic cells. Intratumoral injection of CpG after radiation therapy promotes systemic lymphoma eradication in mice [27], with interesting results in humans [28]. This type of treatment is practical, well tolerated, and repeatable and warrants further investigation in ovarian cancer.

EFFECTOR T-CELL ACTIVATION

T-cell activation requires a primary signal through TCR and cognate antigen complex with MHC and a secondary signal, such as costimulatory CD28 and

inducible T-cell costimulator (ICOS). This activation is regulated by inhibitory receptors, such as CTLA-4 and programmed death-1. New therapies aimed at overcoming these mechanisms of peripheral tolerance by blocking the inhibitory checkpoints CTLA-4-CD80/86, programmed death-1, PD-L1, T-cell immunoglobulin-mucin-3-galectin-9 (TIM-3-GAL9) and lymphocyte-activation gene 3 (LAG3)-MHC-II with mAb to sustain the activation and proliferation of tumor-specific T-cells and to prevent energy or exhaustion resulting in an effective tumor-specific immune response [29[■]]. Additional check-point inhibitors include transforming growth factor beta (TGF β) and interleukin-10 [30].

The majority of CTLA-4 blockade (ipilimumab) clinical data come from studies in patients with melanoma [31,32]. In ovarian cancer, Hodi *et al.* [33] reported that one patient vaccinated with granulocyte-macrophage colony-stimulating factor modified irradiated autologous tumor cells and treated with ipilimumab demonstrated an objective response with durable remission for 4 years while receiving multiple infusions of CTLA-4 antibody. Tumor regression correlated with the CD8⁺/Treg ratio, suggesting that Treg depletion may provide a highly effective form of treatment when given in combination with the tumor vaccine and CTLA-4

antibody [33]. Future studies will help answer the role of anti-CTLA-4 therapy in ovarian cancer. A new phase II clinical trial is being conducted to evaluate ipilimumab as monotherapy in ovarian cancer patients in the United States of America (NCT01611558).

Programmed death-1 is an immunoglobulin superfamily surface molecule that upon binding to PD-L1 and PD-L2 promotes apoptosis in self-reacting T-cells through a mechanism that is distinct from CTLA-4 [34]. Elimination of the programmed death-1 pathway can result in the breakdown of immune tolerance against tumors. In preclinical studies, antibodies blocking PD-L1 or programmed death-1 profoundly enhanced the efficacy of immunotherapy in tumors with constitutive expression of PD-L1, including ovarian cancer. A phase I study using programmed death-1 blocking antibody (CT-011) demonstrated a 33% clinical benefit, including one patient with complete remission [35].

Dual blockade of programmed death-1 and CTLA-4 resulted in reversal of CD8⁺TIL dysfunction and led to tumor rejection in two-thirds of an ovarian cancer mouse model [36,37]. The potential of this combination has recently been highlighted in a phase I-II trial of ipilimumab and nivolumab (programmed death-1 antibody) in melanoma patients [38[•]], however yet to be explored in patients with ovarian cancer. Currently, a phase I trial is being conducted to investigate the safety and tolerability of anti-PD-L1 antibody as monotherapy in patients with metastatic or advanced solid tumors, including ovarian cancer patients (NCT01772004) [39].

Dual programmed death-1 and LAG-3 blockade has a synergistic effect on tumor clearance, and thus could be used as one of the future combinatorial immunotherapeutic approaches [40]. Dual blockade of these two molecules during T-cell priming efficiently augmented proliferation and cytokine production by NY-ESO-1-specific CD8⁺ T-cells in ovarian cancer patients [41]. A phase I trial is currently investigating the combination of anti-LAG-3 (BMS-986016) and programmed death-1 antibodies (BMS-936558) in melanoma and advanced solid tumors; however, its effect on ovarian cancer patients has not been tested.

T REGULATORY CELL DEPLETION

Tregs are potent immunosuppressive cells that promote progression of cancer through their ability to limit antitumor immunity and enhance angiogenesis. Several Treg-depleting strategies have been frequently used in clinical settings, such as metronomic low-dose oral (50 mg twice daily) or intravenous cyclophosphamide [42]. Cyclophosphamide

has been successfully combined with different types of cancer vaccines [43,44]. In a phase II trial, p53-synthetic long peptide vaccine was combined with low-dose cyclophosphamide and found to induce higher p53-specific responses compared with p53-synthetic long peptide monotherapy and produced a clinical response in 20% of patients [45].

Other approaches for Treg depletion include targeting the interleukin-2 receptor alpha chain, also known as CD25. Preclinical models showed that anti-CD25 monoclonal antibody treatment before vaccination led to complete tumor rejection and establishment of long-lasting tumor immunity with long-lasting CD4⁺CD25⁺Treg cells reduction [46]. Daclizumab, the Food and Drug Administration-approved, cell depleting, humanized immunoglobulin G1 (IgG1)-kappa mAb that binds specifically to CD25, has been well tolerated in patients, and demonstrated durable Treg suppression when in combination with hTERT peptide vaccine in breast cancer patients [47]. It is currently being evaluated in clinical trials for a variety of cancers, including ovarian [48].

ROLE OF NONIMMUNOTHERAPEUTICS IN IMMUNOMODULATION

Immunomodulatory effects on cytokine and cell-mediated responses can be induced also by non-immunotherapeutics, such as standard chemotherapy and radiotherapy.

STANDARD CHEMOTHERAPY

Chemotherapy is generally thought to act through selective killing of tumor cells or by irreversibly arresting their growth. Accumulating evidence indicates that several chemotherapeutics are more efficacious in immunocompetent versus immunodeficient hosts [49] and in fact have important 'off-target' immunologic effects to play.

Interestingly, treatment with Taxol (paclitaxel), an antimicrotubule agent, in advanced ovarian cancer was shown to upregulate the cytotoxic T-cell function that has been attributed to Taxol-induced tumor apoptosis and the accessibility of tumor antigens. Platinum-based therapies have also been shown to enhance the immunostimulatory potential of dendritic cells and decrease the immunosuppressive capacity of tumor cells through a STAT6-mediated pathway [50]. Moreover, an enhancement of the overall antitumor effect was observed in mice treated with an ovalbumin vaccine combined with cytotoxic doses of carboplatin and paclitaxel when compared with ovalbumin vaccine alone [51^{••}]. More recently, a clinical trial in advanced cervical

carcinoma demonstrated that depletion of myeloid suppressor cells by a carboplatin/paclitaxel chemotherapy standard regimen enhances the immune response to human papillomavirus vaccine, with sustained human papillomavirus 16-specific T-cell responses throughout several cycles of chemotherapy [52].

PEGylated liposomal doxorubicin (PLD) has also been shown to synergize with pleiomorphic immunomodulatory drugs, such as interleukin-18 [53], as well as vaccines and adoptive T-cell therapy. Facciabene *et al.* [54] have successfully combined PLD with a TLR-8 agonist (VTX-2337) in a humanized mouse model and have translated this to a phase I trial in advanced, platinum-resistant ovarian cancer patients (NCT01294293, $n=13$) in which the combination proved superior to historic data obtained with PLD monotherapy. This is at present tested through a randomized placebo-controlled, phase II trial comparing Doxil versus Doxil + VTX-2337 [Gynecological Oncology Group (GOG)-3003, NCT01666444]. Other agents include 5-fluorouracil, which rendered human colon carcinoma cells more sensitive to lysis by tumor-specific cytotoxic T-cells [55]. Bevacizumab, which is a humanized monoclonal antibody that neutralizes human vascular endothelial growth factor, also has a role in immunomodulation as it restores dendritic cell function and enhances antitumor immune responses and the efficacy of tumor vaccines.

Understanding the immunological events occurring in both animal models and patients undergoing chemotherapy should guide decisions about the development of appropriate combinations and scheduling for the integration of chemotherapy with immunotherapy, which can certainly improve therapeutic management of this disease.

RADIOTHERAPY

Radiotherapy is perceived as an immunosuppressive modality; however, recent data from multiple cancer models have provided plentiful evidence that some of the effects of ionizing radiation are recognized as powerful adjuvant to systemic antitumor immunity. Radiotherapy is able to induce an immune-mediated abscopal effect whereby the radiation causes indirect antitumor effects in non-treated areas, destroying tumors outside the radiation field. The abscopal effect is mediated by the increased cell surface expression of MHC class I molecules, cytokine release of interferon- β and tumor necrosis factor α and increased priming of antigen-specific dendritic cells and by stimulation of systemic innate (natural killer cell-mediated) and adaptive (dendritic cell and cytotoxic T lymphocyte-mediated) immunity against the tumor [56].

In preclinical models, promising results have been obtained by combining radiotherapy with targeted interventions, such as anti-CTLA-4, dendritic cell vaccines and adoptive T-cell therapy. First, the combination of subcutaneous granulocyte-macrophage colony-stimulating factor with local radiotherapy was tested in patients with metastatic solid tumors [57], and as a result, an abscopal response was detected in 30% of the patients. Fractionated radiotherapy given as neoadjuvant was combined with dendritic cell vaccine in sarcoma patients and showed remarkable progression-free survival results [58]. Similar results were seen in glioblastoma patients with a combination approach of dendritic cell vaccine, radiotherapy and chemotherapy [59]. The combination of radiotherapy with ipilimumab has also been tested in a phase I/II trial in patients with metastatic prostate cancer [60]. Randomized phase III trials are underway to further test the possible benefits of this combination. The optimal radiation regimens (doses and fractionations) to be used for immunomodulation remain to be defined [61]. In ovarian cancer, only radioactively labelled antibody therapies, such as the radionuclide therapy with (90)Y- or (177)Lu-CC49 [62], have been tested so far. The combination of radiation therapy and one or more immune-based modalities should be a promising synergistic approach to providing long-term survival with minimal toxicity.

RATIONALE FOR CELLULAR-BASED THERAPIES

Cell-based therapy is founded on the facts that cells can perform complex biological functions and therapeutic tasks, can have exquisite sensitivity and specificity and can be engineered to increase efficacy [63]. (Different approaches are depicted in Fig. 2.)

VACCINES

Consistent with experience in other immunogenic tumors [64], vaccines have shown limited efficacy as monotherapy in patients with advanced recurrent disease but the results are notable and provide basis for further optimization. In a retrospective review of patients treated in the adjuvant setting after secondary complete response, Sabbatini *et al.* [65] demonstrated that patients vaccinated with monovalent or heptavalent vaccines against carbohydrate epitopes experienced significantly longer time to progression and higher progression-free survival rates relative to controls treated with alternative consolidation therapies. Various phase I studies in which advanced ovarian cancer patients were vaccinated with different vaccines including antiidiotype ACA-125 (antiidiotypic cancer antigen-125), an

analog of cancer antigen-125 (CA-125) [66], carcino-embryonic antigen-mucin-1-triad of costimulatory molecules (CEA-MUC-1-TRICOM) poxviral-based vaccine [67], HER2 vaccine [68] and p53 peptide antigen vaccine [45,69] have resulted in improved survival and the induction of antigen-specific T-cell and humoral immunity. Odunsi *et al.* [70] used recombinant poxviruses (vaccinia and fowlpox)-expressing tumor-associated antigens (NY-ESO-1) as cancer vaccines to induce tumor-specific immune response in 22 EOC patients with advanced disease with a median progression-free survival of 21 months (95% confidence interval, 16–29 months), and median overall survival of 48 months. However, a phase III placebo-controlled maintenance therapy trial of abagovomab (anti-CA-125) 2mg versus placebo failed to prolong progression-free survival or overall survival [71²²]. An alternative approach to vaccines directed toward specific antigens is whole tumor antigen vaccines created using tumor cells, autologous tumor lysate or tumor-derived RNA (Fig. 2) [72–74]. We and others have also shown objective responses in recurrent advanced ovarian cancer patients when vaccinated with dendritic cell-based whole tumor vaccination [75], or viral oncolysate vaccine generated from ovarian cancer cell lines infected with influenza-A virus [76,77] or with autologous tumor cells infected with Newcastle disease virus [78].

A major limitation of cancer vaccines stems from the inability to elicit a rapid and overwhelming T-cell response, which is required to reject established

tumors. This problem is magnified in ovarian cancer by the paucity of well characterized rejection antigens and by the significant molecular heterogeneity of the disease [79]. Even when a defined target is available, and vaccination successfully induces an immune response, the long-term benefit can be limited by tumor evolution. In a recent study, one patient experienced complete objective response to NY-ESO-1 peptide vaccine, but later recurred with a NY-ESO-1-negative tumor, proving that single-target immunization can result in immune escape tumor variants following initial response [80]. Recent advances in the clinical application of immunotherapy suggest that immunotherapy with ‘personalized’ private antigens (that arise from mutations) could also be expected to induce rapid and strong secondary immune responses (reviewed in [81,82]). The current view is that both approaches, targeting public or targeting private antigens, can be beneficial. However, to increase the clinical benefits one should choose a patient population with small volume or no disease with good performance status and should vaccinate in combination with other immunomodulatory agents to maximize the efficacy of the vaccine.

ADOPTIVE T-CELL THERAPY

To be effective, cancer immunotherapy is dependent on the presence of sufficient numbers of anti-tumor lymphocytes with appropriate homing and effector functions that enable them to seek out and destroy cancer cells *in vivo*. The adoptive transfer of ex-vivo expanded, tumor-reactive T-cells holds the




FIGURE 2. Immunotherapeutic approaches for cellular therapies. 1, tumor infiltrating T-cells; 2, dendritic cell vaccines; 3, genetically engineered T-cells.

potential of achieving this for the treatment of established tumors [82].

In an early pilot T-cell transfer trial in which autologous tumor-infiltrating lymphocytes were administered after surgical resection and cisplatin chemotherapy, the disease-free survival and overall survival was found to be prolonged in ovarian cancer patients (Fig. 2) [83]. In another study, administration of TILs (alone or in combination with chemotherapy) was shown to induce objective cancer regressions [84]. Although TILs represent the candidate tumor-reactive cell population for adoptive immunotherapy, alternative approaches have been employed to generate large numbers of tumor-reactive CD8⁺ and CD4⁺ T-cells against ovarian cancer. Killing of ovarian cancer cells can be achieved by retargeting of T lymphocytes using F (Ab')₂ fragments of a bispecific monoclonal antibody (OC/TR), directed to the CD3 molecule on T lymphocytes and to the folate receptor on ovarian carcinoma cells [85]. Intraperitoneal infusions of activated autologous OC/TR-retargeted peripheral blood T-cells with recombinant interleukin-2 induced objective tumor regression in three out of 19 patients [86], with detectable OC/TR-directed lymphocytes in the peritoneal cavity of treated patients after infusion.

We have recently reported a phase I study of a combinatorial approach encompassing dendritic cell-based autologous whole tumor vaccination and antiangiogenesis therapy, followed by the adoptive transfer of autologous vaccine-primed CD3/CD28-costimulated lymphocytes [75]. Three patients with residual measurable disease who have been previously vaccinated with a whole tumor lysate vaccine received outpatient lymphodepletion and adoptive T-cell transfer, which was well tolerated and resulted in a durable reduction of circulating regulatory T-cells and in increased CD8⁺ lymphocyte counts. The vaccine-induced restoration of antitumor immunity was achieved in two individuals, who also demonstrated clinical benefits, including one complete response.

Adoptive T-cell therapy can become more effective and powerful by genetically engineering patients' lymphocytes, endowing them with more tumor specificity (Fig. 2). Genes used to modify T-cells include those encoding TCRs and chimeric antigen receptors (CARs). TCR-based engineering represents a compelling strategy for ovarian cancer therapy as TCRs that recognize human leukocyte antigen-A2 restricted epitopes from known ovarian cancer antigens, such as NY-ESO-1, p53 and others [87].

Engineering T-cells with redirected specificity to recognize antigens in an MHC-unrestricted fashion can be achieved through the use of CARs. In this

case, T-cells are transduced with fusion genes encoding an extracellular domain that specifically binds to tumor epitopes through a single-chain variable fragment antibody, linked to intracellular signaling modules that mediate T-cell activation [88]. Some of the generated CARs, which have been investigated *in vitro* and *in vivo* and are relevant to ovarian cancer, are folate receptor-alpha, MUC-16, HER-2 [89,90] and mesothelin [91]. One study of adoptive transfer of CARs in ovarian cancer demonstrated safety but showed no clinical response because of low expression of the transgenic CAR and poor persistence of the transferred T-cells [85]. Improved success of CARs in the clinic requires a panel of bioengineered T-cells with different specificities, custom-made for each individual, which is technically and economically challenging.

CONCLUSION

In the past decades, we have seen a dramatic increase in the number of immunotherapy clinical trials to enhance antitumor immune response and cancer vaccine efficacy. Sufficient evidence indicates that ovarian cancers are indeed immunogenic tumors and excellent candidates for immunotherapy. Both passive and active immunotherapeutic modalities have shown potential clinical benefit in at least a subset of these patients. Future challenge for immunotherapy against ovarian cancer is to use a combinatorial approach to test rational immunomodulatory combinations that can induce efficient antitumor immunity that may achieve prolonged patient survival. The other major challenge for cellular immunotherapeutics is the development of manufacturing technologies that are less costly and do not require a very complex infrastructure so therapy can be accessible by the masses.

Acknowledgements

This study was supported by National Institution of Health R01 grant FD003520-02, the Ovarian Cancer Immunotherapy Initiative and the Marcus Foundation.

Conflicts of interest

The authors declare no conflict of interest.

REFERENCES AND RECOMMENDED READING

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Jemal A, Siegel R, Ward E, *et al.* Cancer statistics, 2009. *CA Cancer J Clin* 2009; 59:225–249.
2. Vaughan S, Coward JI, Bast RC, *et al.* Rethinking ovarian cancer: recommendations for improving outcomes. *Nat Rev Cancer* 2011; 11:719–725.

3. Zhang L, Conejo-Garcia JR, Katsaros D, *et al.* Intratumoral T cells, recurrence, and survival in epithelial ovarian cancer. *N Engl J Med* 2003; 348:203–213.
4. Schlienger K, Chu CS, Woo EY, *et al.* TRANCE- and CD40 ligand-matured dendritic cells reveal MHC class I-restricted T cells specific for autologous tumor in late-stage ovarian cancer patients. *Clin Cancer Res* 2003; 9:1517–1527.
5. Goodell V, Salazar LG, Urban N, *et al.* Antibody immunity to the p53 oncogenic protein is a prognostic indicator in ovarian cancer. *J Clin Oncol* 2006; 24:762–768.
6. Santin AD, Hermonat PL, Ravaggi A, *et al.* In vitro induction of tumor-specific human lymphocyte antigen class I-restricted CD8 cytotoxic T lymphocytes by ovarian tumor antigen-pulsed autologous dendritic cells from patients with advanced ovarian cancer. *Am J Obstet Gynecol* 2000; 183:601–609.
7. Gnjatich S, Ritter E, Buchler MW, *et al.* Seromic profiling of ovarian and pancreatic cancer. *Proc Natl Acad Sci U S A* 2010; 107:5088–5093.
8. Reuschenbach M, von Knebel Doeberitz M, Wentzensen N. A systematic review of humoral immune responses against tumor antigens. *Cancer Immunol Immunother* 2009; 58:1535–1544.
9. Cancer Genome Atlas Research Network. Integrated genomic analyses of ovarian carcinoma. *Nature* 2011; 474:609–615.
10. Chatterjee M, Wojciechowski J, Tainsky MA. Discovery of antibody biomarkers using protein microarrays of tumor antigens cloned in high throughput. *Methods Mol Biol* 2009; 520:21–38.
11. Hwang WT, Adams SF, Tahirovic E, *et al.* Prognostic significance of tumor-infiltrating T cells in ovarian cancer: a meta-analysis. *Gynecol Oncol* 2012; 124:192–198.
12. Bachmayr-Heyda A, Aust S, Heinze G, *et al.* Prognostic impact of tumor-infiltrating CD8+ T cells in association with cell proliferation in ovarian cancer patients—a study of the OVCAD consortium. *BMC Cancer* 2013; 13:422.
13. Adams SF, Levine DA, Cadungog MG, *et al.* Intraepithelial T cells and tumor proliferation: impact on the benefit from surgical cytoreduction in advanced serous ovarian cancer. *Cancer* 2009; 115:2891–2902.
14. Curiel TJ, Coukos G, Zou L, *et al.* Specific recruitment of regulatory T cells in ovarian carcinoma fosters immune privilege and predicts reduced survival. *Nat Med* 2004; 10:942–949.
15. Sato E, Olson SH, Ahn J, *et al.* Intraepithelial CD8+ tumor-infiltrating lymphocytes and a high CD8+regulatory T cell ratio are associated with favorable prognosis in ovarian cancer. *Proc Natl Acad Sci U S A* 2005; 102:18538–18543.
16. Hamanishi J, Mandai M, Iwasaki M, *et al.* Programmed cell death 1 ligand 1 and tumor-infiltrating CD8+ T lymphocytes are prognostic factors of human ovarian cancer. *Proc Natl Acad Sci U S A* 2007; 104:3360–3365.
17. Buckanovich RJ, Facciabene A, Kim S, *et al.* Endothelin B receptor mediates the endothelial barrier to T cell homing to tumors and disables immune therapy. *Nat Med* 2008; 14:28–36.
18. Kandalafi LE, Facciabene A, Buckanovich RJ, Coukos G. Endothelin B receptor, a new target in cancer immune therapy. *Clin Cancer Res* 2009; 15:4521–4528.
19. Japanese Serous Ovarian Cancer Study Group. Yoshihara K, Tsunoda T, Shigemizu D, *et al.* High-risk ovarian cancer based on 126-gene expression signature is uniquely characterized by downregulation of antigen presentation pathway. *Clin Cancer Res* 2012; 18:1374–1385.
20. Rosenberg SA, Dudley ME. Adoptive cell therapy for the treatment of patients with metastatic melanoma. *Curr Opin Immunol* 2009; 21:233–240.
21. Fehres CM, Unger WW, Garcia-Vallejo JJ, van Kooyk Y. Understanding the biology of antigen cross-presentation for the design of vaccines against cancer. *Front Immunol* 2014; 5:149.
- Thorough review of the mechanisms of immune response and helps to understand vaccine design.
22. Beatty GL, Torigian DA, Chiorean EG, *et al.* A phase I study of an agonist CD40 monoclonal antibody (CP-870,893) in combination with gemcitabine in patients with advanced pancreatic ductal adenocarcinoma. *Clin Cancer Res* 2013; 19:6286–6295.
23. Vonderheide RH, Glennie MJ. Agonistic CD40 antibodies and cancer therapy. *Clin Cancer Res* 2013; 19:1035–1043.
- Comprehensive review of CD40 in cancer therapy
24. Vonderheide RH, Flaherty KT, Khalil M, *et al.* Clinical activity and immune modulation in cancer patients treated with CP-870,893, a novel CD40 agonist monoclonal antibody. *J Clin Oncol* 2007; 25:876–883.
25. Jackaman C, Nelson DJ. Intratumoral interleukin-2/agonist CD40 antibody drives CD4+ independent resolution of treated-tumors and CD4+ dependent systemic and memory responses. *Cancer Immunol Immunother* 2012; 61:549–560.
26. Scarlett UK, Cubillos-Ruiz JR, Nesbeth YC, *et al.* In situ stimulation of CD40 and Toll-like receptor 3 transforms ovarian cancer-infiltrating dendritic cells from immunosuppressive to immunostimulatory cells. *Cancer Res* 2009; 69:7329–7337.
27. Houot R, Levy R. T-cell modulation combined with intratumoral CpG cures lymphoma in a mouse model without the need for chemotherapy. *Blood* 2009; 113:3546–3552.
28. Brody JD, Ai WZ, Czerwinski DK, *et al.* In situ vaccination with a TLR9 agonist induces systemic lymphoma regression: a phase I/II study. *J Clin Oncol* 2010; 28:4324–4332.
29. Perez-Gracia JL, Labiano S, Rodriguez-Ruiz ME, *et al.* Orchestrating immune check-point blockade for cancer immunotherapy in combinations. *Curr Opin Immunol* 2014; 27:89–97.
- Comprehensive review of immune check-point blockade in cancer immunotherapy.
30. Sun H, Sun C, Xiao W. Expression regulation of co-inhibitory molecules on human natural killer cells in response to cytokine stimulations. *Cytokine* 2014; 65:33–41.
31. Fong L, Small EJ. Anti-cytotoxic T-lymphocyte antigen-4 antibody: the first in an emerging class of immunomodulatory antibodies for cancer treatment. *J Clin Oncol* 2008; 26:5275–5283.
32. Hodi FS, O'Day SJ, McDermott DF, *et al.* Improved survival with ipilimumab in patients with metastatic melanoma. *N Engl J Med* 2010; 363:711–723.
33. Hodi FS, Butler M, Oble DA, *et al.* Immunologic and clinical effects of antibody blockade of cytotoxic T lymphocyte-associated antigen 4 in previously vaccinated cancer patients. *Proc Natl Acad Sci U S A* 2008; 105:3005–3010.
34. Pedoeem A, Azoulay-Alfaguter I, Strazza M, *et al.* Programmed death-1 pathway in cancer and autoimmunity. *Clin Immunol* 2014; 153:145–152.
35. Berger R, Rotem-Yehudar R, Slama G, *et al.* Phase I safety and pharmacokinetic study of CT-011, a humanized antibody interacting with PD-1, in patients with advanced hematologic malignancies. *Clin Cancer Res* 2008; 14:3044–3051.
36. Govindaraj C, Scalzo-Inguanti K, Madondo M, *et al.* Impaired Th1 immunity in ovarian cancer patients is mediated by TNFR2+ Tregs within the tumor microenvironment. *Clin Immunol* 2013; 149:97–110.
37. Duraiswamy J, Freeman G, Coukos G. Replenish the source within: Rescuing tumor-infiltrating lymphocytes by double checkpoint blockade. *Oncoimmunology* 2013; 2:e25912.
38. Wolchok JD, Kluger H, Callahan MK, *et al.* Nivolumab plus ipilimumab in advanced melanoma. *N Engl J Med* 2013; 369:122–133.
- This study demonstrated prolonged survival in patients treated with dual blocking of CTLA-4 and programmed death-1.
39. ClinicalTrials.gov. NCT01772004. A phase I, open-label, multiple-ascending dose trial to investigate the safety, tolerability, pharmacokinetics, biological and clinical activity of MSB0010718c in subjects with metastatic or locally advanced solid tumors and expansion to selected indications: NIH; 2014 [cited 1 May 2014]. Available from: <http://www.clinicaltrials.gov/ct2/show/NCT01772004?term=NCT01772004&rank=1>.
40. Turnis ME, Korman AJ, Drake CG, Vignali DA. Combinatorial Immunotherapy: PD-1 may not be LAG-ing behind any more. *Oncoimmunology* 2012; 1:1172–1174.
41. Matsuzaki J, Gnjatich S, Mhawech-Fauceglia P, *et al.* Tumor-infiltrating NY-ESO-1-specific CD8+ T cells are negatively regulated by LAG-3 and PD-1 in human ovarian cancer. *Proc Natl Acad Sci U S A* 2010; 107:7875–7880.
42. Le DT, Jaffee EM. Regulatory T-cell modulation using cyclophosphamide in vaccine approaches: a current perspective. *Cancer Res* 2012; 72:3439–3444.
43. Ellebaek E, Engell-Noerregaard L, Iversen TZ, *et al.* Metastatic melanoma patients treated with dendritic cell vaccination, Interleukin-2 and metronomic cyclophosphamide: results from a phase II trial. *Cancer Immunol Immunother* 2012; 61:1791–1804.
44. Camisaschi C, Filipazzi P, Tazzari M, *et al.* Effects of cyclophosphamide and IL-2 on regulatory CD4+ T cell frequency and function in melanoma patients vaccinated with HLA-class I peptides: impact on the antigen-specific T cell response. *Cancer Immunol Immunother* 2013; 62:897–908.
45. Vermeij R, Leffers N, Hoogeboom BN, *et al.* Potentiation of a p53-SLP vaccine by cyclophosphamide in ovarian cancer: a single-arm phase II study. *Int J Cancer* 2012; 131:E670–E680.
46. Powell DJ Jr, Attia P, Ghetie V, *et al.* Partial reduction of human FOXP3+ CD4 T cells in vivo after CD25-directed recombinant immunotoxin administration. *J Immunother* 2008; 31:189–198.
47. Rech AJ, Vonderheide RH. Clinical use of anti-CD25 antibody daclizumab to enhance immune responses to tumor antigen vaccination by targeting regulatory T cells. *Ann N Y Acad Sci* 2009; 1174:99–106.
48. Vasievich EA, Huang L. The suppressive tumor microenvironment: a challenge in cancer immunotherapy. *Mol Pharm* 2011; 8:635–641.
49. Apetoh L, Ghiringhelli F, Tesniere A, *et al.* Toll-like receptor 4-dependent contribution of the immune system to anticancer chemotherapy and radiotherapy. *Nat Med* 2007; 13:1050–1059.
50. Hato SV, de Vries IJ, Lesterhuis WJ. STATing the importance of immune modulation by platinum chemotherapeutics. *Oncoimmunology* 2012; 1:234–236.
51. Diaz Y, Tundidor Y, Lopez A, Leon K. Concomitant combination of active immunotherapy and carboplatin- or paclitaxel-based chemotherapy improves antitumor response. *Cancer Immunol Immunother* 2013; 62:455–469.
- This study demonstrated that carboplatin and paclitaxel concomitantly combined with active immunotherapies is beneficial for advanced cancer patients.
52. Van Der Burgsh, van Meir H, Kroep JR, *et al.* Synergistic effects of properly timed HPV16 synthetic long peptide vaccination during standard carboplatin-paclitaxel chemotherapy in animals and in patients with metastatic cervical carcinoma. Conference Abstract #2938, 105th Annual meeting of American Association for Cancer Research, San Diego, CA, April 2014.
53. Alagkiozidis I, Facciabene A, Carpenito C, *et al.* Increased immunogenicity of surviving tumor cells enables cooperation between liposomal doxorubicin and IL-18. *J Transl Med* 2009; 7:104.

54. Facciabene A, Malykhin A, Mathews M, et al. TLR8 agonist and Doxil chemotherapy potentially activate human antitumor immune response in a human immune system mouse model. 102nd Annual meeting of AACR, Orlando FL, 2011, Abstract# 4575. *Cancer Res* 2011; 71(8 Suppl 1). doi: 10.1158/1538-7445.AM2011-4745.
55. Bergmann-Leitner ES, Abrams SI. Treatment of human colon carcinoma cell lines with antineoplastic agents enhances their lytic sensitivity to antigen-specific CD8+ cytotoxic T lymphocytes. *Cancer Immunol Immunother* 2001; 50:445–455.
56. Ahmed MM, Hodge JW, Guha C, et al. Harnessing the potential of radiation-induced immune modulation for cancer therapy. *Cancer Immunol Res* 2013; 1:280–284.
57. Formenti SC, Demaria S. Systemic effects of local radiotherapy. *Lancet Oncol* 2009; 10:718–726.
58. Finkelstein SE, Icloaz C, Bui MM, et al. Combination of external beam radiotherapy (EBRT) with intratumoral injection of dendritic cells as neo-adjuvant treatment of high-risk soft tissue sarcoma patients. *Int J Radiat Oncol Biol Phys* 2012; 82:924–932.
59. Pellegatta S, Eoli M, Frigerio S, et al. The natural killer cell response and tumor debulking are associated with prolonged survival in recurrent glioblastoma patients receiving dendritic cells loaded with autologous tumor lysates. *Oncoimmunology* 2013; 2:e23401.
60. Slovin SF, Higano CS, Hamid O, et al. Ipilimumab alone or in combination with radiotherapy in metastatic castration-resistant prostate cancer: results from an open-label, multicenter phase I/II study. *Ann Oncol* 2013; 24:1813–1821.
61. Schae D, Ratikan JA, Iwamoto KS, McBride WH. Maximizing tumor immunity with fractionated radiation. *Int J Radiat Oncol Biol Phys* 2012; 83:1306–1310.
62. Meredith R, You Z, Alvarez R, et al. Predictors of long-term outcome from intraperitoneal radioimmunotherapy for ovarian cancer. *Cancer Biother Radiopharm* 2012; 27:36–40.
63. Fischbach MA, Bluestone JA, Lim WA. Cell-based therapeutics: the next pillar of medicine. *Sci Transl Med* 2013; 5:179ps7.
- Comprehensive review of cell-based therapies
64. Rosenberg SA, Yang JC, Restifo NP. Cancer immunotherapy: moving beyond current vaccines. *Nat Med* 2004; 10:909–915.
65. Sabbatini P, Spriggs D, Aghajanian C, et al. Consolidation strategies in ovarian cancer: observations for future clinical trials. *Gynecol Oncol* 2009; 116:66–71.
66. Reinartz S, Kohler S, Schlebusch H, et al. Vaccination of patients with advanced ovarian carcinoma with the antiidiotype ACA125: immunological response and survival (phase Ib/II). *Clin Cancer Res* 2004; 10:1580–1587.
67. Gulley JL, Arlen PM, Tsang KY, et al. Pilot study of vaccination with recombinant CEA-MUC-1-TRICOM poxviral-based vaccines in patients with metastatic carcinoma. *Clin Cancer Res* 2008; 14:3060–3069.
68. Disis ML, Goodell V, Schiffman K, Knutson KL. Humoral epitope-spreading following immunization with a HER-2/neu peptide based vaccine in cancer patients. *J Clin Immunol* 2004; 24:571–578.
69. Leffers N, Lambeck AJ, Gooden MJ, et al. Immunization with a P53 synthetic long peptide vaccine induces P53-specific immune responses in ovarian cancer patients, a phase II trial. *Int J Cancer* 2009; 125:2104–2113.
70. Odunsi K, Matsuzaki J, Karbach J, et al. Efficacy of vaccination with recombinant vaccinia and fowlpox vectors expressing NY-ESO-1 antigen in ovarian cancer and melanoma patients. *Proc Natl Acad Sci U S A* 2012; 109:5797–5802.
71. Sabbatini P, Harter P, Scambia G, et al. Abagovomab as maintenance therapy in patients with epithelial ovarian cancer: a phase III trial of the AGO OVAR, COG, GINECO, and GEICO—the MIMOSA study. *J Clin Oncol* 2013; 31:1554–1561.
- This study evaluated the effect of anti-CA-125 antibody in ovarian cancer and showed favorable safety profile.
72. Chianese-Bullock KA, Irvin WP Jr, Petroni GR, et al. A multipptide vaccine is safe and elicits T-cell responses in participants with advanced stage ovarian cancer. *J Immunother* 2008; 31:420–430.
73. Tsuda N, Mochizuki K, Harada M, et al. Vaccination with predesignated or evidence-based peptides for patients with recurrent gynecologic cancers. *J Immunother* 2004; 27:60–72.
74. Chu C, June BJ, Boyer et al., Autologous dendritic cell (IDD-6) vaccination as consolidation for advanced ovarian cancer. In: Annual Meeting on Women's Cancer; Society of Gynecologic Oncologists; 2014. *Gynecol Oncol* 2014; 134. [Abstract]
75. Chiang CL, Kandalaf LE, Tanyi J, et al. A dendritic cell vaccine pulsed with autologous hypochlorous acid-oxidized ovarian cancer lysate primes effective broad antitumor immunity: from bench to bedside. *Clin Cancer Res* 2013; 19:4801–4815.
76. Ioannides CG, Platsoucas CD, Freedman RS. Immunological effects of tumor vaccines: II. T cell responses directed against cellular antigens in the viral oncolysates. *In Vivo* 1990; 4:17–24.
77. Ioannides CG, Platsoucas CD, Pateria R, et al. T-cell functions in ovarian cancer patients treated with viral oncolysates: I. Increased helper activity to immunoglobulins production. *Anticancer Res* 1990; 10:645–653.
78. Schirmacher V. Clinical trials of antitumor vaccination with an autologous tumor cell vaccine modified by virus infection: improvement of patient survival based on improved antitumor immune, memory. *Cancer Immunol Immunother* 2004; 54:587–598.
79. Carter SL, Cibulskis K, Helman E, et al. Absolute quantification of somatic DNA alterations in human cancer. *Nat Biotech* 2012; 30:413–421.
80. Odunsi K, Qian F, Matsuzaki J, et al. Vaccination with an NY-ESO-1 peptide of HLA class I/II specificities induces integrated humoral and T cell responses in ovarian cancer. *Proc Natl Acad Sci U S A* 2007; 104:12837–12842.
81. Gulley JL. Therapeutic vaccines: the ultimate personalized therapy? *Hum Vaccin Immunother* 2013; 9:219–221.
82. June CH. Adoptive T cell therapy for cancer in the clinic. *J Clin Invest* 2007; 117:1466–1476.
83. Fujita K, Ikarashi H, Takakuwa K, et al. Prolonged disease-free period in patients with advanced epithelial ovarian cancer after adoptive transfer of tumor-infiltrating lymphocytes. *Clin Cancer Res* 1995; 1:501–507.
84. Aoki Y, Takakuwa K, Kodama S, et al. Use of adoptive transfer of tumor-infiltrating lymphocytes alone or in combination with cisplatin-containing chemotherapy in patients with epithelial ovarian cancer. *Cancer Res* 1991; 51:1934–1939.
85. Kershaw MH, Westwood JA, Parker LL, et al. A phase I study on adoptive immunotherapy using gene-modified T cells for ovarian cancer. *Clin Cancer Res* 2006; 12 (20 Pt 1):6106–6115.
86. Freedman RS, Platsoucas CD, Deisseroth AB. Use of a retroviral vector to study the trafficking patterns of purified ovarian tumor infiltrating lymphocytes (TIL) used in intraperitoneal adoptive immunotherapy of ovarian cancer patients. A pilot study. *Hum Gene Ther* 1996; 7:1351–1365.
87. Riley JL, June CH, Blazar BR. Human T regulatory cell therapy: take a billion or so and call me in the morning. *Immunity* 2009; 30:656–665.
88. Sadelain M, Brentjens R, Riviere I. The promise and potential pitfalls of chimeric antigen receptors. *Curr Opin Immunol* 2009; 21:215–223.
89. Lanitis E, Dangaj D, Hagemann IS, et al. Primary human ovarian epithelial cancer cells broadly express HER2 at immunologically-detectable levels. *PLoS One* 2012; 7:e49829.
90. Kandalaf LE, Powell DJ Jr, Coukos G. A phase I clinical trial of adoptive transfer of folate receptor-alpha redirected autologous T cells for recurrent ovarian cancer. *J Transl Med* 2012; 10:157.
91. Chekmasova AA, Rao TD, Nikhamin Y, et al. Successful eradication of established peritoneal ovarian tumors in SCID-Beige mice following adoptive transfer of T cells genetically targeted to the MUC16 antigen. *Clin Cancer Res* 2010; 16:3594–3606.