

The evolution of checkpoint blockade as a cancer therapy: what's here, what's next?

Daniel Sanghoon Shin^{1,5} and Antoni Ribas^{1,2,3,4,5}



Unleashing the immune system to fight cancer has become one of the main treatment modalities since the anti-CTLA-4 antibody, ipilimumab was approved for patients with advanced melanoma in 2011. Pembrolizumab and nivolumab, two anti-PD-1 antibodies recently approved for the treatment of patients with metastatic melanoma, are being actively investigated for the treatment of multiple cancers including lung, breast, bladder and renal cancers along with other anti-PD-1/L1 antibodies. Early results of combining of anti-CTLA-4 antibody and anti-PD-1 antibody treatment for advanced melanoma patients are showing impressive response rates with manageable toxicity profiles. There are several other checkpoint molecules that are likely potential inhibitory targets. The outcome of blocking some of these negative immune regulators, such as LAG-3 or TIM-3, is being pursued in the clinic or about to enter clinical development. Blockade of these molecules is demonstrating promising preclinical activity alone or when combined with anti-PD-1/L1. Future studies will define bio-markers of these therapies and how to target them alone or in combination with other immunotherapies, chemotherapy, radiotherapy and small molecule inhibitors.

Addresses

¹ Department of Medicine, Division of Hematology-Oncology, University of California Los Angeles (UCLA), Los Angeles, CA, USA

² Department of Molecular and Medical Pharmacology, UCLA, Los Angeles, CA, USA

³ Department of Surgery, Division of Surgical-Oncology, UCLA, Los Angeles, CA, USA

⁴ Jonsson Comprehensive Cancer Center at UCLA, 10833 Le Conte Avenue, Los Angeles, CA 90095-1782, USA

⁵ Department of Molecular, Cellular and Integrative Physiology, UCLA, Los Angeles, CA, USA

Corresponding author: Ribas, Antoni (aribas@mednet.ucla.edu)

Current Opinion in Immunology 2015, **33**:23–35

This review comes from a themed issue on **Tumour immunology**

Edited by **Hans Schreiber** and **Philip D Greenberg**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 23rd January 2015

<http://dx.doi.org/10.1016/j.coi.2015.01.006>

0952-7915/© 2015 Published by Elsevier Ltd.

Introduction

Our immune system has the capability to identify and eliminate cells expressing tumor antigens [1*,2]. At the same time, it has developed sophisticated ways of

controlling its activity to maintain tolerance to self-antigens [3]. Various diseases arise from defects in this system such as autoimmune diseases [4], and cancer cells can exploit this system escaping from immune attack (immune evasion), which has been a major limitation of cancer immunotherapy [5–9]. Efforts made over past decades have allowed for our current understanding about immune escape mechanisms, one of which is the manipulation of immune cell-intrinsic checkpoints that are induced upon T cell activation [10]. First evidence of improved survival with blockade of these checkpoint molecules came from patients with advanced melanoma treated with the anti-CTLA-4 antibody ipilimumab, which demonstrated 10–15% of durable response [11**]. Blockade of the PD-1/PD-L1 pathway demonstrated unprecedented durable response in 30–35% of patients with advanced melanoma [12], which led to the FDA approval of pembrolizumab (anti-PD-1 antibody) in September 2014, and nivolumab in December 2014. This class of immune modulatory antibodies blocking PD-1 or PD-L1 are arguably the most exciting development in current cancer drug development (Figure 1).


A great deal of research has uncovered the complex interplay of immune regulatory molecules (co-inhibitory and co-stimulatory) that are regulating T cell activity [13**,14–16], and as CTLA-4 and PD-1 inhibitors have taught us, we can actually manipulate these molecules to activate our immune system to fight against cancers. However, many other studies will be required to understand how we can improve anti-tumor activity by combining checkpoint inhibitors or co-stimulatory activators, and possibly combining with conventional oncological therapies. Some of the new checkpoint inhibitors, such as LAG-3 and TIM-3, are close to clinical development, especially combined with PD-1 inhibitors [17–19]. There are many other potential candidates for further development. We currently do not have validated bio-markers to predict the likelihood of response for each patient for these checkpoint inhibitors. However, it is anticipated that in the near future, we will have tools to predict the chance of response of each patient before the initiation of these therapies.

Blockade of CTLA-4

Biology

Cytotoxic T lymphocyte-associated antigen-4 (CTLA-4, CD152) was initially discovered through cDNA library screening as one of the members of the immunoglobulin superfamily and found to be a co-inhibitory molecule expressed on T cells evidenced by profound autoimmune

Figure 1



Schematic representation of inhibitory immune checkpoints between antigen presenting cell (APC), T cell, cancer cell and normal cell. Upon T cell activation via T cell receptor (TCR) engagement by MHC-peptide complex between APC, inhibitory checkpoints are interacting with their cognate ligands expressed on APC. When T cell encounters antigen presented by MHC class I molecule on cancer cell, cancer cell expresses cognate ligands to interact with inhibitory checkpoints expressed on T cell. Inhibitory immune checkpoints also play a role in self-tolerance when T cell encounters normal cell. Most of inhibitory immune checkpoint molecules are members of immunoglobulin superfamily (IgSF), except HVEM, which is a member of TNF (Tumor necrosis factor) receptor superfamily. Molecules cited; CTLA-4 (Cytotoxic T lymphocyte-associated antigen-4, CD152); PD-1/L-1 (Programmed cell death-1 or CD279/Programmed cell death ligand-1 or CD274); TIM-3 (T-cell immunoglobulin and mucin containing protein-3); LAG-3 (Lymphocyte activation gene-3 or CD223); B7-H3 (B7 homolog 3, CD276); B7-H4 (B7 homolog 4, B7S1, B7x, VTCN1); VISTA (V-domain Ig suppressor of T cell activation, B7-H5 or PD-1Homolog, differentiation of embryonic stem cells 1; Dies 1); CEACAM1 (Carcinoembryonic antigen-related cell adhesion molecule 1, CD66a); BTLA (B and T lymphocyte attenuator, CD272); CD200R (CD200 receptor), CD200 (OX2); TIGIT (T cell Ig and ITIM doman, Vsig 9, Vstm3 or WUCAM).

phenomenon in knockout mouse models [20–23]. It has approximately 30% of sequence homology with CD28, which is a critical co-stimulatory molecule constitutively expressed on T cells. CTLA-4 is not expressed by naïve CD4⁺ or CD8⁺ T cells; however, it is contained intracellularly and TCR engagement induces trafficking to the cell surface where it can become engaged in signaling [24]. CTLA-4 binds to B7-1 (CD80) and B7-2 (CD86) as CD28 does, but with much higher affinity (10–40-fold) so that it arrests CD28 induced T cell activation by competing for ligand engagement, resulting in decreased cytokine production and cell cycle arrest [25,26]. It is also reported that CTLA-4 is constitutively expressed by Tregs, which play an important role in peripheral tolerance via suppressive activity on cytotoxic T cells [27–29].

Preclinical data

Based on *in vitro* studies suggesting that CTLA-4 is a key checkpoint molecule in adaptive immune responses, its

antitumor activity was first tested in various animal tumor models, such as breast [30], prostate [31], lymphoma [32], sarcoma [33], colon [34] and melanoma [35] via anti-CTLA-4 blocking antibody therapy. The first preclinical report was published in 1996 by Allison and co-workers, a study that demonstrated that anti-CTLA-4 antibody effectively enhances immune response to reject tumors [36]. The efficacy was restricted to a few cancer cell lines, and the B16 melanoma model was less sensitive and therapeutic efficacy was only achieved when it was combined with a GM-CSF vaccine [37,38]. These animal studies provided the evidence that the CTLA-4 blockade could result in significant anti-tumor activity by enhancing naturally or vaccine induced T cells, which led to testing it in a clinical setting.

Clinical data

Two anti-CTLA-4 fully human monoclonal antibodies, ipilimumab (IgG1 isotype) [39] and tremelimumab (IgG2

isotype) [40] reached clinical trials, mostly tested in patients with advanced melanoma. A pivotal phase III randomized trial was published for ipilimumab in 2010 [11^{••}]. This trial demonstrated significantly improved overall survival for patients with previously treated unresectable stage III or stage IV melanoma compared to a peptide vaccine. A subsequent phase III randomized trial for treatment-naïve patients with advanced melanoma showed that ipilimumab plus dacarbazine improved overall survival compared with treatment with dacarbazine alone [41^{••}]. These studies led to the FDA approval of ipilimumab for metastatic melanoma patients in 2011. Tremelimumab entered phase III trial for previously untreated patients with promising early phase I and II study results [42,43]. However, phase III data did not show survival benefit. There are several potential reasons for the difference between ipilimumab and tremelimumab: different dosing regimens (every 3 weeks for ipilimumab, every 3 months for tremelimumab), different exclusion criteria (tremelimumab trial excluded patients with LDH > 2X ULN) as well as unintended crossover patients in the control chemotherapy arm to ipilimumab which was easily accessible during the tremelimumab trial [44]. Currently, tremelimumab continues to be evaluated as a treatment in combination with other anticancer agents for melanoma and other tumor types [45,46].

Blockade of PD-1/PD-L1 pathway

Biology

Programmed cell death-1 (PD-1, CD279) is a type I transmembrane receptor member of the immunoglobulin superfamily, expressed by activated T cells, and binds to two ligands, PD-L1 (B7-H1, CD274) and PD-L2 (B7-DC, CD273), both of which are part of the B7 immunoglobulin superfamily [47]. PD-L1 is expressed on multiple normal tissues and malignant cells whereas PD-L2 is

mainly expressed by antigen presenting cells [48,49]. The critical role of PD-1 in immune regulatory function has been demonstrated by inhibition of the effector phase of T cells primarily within the tumor microenvironment as well as mouse strain-specific development of autoimmune disease with long latency in a PD-1 knockout mouse model [50–54]. Given the selective immune suppressive signals delivered by cancer, it was predicted that the blockade of PD-1/PD-L1 pathway will have greater antitumor activity and fewer side effects compared to CTLA-4 blockade [13^{••},48].

Preclinical/clinical data

Blockade of PD-1 has been tested in animal models such as murine pancreatic carcinoma [55], B16 melanoma [56], squamous cell carcinoma [57] and CT26 colon carcinoma [58,59], which demonstrated effective anti-tumor T cell responses. A PD-L1 blocking antibody was tested in 4T1 breast carcinoma model which overcame the resistance to immune mediated tumor rejection when treated together with a 4-1BB agonist antibody [60]. These studies confirmed the therapeutic potential of targeting this immune checkpoint and multiple anti-PD-1/PD-L1 antibodies have been evaluated in clinical trials as Table 1 shows. These studies have demonstrated an unprecedented 30–35% durable response in patients with advanced melanoma, which led to a first FDA approval for pembrolizumab (anti-PD-1 antibody) in September of 2014 [12,61^{••},62,63,64[•],65], followed by nivolumab in December 2014. Anti-PD-1/L1 blocking antibodies are currently being evaluated in other malignancies, such as lung [66], breast [67–69], bladder [70,71,72^{••}], renal cancers [73,74^{••}] and chemotherapy-refractory Hodgkin disease [75^{••}] with promising results. Arguably, this is one of the most exciting developments in cancer therapeutics and has illuminated the path to a new era in treatment of patients with cancer [76].

Table 1

Checkpoint blockade agents in clinical development

Target	Agent	Class
PD-1	Nivolumab (MDX1106, BMS936558, BMS-ONO)	IgG4 fully human antibody
	Pembrolizumab (MK3475M, lambrolizumab, Merck) ^a	IgG4 engineered humanized antibody
	Pidilizumab (CT-011, Cure Tech)	IgG1 humanized antibody
	AMP-224 (Amplimmune-GSK)	Fc-PD-L2 fusion protein
	AMP-514 (MED10680, Amplimmune) ^b	Monoclonal antibody
	AUNP-12 (Pierre Fabre, Aurigene)	Branched 29-amino acid peptide
PD-L1	BMS935559 (MDX-1105, BMS-ONO)	IgG4 fully human antibody
	BMS936559 (BMY)	IgG4 fully human antibody
	MPDL3280A (Genentech)	IgG1 engineered humanized antibody
	MED14736 (MedImmune, AZ)	IgG1 engineered humanized antibody
	MSB0010718C (Merck-Serono)	IgG1 antibody
CTLA-4	Tremelimumab (ticilimumab, CP-675,206, AZ)	IgG2 fully human antibody
LAG-3	BMS-986016	Monoclonal antibody
B7-H3	MGA271 (Macrogenics)	IgG1k humanized antibody

^a FDA approved for patients with advanced melanoma.

^b NCT02013804. Phase I (clinicaltrials.gov).

Biomarker/patient selection

With the success of the PD-1/PD-L1 blockade, it has become a top priority to identify and characterize the factors in the tumor microenvironment that predict which patients are likely to respond to this therapy. Tumor PD-L1 expression has been pursued as a potential biomarker and studies have demonstrated that PD-L1 baseline expression level showed a strong association with response to anti-PD-1 therapy. However, clinical responses were observed also in patients considered to be negative for PD-L1 expression in the tumor [77]. Therefore, we need to integrate it into a more complex equation. In order for PD-1/PD-L1 blockade therapy to work, tumor antigen specific pre-existing T cells that become disabled by tumor PD-L1 expression are likely required [72^{••}]. PD-L1 expression in the absence of T cells in a tumor microenvironment (constitutive expression by oncogenic process) as well as the presence of T cells without an adaptive PD-L1 expression (inducible PD-L1 expression upon cytokine stimulation mainly by interferons from immune cells) [13^{••},78–81] is of unknown significance. Many endeavors are taking place to develop highly sensitive and specific assays to detect and quantify PD-L1 expression and yet PD-L1 expression alone will not likely capture the full biology of PD-1/PD-L1 interaction. The presence of cytotoxic CD8⁺ T cells is a standard IHC protocol. However, it does not provide tumor specificity. An indirect way of measuring tumor specific T cells is the expression of PD-1, PD-L1, CD137, OX40 or TIM3, which are TCR activation markers [74^{••},82^{••}]. Clonality of TCR can be analyzed by deep sequencing [83] and specific TCR engagement with tumor antigen led to release of cytokines, especially interferons which can be detected by laser capture microscopy, PCR or in situ hybridization methods [84^{••}]. These efforts to characterize each variable in this complex equation will inform oncologists of a patient's predisposition to respond to anti-PD-1/L1 antibody therapy [85].

Blockade of TIM-3, LAG-3, B7-H3, B7-H4, VISTA, CEACAM1, BTLA, CD200-CD200R and TGIT

TIM-3

T-cell immunoglobulin and mucin containing protein-3 (TIM-3) was discovered a decade ago as a molecule expressed on IFN- γ producing CD4⁺ T helper 1 (Th1) and CD8⁺ T cytotoxic 1 (Tc1) cells [85]. The functional significance of TIM-3 was studied in the setting of autoimmunity initially [85–87], and its function as a T cell inhibitory receptor became clear as galectin-9 (C-type lectin) was identified for its ligand, which is mainly expressed on Treg [88]. TIM-3 is believed to play a critical role in inhibiting Th1 responses through Treg expressing galectin-9, evidenced by TIM-3 blockade administration at the time of allogeneic transplantation which resulted in loss of tolerogenic effects of donor-derived Treg cell transfusion [87]. Its critical role in

tumor-induced immune suppression at the immune checkpoint came from a study showing that TIM-3 marks most suppressed or dysfunctional populations of CD8⁺ T cells in animal models of solid and hematologic malignancies [89[•],90–92]. TIM-3 blockade in these animal models has successfully demonstrated similar anti-tumor activity compared to PD-1 pathway blockade [93], with greater efficacy with the combined use of TIM-3 and PD-1 blockade [19]. Increasing data support the relevance of TIM-3 blockade in human cancer as TIM-3 positive NY-ESO-1 specific CD8⁺ T cells in patients with melanoma showed dysfunctional phenotypes and TIM-3 blockade restored IFN- γ and TNF- α production as well as the proliferation of NY-ESO-1-specific CD8⁺ T cells in response to antigenic stimulation [89[•]]. TIM-3 has now emerged as an immune checkpoint receptor with its selective expression in tumor tissue as well as its critical role in multiple immune suppressive mechanisms, which strongly supports TIM-3 targeted immunotherapies as single or combined modalities. Monoclonal antibodies blocking TIM-3 are under preclinical development by several companies; however detailed information about these is not yet available.

LAG-3

Lymphocyte activation gene-3 (LAG-3, CD223) is another surface molecule of the immunoglobulin superfamily, expressed on activated T cells, NK cells, B cells and plasmacytoid dendritic cells, that plays an important role in negative regulation of T cell proliferation via binding to MHC class II with high affinity [94–96]. LAG-3 is also required for optimal function of Treg. Its co-expression with PD-1 in tumor infiltrating lymphocytes in certain malignancies correlates with impaired CD8⁺ effector T cell function [97,98]. LAG-3-Ig (Immutep, IMP321, Paris), a fusion protein (antigen presenting cell activator), has been tested in animal models as well as in human clinical trials. Early trial with standard influenza vaccine or hepatitis B vaccine showed high tolerability and Th1 CD4 T cell response was observed in several participants [99,100]. This led to a clinical trial in patients with renal cell carcinoma as a single agent, dose escalation trial [101]. This agent was well tolerated and it appeared to be correlated to development of effector phenotype of CD8⁺ T cells. There were no objective responses, although several patients showed stabilization of disease. It was then tested in patients with metastatic breast cancer in combination with first line paclitaxel [102,103], which demonstrated an objective response rate of 50% compared to an historical response rate of 25%. Phase III of the clinical trial is expected to be launched in 2015. As eluded to above, previous studies have suggested that dual inhibition of LAG-3 and PD-1 could enhance T effector activity and antitumor activity [17,104] and a clinical trial is actively recruiting patients to assess the safety and tolerability as well as dose limiting toxicity of BMS-986016 alone and in combination with nivolumab in

subjects with advanced solid malignancies (metastatic and/or unresectable; NCT01968109, clinicaltrials.gov). Based on the concurrent expression of both PD-1 and LAG-3 being largely limited to tumor infiltrating lymphocytes, it is expected to have less toxicity than CTLA-4, likely similar to anti-PD-1 antibody therapy.

B7-H3

B7-H3 (B7 homolog 3, CD276) is a type I transmembrane protein which shares 31% of sequence homology with B7-H1 [105,106]. Its transcript has a ubiquitous expression pattern, however, protein expression is limited. It is also found to be highly expressed in a variety of cancer types, including breast, pancreatic, prostate, ovarian and renal cell carcinoma [107–112]. The function of B7-H3 is controversial since both co-stimulatory and co-inhibitory properties have been described [105,113,114], and its expression in cancer is associated with either prolonged survival or poor prognosis. These studies indicate the complex role of B7-H3 in tumor immunity and it appears to be dependent on the contexts of tumor specific micro-environmental factors [115,116]. MGA271, a fully human monoclonal antibody, has been developed to target B7-H3, which demonstrated potent anti-tumor activity in xenograft models of renal cell and bladder carcinoma [117]. Clinical trials with MG271 for multiple refractory cancers that express B7-H3 are ongoing and the result will provide further insights on role of B7-H3 on tumor immunity.

B7-H4

B7-H4 (B7 homolog 4) is a member of the B7 family transmembrane protein identified in 2003, also known as B7S1, B7x and V-set domain containing T cell activation inhibitor (Vtcn1), which binds to unknown receptors on activated T cells and results in inhibition of effector function [118–120]. Its protein expression appeared to be limited to APC as well as cancer cells, although mRNA is detected in most nonhematopoietic tissues [116,121]. Studies have demonstrated the association of high expression of B7-H4 with poor prognosis in human cancers [122–128], inverse correlation with its expression by cancer cells as well as tumor associated macrophages (TAM) and tumor infiltrating lymphocytes, which had been tested in xenograft models by blocking B7-H4 expression [129,130]. Although it has not been yet tested in the clinical setting, B7-H4 may be an attractive target based on its limited expression on tumors as well as TAM and its important roles in immune evasion and tumorigenesis.

VISTA

V-domain Ig suppressor of T cell activation (VISTA, also known as B7-H5 or PD-1 Homolog and differentiation of embryonic stem cells 1; Dies 1) is a negative checkpoint regulator and shares homology with PD-L1 [131*,132,133]. It is mainly expressed on hematopoietic compartment and highly expressed on myeloid lineage,

especially APCs. VISTA induces forkhead box p3 (FOXP3) expression in T cells, and VISTA-expressing APCs or VISTA-Ig fusion protein inhibits T cell proliferation and *in vitro* cytokine production. [134]. In mouse models, its expression is positively correlated with CD11b, also found on tumor infiltrating lymphocytes, and the blockade of VISTA enhances antitumor immunity [135*]. Its expression on cancer cells enhances tumor-invasion by augmenting type I matrix metalloproteinase (MMP) [136,137] and it may offer therapeutic application for human cancer.

CEACAM1

Carcinoembryonic antigen-related cell adhesion molecule 1 (CEACAM1, CD66a) is a transmembrane glycoprotein that belongs to the carcinoembryonic antigen family and is also a member of the immunoglobulin superfamily [138–140]. It is composed of sequentially ordered immunoglobulin-like domains, expressed on a variety of cells, and has multiple biological functions, especially cell to cell adhesion [141]. It has been reported that CEACAM1 is an independent poor prognostic factor for patients with melanoma and it strongly predicted the development of metastatic disease [142]. CEACAM1 acts as an inhibitory molecule that negatively regulates cytotoxic T cell proliferation, which is activity evidenced by direct T cell receptor cross linking [143], and its homotypic interaction inhibits natural killer cell mediated killing as well [144,145]. Interestingly, a recent study by Huang *et al.* showed that the presence of CEACAM1 endows TIM-3 inhibitory function. CEACAM1 facilitates the maturation and surface expression of TIM-3 by forming a heterodimeric interaction, and co-blockade of CEACAM1 and TIM-3 enhanced the anti-tumor immune response [146**]. Monoclonal antibody targeting CEACAM1 was developed and showed that it efficiently blocks CEACAM1 homophilic interactions and renders melanoma cells susceptible to T cell mediated cytotoxicity both *in vitro* and *in vivo* [147]. Other than melanoma, several other malignancies, such as NSCLC have a prognostic association between CEACAM1 expression. Therefore, CEACAM1 targeted immunotherapy will be developed clinically soon.

BTLA

B and T lymphocyte attenuator (BTLA, CD272) is a type I membrane glycoprotein identified as an inhibitory receptor, which has structural similarities with CTLA-4 and PD-1 and is mainly expressed by immune cells [148,149]. Contrary to other B7 family members (immunoglobulin superfamily), its ligand, Herpesvirus entry mediator (HVEM) is a member of TNF superfamily [150,151]. HVEM has additional partners in addition to BTLA, such as CD160, LT α and LIGHT. Depending on its partner, CD160, BTLA, LIGHT/HVEM pathway delivers bidirectional signaling [152–154]. BTLA over-expression has been reported in hematological malignancies as well as

melanoma and it appears to be associated with impaired tumor-specific T cell activity, especially associated with PD-1 expression [155–158]. In one study, coupled with PD-1 and TIM-3 blockade, BTLA inhibition enhanced the expansion, proliferation and cytokine production of tumor specific CD8⁺ T cells [159] and this suggests that these three checkpoint molecules play an important role in immune evasion in patients with melanoma.

CD200–CD200R

CD200R is a type I membrane glycoprotein, which belongs to the immunoglobulin superfamily, expressed on myeloid cell as well as lymphoid cells, such as NK cells, CD4⁺ and CD8⁺ T cells [160,161]. In contrast to CD200R, CD200 (OX2) is widely expressed on immune and non-immune cells [162], CD200–CD200R pathway has an important role in regulating the strength of the initial response to infectious challenge in order to maintain homeostasis [163,164] and the role of anti-cancer immunity emerged from these studies have shown poor prognostic implication of CD200 expression on multiple myeloma, acute myeloid leukemia, B cell chronic leukemia, ovarian cancer and melanoma [165–167]. Studies have revealed that suppression of anti-tumor activity comes from direct CD200–CD200R inhibitory signaling on NK and myeloid cells [168] and indirect effect on T cells via DCs or macrophages [168]. CD200 expression on non-malignant cells appears to play a significant role as evidenced by CD200 transgenic mice, with larger primary tumor formation as well as lymph node metastases [169]. This implies a systemic effect of CD200–CD200R pathway on anti-tumor immunity and it can be targeted even with CD200 negative tumors. A monoclonal antibody blocking CD200 (ALXN6000: samalizumab) has been tested in patients with B-CLL and multiple myeloma (Clinical trials.gov Identifier: NCT00648739). The study showed 36% (8/22) of evaluable patients demonstrated 10% reduction in bulky disease. One patient had confirmed partial response (71% reduction in bulky disease). One thing to pay attention to block this pathway is that CD200–CD200R is an important regulator of inflammation that may promote cancer growth [170]. Increased inflammation from CD200–CD200R blockade, therefore, may result in unwanted effect of tumor growth or progression [171]. Thus, it would be important to find the fine balance in order to block this pathway for further clinical development [172*].

TIGIT

TIGIT (T cell Ig and ITIM domain, also called Vsig 9, Vstm3 or WUCAM) is a CD28 family receptor protein expressed on T cells and NK cells upon activation, binds to two ligands, CD155 (PVR: poliovirus receptor) and CD112 (PVRL2, nectin-2) on antigen presenting cells that leads to tolerogenic phenotype [173,174]. It shares ligands with CD226 (DNAM-1) and TIGIT/CD226 forms a network that regulates human T cell function

as CD28/CTLA-4–CD80/CD86 pathway does. CD226 delivers positive signal, and TIGIT delivers inhibitory signal [175,176]. This pathway appears to be involved in multiple autoimmune diseases, such as type I diabetes and multiple sclerosis [177,178]. Transgenic mouse studies have shown that TIGIT deficiency per se does not cause spontaneous autoimmunity. However, these mice do have augmented immunity upon immunization [179]. Inhibitory role in CNS autoimmunity was observed in a series of studies which showed increased experimental autoimmune encephalitis (EAE) severity upon TIGIT blockade [180]. These studies suggest that TIGIT plays a role in regulating peripheral immune tolerance and future studies will define its role in anti-tumor activity as well as potential therapeutic application.

Combined checkpoint blockade

Preclinical studies combining CTLA-4 and PD-1 blocking antibodies demonstrated superior antitumor activity [70,181**,182] with similar toxicity profile as anti-CTLA-4 monotherapy. The first phase I clinical trial combining ipilimumab and nivolumab [183**] was updated in 2014 ASCO annual meeting [184] with an impressive 2 year survival of 79% (objective response rate of 43%) among patients with advanced melanoma, although it did show increased toxicity compared to either agents alone (62% of grade 3/4 toxicities). A phase III study has completed the accrual, and this combinatorial treatment is being tested in other cancers. This combination strategy was also studied with a tumor vaccine, which showed effective restoration of T cell rejection function in animal models [185].

Combination treatment with checkpoint blockade and small molecule inhibitors is an attractive option given the potential for increased tumor antigen presentation, a larger window period for immunotherapy and the feasibility to deliver full doses of each agent due to different mechanisms of action [186,187*,188–190]. Combination treatment of ipilimumab and vemurafenib was studied for patients with advanced melanoma; however, the trial was terminated prematurely due to significant hepatotoxicity [191]. The etiology of hepatotoxicity is not clear, but likely related to paradoxical activation of the MAPK pathway. Early clinical studies has been launched with anti-PD-1 or anti-PD-L1 antibodies combined with BRAF plus MEK inhibitors for patients with BRAF mutated melanoma given the potential benefit of mitigating paradoxical activation of MAPK pathway although there was a concern for immune cell inhibition [192]. These studies will address the feasibility of a triple combination, toxicity profile as well as the effect on immune cell that is highly relevant to anti-tumor activity. Another attractive strategy is to combine checkpoint blockade with B cell receptor signaling inhibitors (ibrutinib and idelalisib), which were approved for B cell lymphoid malignancies. Recent studies demonstrated

the effects of these agents on T cell activities, especially on Tregs [193^{••},194,195^{••},196] and future studies will need to define clinical activity of this combinatorial treatment.

Other checkpoint blockade combinations are in the early clinical developmental stage, combining anti-PD-1 with antibodies blocking LAG-3 or TIM-3. LAG-3 and TIM-3 are being considered as T cell exhaustion markers and can be targeted with PD-1/L1 pathway blockade supported by pre-clinical models demonstrated promising anti-tumor activity [17,19,92,97]. Phase I study comparing LAG-3 vs. LAG-3 plus PD-1 blockade among patients with advanced solid tumors has just launched (NCT01968109) and future studies will tell us whether these combinations are feasible with manageable toxicity and improved responses compared to anti-PD-1/L1 monotherapy.

Future direction and concluding remarks

It is exciting time to witness how immunotherapy, specifically treatment with checkpoint blockade monoclonal antibodies, is becoming one of the main armamentarium for cancer therapy. Unprecedented durable response among patients with advanced melanoma, Hodgkin disease, renal cell carcinoma, lung and bladder cancers, among others, with anti-PD-1 or anti-PD-L1 antibody monotherapy has set the stage to revolutionize treatment approaches for patients with advanced cancer. However, a lot of further work needs to be done in order to define biomarkers to predict the response to anti-PD-1 therapies and identify ideal combinations among checkpoint blockade as well as small molecule inhibitors, chemotherapy and radiotherapy through rational design of the study based on scientific evidence. There are several emerging checkpoint molecules that can be targeted. We have mainly discussed checkpoint blockade, however, immune receptor stimulators are also in active investigation along with others in pre-clinical development, such as ICOS (CD278) agonistic antibody which is being evaluated in combination with ipilimumab. Another promising strategy is the combination of agonistic antibodies targeting CD137 (41BB) (BMS-66513 or PF-05082566) have been tested in phase I and II studies for patients with solid tumors, including melanoma, renal carcinoma, and ovarian cancer. The utilization of these agents to modulate our immune system to against cancer is eagerly anticipated, in particular in combination with checkpoint blockade therapy.

Conflict of interest

A.R. has served as a compensated consultant for Amgen, Compugen, Daiichi-Sankyo, Genentech-Roche, GSK, Merck and Novartis with the honoraria paid to the institution.

Acknowledgements

This work was funded by the National Institutes of Health (NIH) grants P01 CA168585, 2U54 CA151819, R01CA170689 and P01 CA132681, Stand

Up to Cancer-Cancer Research Institute (SU2C-CRI) grant SUC2CR-AACR-DT10, the Melanoma Research Foundation, the Dr. Robert Vigen Memorial Fund, the Ressler Family Foundation and the Wesley Coyle Memorial Fund (to A.R.), Oncology (5T32CA009297-30) and Dermatology (5T32AR058921-05) training grants.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Vesely MD, Kershaw MH, Schreiber RD, Smyth MJ: **Natural innate and adaptive immunity to cancer.** *Annu Rev Immunol* 2011, **29**:235-271.
- Excellent review on the current understanding of concept of the cancer immunoediting.
2. Burnet FM: **The concept of immunological surveillance.** *Prog Exp Tumor Res* 1970, **13**:1-27.
3. Dunn GP, Old LJ, Schreiber RD: **The three Es of cancer immunoediting.** *Annu Rev Immunol* 2004, **22**:329-360.
4. Goodnow CC: **Multistep pathogenesis of autoimmune disease.** *Cell* 2007, **130**:25-35.
5. Zitvogel L, Tesniere A, Kroemer G: **Cancer despite immunosurveillance: immunoselection and immunosubversion.** *Nat Rev Immunol* 2006, **6**:715-727.
6. Rabinovich GA, Gabrilovich D, Sotomayor EM: **Immunosuppressive strategies that are mediated by tumor cells.** *Annu Rev Immunol* 2007, **25**:267-296.
7. Finn OJ: **Immuno-oncology: understanding the function and dysfunction of the immune system in cancer.** *Ann Oncol* 2012, **23**(Suppl. 8):viii6-viii9.
8. Zou W: **Immunosuppressive networks in the tumour environment and their therapeutic relevance.** *Nat Rev Cancer* 2005, **5**:263-274.
9. Schwartz RH: **T cell anergy.** *Annu Rev Immunol* 2003, **21**:305-334.
10. Callahan MK, Wolchok JD: **At the bedside: CTLA-4- and PD-1-blocking antibodies in cancer immunotherapy.** *J Leukoc Biol* 2013, **94**:41-53.
11. Hodi FS, O'Day SJ, McDermott DF, Weber RW, Sosman JA, Haanen JB, Gonzalez R, Robert C, Schadendorf D, Hassel JC *et al.*: **Improved survival with ipilimumab in patients with metastatic melanoma.** *N Engl J Med* 2010, **363**:711-723.
- Pivotal phase III trial comparing ipilimumab with or without gp100 vs. gp100 alone. Trial demonstrated improved overall survival in patients with previously treated metastatic melanoma.
12. Hamid O, Robert C, Daud A, Hodi FS, Hwu WJ, Kefford R, Wolchok JD, Hersey P, Joseph RW, Weber JS *et al.*: **Safety and tumor responses with lambrolizumab (anti-PD-1) in melanoma.** *N Engl J Med* 2013, **369**:134-144.
13. Pardoll DM: **The blockade of immune checkpoints in cancer immunotherapy.** *Nat Rev Cancer* 2012, **12**:252-264.
- Outstanding review of current understanding and clinical development of immune checkpoint blockade.
14. Butt AQ, Mills KH: **Immunosuppressive networks and checkpoints controlling antitumor immunity and their blockade in the development of cancer immunotherapeutics and vaccines.** *Oncogene* 2014, **33**:4623-4631.
15. Nirschl CJ, Drake CG: **Molecular pathways: coexpression of immune checkpoint molecules: signaling pathways and implications for cancer immunotherapy.** *Clin Cancer Res* 2013, **19**:4917-4924.
16. Weber J: **Immune checkpoint proteins: a new therapeutic paradigm for cancer — preclinical background: CTLA-4 and PD-1 blockade.** *Semin Oncol* 2010, **37**:430-439.
17. Woo SR, Turnis ME, Goldberg MV, Bankoti J, Selby M, Nirschl CJ, Bettini ML, Gravano DM, Vogel P, Liu CL *et al.*: **Immune inhibitory molecules LAG-3 and PD-1 synergistically regulate T-cell**

- function to promote tumoral immune escape. *Cancer Res* 2012, **72**:917-927.
18. Turnis ME, Korman AJ, Drake CG, Vignali DA: **Combinatorial immunotherapy: PD-1 may not be LAG-ing behind any more.** *Oncoimmunology* 2012, **1**:1172-1174.
 19. Sakuishi K, Apetoh L, Sullivan JM, Blazar BR, Kuchroo VK, Anderson AC: **Targeting Tim-3 and PD-1 pathways to reverse T cell exhaustion and restore anti-tumor immunity.** *J Exp Med* 2010, **207**:2187-2194.
 20. Linsley PS, Brady W, Grosmaire L, Aruffo A, Damle NK, Ledbetter JA: **Binding of the B cell activation antigen B7 to CD28 costimulates T cell proliferation and interleukin 2 mRNA accumulation.** *J Exp Med* 1991, **173**:721-730.
 21. Brunet JF, Denizot F, Luciani MF, Roux-Dosseto M, Suzan M, Mattei MG, Golstein P: **A new member of the immunoglobulin superfamily — CTLA-4.** *Nature* 1987, **328**:267-270.
 22. Waterhouse P, Penninger JM, Timms E, Wakeham A, Shahinian A, Lee KP, Thompson CB, Griesser H, Mak TW: **Lymphoproliferative disorders with early lethality in mice deficient in Ctlα-4.** *Science* 1995, **270**:985-988.
 23. Tivol EA, Borriello F, Schweitzer AN, Lynch WP, Bluestone JA, Sharpe AH: **Loss of CTLA-4 leads to massive lymphoproliferation and fatal multiorgan tissue destruction, revealing a critical negative regulatory role of CTLA-4.** *Immunity* 1995, **3**:541-547.
 24. Walunas TL, Lenschow DJ, Bakker CY, Linsley PS, Freeman GJ, Green JM, Thompson CB, Bluestone JA: **CTLA-4 can function as a negative regulator of T cell activation.** *Immunity* 1994, **1**:405-413.
 25. Chen L: **Co-inhibitory molecules of the B7-CD28 family in the control of T-cell immunity.** *Nat Rev Immunol* 2004, **4**:336-347.
 26. Pentcheva-Hoang T, Egen JG, Wojnoonski K, Allison JP: **B7-1 and B7-2 selectively recruit CTLA-4 and CD28 to the immunological synapse.** *Immunity* 2004, **21**:401-413.
 27. Wing K, Onishi Y, Prieto-Martin P, Yamaguchi T, Miyara M, Fehervari Z, Nomura T, Sakaguchi S: **CTLA-4 control over Foxp3⁺ regulatory T cell function.** *Science* 2008, **322**:271-275.
 28. Shevach EM: **Mechanisms of foxp3⁺ T regulatory cell-mediated suppression.** *Immunity* 2009, **30**:636-645.
 29. Sakaguchi S, Wing K, Onishi Y, Prieto-Martin P, Yamaguchi T: **Regulatory T cells: how do they suppress immune responses?** *Int Immunol* 2009, **21**:1105-1111.
 30. Hurwitz AA, Yu TF, Leach DR, Allison JP: **CTLA-4 blockade synergizes with tumor-derived granulocyte-macrophage colony-stimulating factor for treatment of an experimental mammary carcinoma.** *Proc Natl Acad Sci U S A* 1998, **95**:10067-10071.
 31. Kwon ED, Hurwitz AA, Foster BA, Madias C, Feldhaus AL, Greenberg NM, Burg MB, Allison JP: **Manipulation of T cell costimulatory and inhibitory signals for immunotherapy of prostate cancer.** *Proc Natl Acad Sci U S A* 1997, **94**:8099-8103.
 32. Van Ginderachter JA, Liu Y, Geldhof AB, Brijs L, Thielemans K, De BP, Raes G: **B7-1, IFN gamma and anti-CTLA-4 co-operate to prevent T-cell tolerization during immunotherapy against a murine T-lymphoma.** *Int J Cancer* 2000, **87**:539-547.
 33. Espenschied J, Lamont J, Longmate J, Pendas S, Wang Z, Diamond DJ, Ellenhorn JD: **CTLA-4 blockade enhances the therapeutic effect of an attenuated poxvirus vaccine targeting p53 in an established murine tumor model.** *J Immunol* 2003, **170**:3401-3407.
 34. Saha A, Chatterjee SK: **Combination of CTL-associated antigen-4 blockade and depletion of CD25 regulatory T cells enhance tumour immunity of dendritic cell-based vaccine in a mouse model of colon cancer.** *Scand J Immunol* 2010, **71**:70-82.
 35. Suttmüller RP, van Duivenvoorde LM, van EA, Schumacher TN, Wildenberg ME, Allison JP, Toes RE, Offringa R, Melief CJ: **Synergism of cytotoxic T lymphocyte-associated antigen 4 blockade and depletion of CD25(+) regulatory T cells in antitumor therapy reveals alternative pathways for suppression of autoreactive cytotoxic T lymphocyte responses.** *J Exp Med* 2001, **194**:823-832.
 36. Leach DR, Krummel MF, Allison JP: **Enhancement of antitumor immunity by CTLA-4 blockade.** *Science* 1996, **271**:1734-1736.
 37. van EA, Hurwitz AA, Allison JP: **Combination immunotherapy of B16 melanoma using anti-cytotoxic T lymphocyte-associated antigen 4 (CTLA-4) and granulocyte/macrophage colony-stimulating factor (GM-CSF)-producing vaccines induces rejection of subcutaneous and metastatic tumors accompanied by autoimmune depigmentation.** *J Exp Med* 1999, **190**:355-366.
 38. Hodi FS: **Cytotoxic T-lymphocyte-associated antigen-4.** *Clin Cancer Res* 2007, **13**:5238-5242.
 39. Boasberg P, Hamid O, O'Day S: **Ipilimumab: unleashing the power of the immune system through CTLA-4 blockade.** *Semin Oncol* 2010, **37**:440-449.
 40. Ribas A: **Clinical development of the anti-CTLA-4 antibody tremelimumab.** *Semin Oncol* 2010, **37**:450-454.
 41. Robert C, Thomas L, Bondarenko I, O'Day S, Weber J, Garbe C, Lebbe C, Baurain JF, Testori A, Grob JJ *et al.*: **Ipilimumab plus dacarbazine for previously untreated metastatic melanoma.** *N Engl J Med* 2011, **364**:2517-2526.
- Pivotal phase III trial comparing ipilimumab in combination with dacarbazine vs. dacarbazine plus placebo. Trial showed improved overall survival in patients with previously untreated metastatic melanoma. Following these studies together with Ref. [11**], ipilimumab was approved by FDA in 2011.
42. Camacho LH, Antonia S, Sosman J, Kirkwood JM, Gajewski TF, Redman B, Pavlov D, Bulanhagui C, Bozon VA, Gomez-Navarro J *et al.*: **Phase I/II trial of tremelimumab in patients with metastatic melanoma.** *J Clin Oncol* 2009, **27**:1075-1081.
 43. Kirkwood JM, Lorigan P, Hersey P, Hauschild A, Robert C, McDermott D, Marshall MA, Gomez-Navarro J *et al.*: **Phase II trial of tremelimumab (CP-675,206) in patients with advanced refractory or relapsed melanoma.** *Clin Cancer Res* 2010, **16**:1042-1048.
 44. Ribas A, Kefford R, Marshall MA, Punt CJ, Haanen JB, Marmol M, Garbe C, Gogas H, Schachter J, Linette G *et al.*: **Phase III randomized clinical trial comparing tremelimumab with standard-of-care chemotherapy in patients with advanced melanoma.** *J Clin Oncol* 2013, **31**:616-622.
 45. Tarhini AA, Cherian J, Moschos SJ, Tawbi HA, Shuai Y, Gooding WE, Sander C, Kirkwood JM: **Safety and efficacy of combination immunotherapy with interferon alfa-2b and tremelimumab in patients with stage IV melanoma.** *J Clin Oncol* 2012, **30**:322-328.
 46. McNeel DG, Smith HA, Eickhoff JC, Lang JM, Staab MJ, Wilding G, Liu G: **Phase I trial of tremelimumab in combination with short-term androgen deprivation in patients with PSA-recurrent prostate cancer.** *Cancer Immunol Immunother* 2012, **61**:1137-1147.
 47. Keir ME, Butte MJ, Freeman GJ, Sharpe AH: **PD-1 and its ligands in tolerance and immunity.** *Annu Rev Immunol* 2008, **26**:677-704.
 48. Okazaki T, Honjo T: **PD-1 and PD-1 ligands: from discovery to clinical application.** *Int Immunol* 2007, **19**:813-824.
 49. Butte MJ, Keir ME, Phamduy TB, Sharpe AH, Freeman GJ: **Programmed death-1 ligand 1 interacts specifically with the B7-1 costimulatory molecule to inhibit T cell responses.** *Immunity* 2007, **27**:111-122.
 50. Dong H, Strome SE, Salomao DR, Tamura H, Hirano F, Flies DB, Roche PC, Lu J, Zhu G, Tamada K *et al.*: **Tumor-associated B7-H1 promotes T-cell apoptosis: a potential mechanism of immune evasion.** *Nat Med* 2002, **8**:793-800.
 51. Zou W, Chen L: **Inhibitory B7-family molecules in the tumour microenvironment.** *Nat Rev Immunol* 2008, **8**:467-477.
 52. Seliger B, Marincola FM, Ferrone S, Abken H: **The complex role of B7 molecules in tumor immunology.** *Trends Mol Med* 2008, **14**:550-559.

53. Nishimura H, Okazaki T, Tanaka Y, Nakatani K, Hara M, Matsumori A, Sasayama S, Mizoguchi A, Hiai H, Minato N *et al.*: **Autoimmune dilated cardiomyopathy in PD-1 receptor-deficient mice.** *Science* 2001, **291**:319-322.
 54. Jacobs JF, Idema AJ, Bol KF, Nierkens S, Grauer OM, Wesseling P, Grotenhuis JA, Hoogerbrugge PM, de Vries IJ, Adema GJ: **Regulatory T cells and the PD-L1/PD-1 pathway mediate immune suppression in malignant human brain tumors.** *NeuroOncology* 2009, **11**:394-402.
 55. Okudaira K, Hokari R, Tsuzuki Y, Okada Y, Komoto S, Watanabe C, Kurihara C, Kawaguchi A, Nagao S, Azuma M *et al.*: **Blockade of B7-H1 or B7-DC induces an anti-tumor effect in a mouse pancreatic cancer model.** *Int J Oncol* 2009, **35**: 741-749.
 56. Blank C, Brown I, Peterson AC, Spiotto M, Iwai Y, Honjo T, Gajewski TF: **PD-L1/B7H-1 inhibits the effector phase of tumor rejection by T cell receptor (TCR) transgenic CD8⁺ T cells.** *Cancer Res* 2004, **64**:1140-1145.
 57. Strome SE, Dong H, Tamura H, Voss SG, Flies DB, Tamada K, Salomao D, Chevillat J, Hirano F, Lin W *et al.*: **B7-H1 blockade augments adoptive T-cell immunotherapy for squamous cell carcinoma.** *Cancer Res* 2003, **63**:6501-6505.
 58. Iwai Y, Terawaki S, Honjo T: **PD-1 blockade inhibits hematogenous spread of poorly immunogenic tumor cells by enhanced recruitment of effector T cells.** *Int Immunol* 2005, **17**:133-144.
 59. Zang X, Allison JP: **The B7 family and cancer therapy: costimulation and coinhibition.** *Clin Cancer Res* 2007, **13**:5271-5279.
 60. Hirano F, Kaneko K, Tamura H, Dong H, Wang S, Ichikawa M, Rietz C, Flies DB, Lau JS, Zhu G, Tamada K, Chen L: **Blockade of B7-H1 and PD-1 by monoclonal antibodies potentiates cancer therapeutic immunity.** *Cancer Res* 2005, **65**:1089-1096.
 61. Topalian SL, Hodi FS, Brahmer JR, Gettinger SN, Smith DC, McDermott DF, Powderly JD, Carvajal RD, Sosman JA, Atkins MB *et al.*: **Safety, activity, and immune correlates of anti-PD-1 antibody in cancer.** *N Engl J Med* 2012, **366**:2443-2454.
- This article reports on the first phase 1 clinical trial with an anti-PD-1 antibody with significant antitumor activity in patients with melanoma, renal cell carcinoma and impressively also with lung cancer. It also provided the first evidence of the correlation of PD-L1 expression and tumor responses.
62. Topalian SL, Sznol M, McDermott DF, Kluger HM, Carvajal RD, Sharfman WH, Brahmer JR, Lawrence DP, Atkins MB, Powderly JD *et al.*: **Survival, durable tumor remission, and long-term safety in patients with advanced melanoma receiving nivolumab.** *J Clin Oncol* 2014, **32**:1020-1030.
 63. Brahmer JR, Tykodi SS, Chow LQ, Hwu WJ, Topalian SL, Hwu P, Drake CG, Camacho LH, Kauh J, Odunsi K *et al.*: **Safety and activity of anti-PD-L1 antibody in patients with advanced cancer.** *N Engl J Med* 2012, **366**:2455-2465.
 64. Robert C, Ribas A, Wolchok JD, Hodi FS, Hamid O, Kefford R, Weber JS, Joshua AM, Hwu WJ, Gangadhar TC *et al.*: **Anti-programmed-death-receptor-1 treatment with pembrolizumab in ipilimumab-refractory advanced melanoma: a randomised dose-comparison cohort of a phase 1 trial.** *Lancet* 2014, **384**:1109-1117.
- This article reports on the antitumor activity of pembrolizumab in patients who had previously become refractory to ipilimumab and led to the first approval of an anti-PD-1 antibody in the US.
65. Ribas A, Hodi FS, Kefford R, Hamid O, Daud A, Wolchok JD, Hwu WJ, Gangadhar TC, Patnaik A, Joshua AM *et al.*: **Efficacy and safety of the anti-PD-1 monoclonal antibody MK-3475 in 411 patients (pts) with melanoma (MEL).** *2014 ASCO Annual Meeting*. 2014. LBA9000 (abstract).
 66. Harvey RD: **Immunologic and clinical effects of targeting PD-1 in lung cancer.** *Clin Pharmacol Ther* 2014, **96**:214-223.
 67. Stagg J, Allard B: **Immunotherapeutic approaches in triple-negative breast cancer: latest research and clinical prospects.** *Ther Adv Med Oncol* 2013, **5**:169-181.
 68. Schalper KA: **PD-L1 expression and tumor-infiltrating lymphocytes: Revisiting the antitumor immune response potential in breast cancer.** *Oncoimmunology* 2014, **3**:e29288.
 69. Segal NH, Antonia SJ, Brahmer JR, Maio M, Blake-Haskins A, Li X, Vasselli J, Ibrahim RA, Lutzky J, Khleif S: **Preliminary data from a multi-arm expansion study of MEDI4736, an anti-PD-L1 antibody.** *2014 ASCO Annual Meeting*. 2014. 3002 (abstract).
 70. Mangsbo SM, Sandin LC, Anger K, Korman AJ, Loskog A, Totterman TH: **Enhanced tumor eradication by combining CTLA-4 or PD-1 blockade with CpG therapy.** *J Immunother* 2010, **33**:225-235.
 71. Raval RR, Sharabi AB, Walker AJ, Drake CG, Sharma P: **Tumor immunology and cancer immunotherapy: summary of the 2013 SITC primer.** *J Immunother Cancer* 2014, **2**:14.
 72. Tumei PC, Harview CL, Yearley JH, Shintaku IP, Taylor EJ, Robert L, Chmielowski B, Spasic M, Henry G, Ciobanu V *et al.*: **PD-1 blockade induces responses by inhibiting adaptive immune resistance.** *Nature* 2014, **515**:568-571.
- This article studied tumor samples from PD-1 clinical trial which led to first FDA approval of anti-PD-1 antibody (pembrolizumab). It showed the pre-existing CD8⁺ T cell located at the invasive tumor margin was associated with expression of PD-1/PD-L1 immune inhibitory axis and suggested that it may predict tumor response.
73. Tykodi SS: **PD-1 as an emerging therapeutic target in renal cell carcinoma: current evidence.** *OncoTargets Ther* 2014, **7**:1349-1359.
 74. Herbst RS, Soria JC, Kowanetz M, Fine GD, Hamid O, Gordon MS, Sosman JA, McDermott DF, Powderly JD, Gettinger SN *et al.*: **Predictive correlates of response to the anti-PD-L1 antibody MPDL3280A in cancer patients.** *Nature* 2014, **515**:563-567.
- This study evaluated the safety, activity and biomarkers of PD-L1 inhibition using engineered humanized antibody MPDL3280A. Among multiple cancer types, responses were observed with tumors expressing high levels of PD-L1, especially when it was expressed by tumor-infiltrating immune cells. And the response was associated with Th1 gene expression pattern.
75. Ansell SM, Lesokhin AM, Borrello I, Halwani A, Scott EC, Gutierrez M, Schuster SJ, Millenson MM, Cattray D, Freeman GJ *et al.*: **PD-1 blockade with nivolumab in relapsed or refractory Hodgkin's lymphoma.** *N Engl J Med* 2014.
- This is a phase I study evaluating nivolumab, an anti-PD-1 antibody, for relapsed/refractory Hodgkin's lymphoma. It demonstrated high levels of PD-L1 and PD-L2 expression on Reed-Sternberg (RS) cells which showed nuclear positivity of phosphorylated STAT3 (indicative of active JAK-STAT pathway). PD-1 blockade on Hodgkin's lymphoma appears to be very promising based on presented data, further larger studies will define its clinical activity.
76. Ribas A: **Tumor immunotherapy directed at PD-1.** *N Engl J Med* 2012, **366**:2517-2519.
 77. Taube JM, Klein A, Brahmer JR, Xu H, Pan X, Kim JH, Chen L, Pardoll DM, Topalian SL, Anders RA: **Association of PD-1, PD-1 ligands, and other features of the tumor immune microenvironment with response to anti-PD-1 therapy.** *Clin Cancer Res* 2014, **20**:5064-5074.
 78. Marzec M, Zhang Q, Goradia A, Raghunath PN, Liu X, Paessler M, Wang HY, Wysocka M, Cheng M, Ruggeri BA *et al.*: **Oncogenic kinase NPM/ALK induces through STAT3 expression of immunosuppressive protein CD274 (PD-L1, B7-H1).** *Proc Natl Acad Sci U S A* 2008, **105**:20852-20857.
 79. Kim J, Myers AC, Chen L, Pardoll DM, Truong-Tran QA, Lane AP, McDyer JF, Fortuno L, Schleimer RP: **Constitutive and inducible expression of b7 family of ligands by human airway epithelial cells.** *Am J Respir Cell Mol Biol* 2005, **33**:280-289.
 80. Jiang X, Zhou J, Giobbie-Hurder A, Wargo J, Hodi FS: **The activation of MAPK in melanoma cells resistant to BRAF inhibition promotes PD-L1 expression that is reversible by MEK and PI3K inhibition.** *Clin Cancer Res* 2013, **19**:598-609.
 81. Atefi M, Avramis E, Lassen A, Wong DJ, Robert L, Foulad D, Cerniglia M, Titz B, Chodon T, Graeber TG *et al.*: **Effects of MAPK and PI3K pathways on PD-L1 expression in melanoma.** *Clin Cancer Res* 2014, **20**:3446-3457.

82. Gros A, Robbins PF, Yao X, Li YF, Turcotte S, Tran E, Wunderlich JR, Mixon A, Farid S, Dudley ME *et al.*: **PD-1 identifies the patient-specific CD8(+) tumor-reactive repertoire infiltrating human tumors.** *J Clin Invest* 2014, **124**:2246-2259.
- In this article, authors evaluated phenotypic traits of CD8⁺ TILs and TCR β chain clonotypic frequency in melanoma in order to identify patient-specific repertoires of tumor reactive lymphocytes. Study demonstrated that PD-1 expression on CD8⁺ TILs also accurately identifies the repertoires of clonally expanded tumor reactive lymphocytes which implies dual importance of PD-1 expression in tumor microenvironment.
83. Robins HS, Ericson NG, Guenthoer J, O'Brian KC, Tewari M, Drescher CW, Bielas JH: **Digital genomic quantification of tumor-infiltrating lymphocytes.** *Sci Transl Med* 2013, **5**:214ra169.
84. Taube JM, Anders RA, Young GD, Xu H, Sharma R, McMiller TL, Chen S, Klein AP, Pardoll DM, Topalian SL *et al.*: **Colocalization of inflammatory response with B7-h1 expression in human melanocytic lesions supports an adaptive resistance mechanism of immune escape.** *Sci Transl Med* 2012, **4**:127ra37.
- Provided the first evidence that PD-L1 (B7-H1) co-localizes with CD8 infiltrates in tumors. This observation led to the realization that the infiltrating T cells were likely the ones that triggered the expression of PD-L1, in what has been termed adaptive immune resistance.
85. Monney L, Sabatos CA, Gaglia JL, Ryu A, Waldner H, Chernova T, Manning S, Greenfield EA, Coyle AJ, Sobel RA *et al.*: **Th1-specific cell surface protein Tim-3 regulates macrophage activation and severity of an autoimmune disease.** *Nature* 2002, **415**:536-541.
86. Kuchroo VK, Umetsu DT, DeKruyff RH, Freeman GJ: **The TIM gene family: emerging roles in immunity and disease.** *Nat Rev Immunol* 2003, **3**:454-462.
87. Anderson AC, Anderson DE: **TIM-3 in autoimmunity.** *Curr Opin Immunol* 2006, **18**:665-669.
88. Zhu C, Anderson AC, Schubart A, Xiong H, Imitola J, Khoury SJ, Zheng XX, Strom TB, Kuchroo VK: **The Tim-3 ligand galectin-9 negatively regulates T helper type 1 immunity.** *Nat Immunol* 2005, **6**:1245-1252.
89. Fourcade J, Sun Z, Benallaoua M, Guillaume P, Luescher IF, Sander C, Kirkwood JM, Kuchroo V, Zarour HM: **Upregulation of Tim-3 and PD-1 expression is associated with tumor antigen-specific CD8⁺ T cell dysfunction in melanoma patients.** *J Exp Med* 2010, **207**:2175-2186.
- In this article, authors performed elegant experiments to evaluate the significance of co-expression of PD-1 as well as TIM-3 in tumor antigen specific CD8⁺ T cells. Study showed that co-expressed T cells displayed more dysfunctional phenotype which can be reversed with TIM-3 blockade. Moreover, combined with PD-1 blockade was synergistic to activate T cells. This study has strong translational relevance to target both checkpoint molecules in clinical trial.
90. Gao X, Zhu Y, Li G, Huang H, Zhang G, Wang F, Sun J, Yang Q, Zhang X, Lu B: **TIM-3 expression characterizes regulatory T cells in tumor tissues and is associated with lung cancer progression.** *PLoS ONE* 2012, **7**:e30676.
91. Yang ZZ, Grote DM, Ziesmer SC, Niki T, Hirashima M, Novak AJ, Witzig TE, Ansell SM: **IL-12 upregulates TIM-3 expression and induces T cell exhaustion in patients with follicular B cell non-Hodgkin lymphoma.** *J Clin Invest* 2012, **122**:1271-1282.
92. Zhou Q, Munger ME, Veenstra RG, Weigel BJ, Hirashima M, Munn DH, Murphy WJ, Azuma M, Anderson AC, Kuchroo VK *et al.*: **Coexpression of Tim-3 and PD-1 identifies a CD8⁺ T-cell exhaustion phenotype in mice with disseminated acute myelogenous leukemia.** *Blood* 2011, **117**:4501-4510.
93. Ngiew SF, von SB, Akiba H, Yagita H, Teng MW, Smyth MJ: **Anti-TIM3 antibody promotes T cell IFN-gamma-mediated antitumor immunity and suppresses established tumors.** *Cancer Res* 2011, **71**:3540-3551.
94. Huard B, Gaulard P, Faure F, Hercend T, Triebel F: **Cellular expression and tissue distribution of the human LAG-3-encoded protein, an MHC class II ligand.** *Immunogenetics* 1994, **39**:213-217.
95. Triebel F, Jitsukawa S, Baixeras E, Roman-Roman S, Genevee C, Viegas-Pequignot E, Hercend T: **LAG-3, a novel lymphocyte activation gene closely related to CD4.** *J Exp Med* 1990, **171**:1393-1405.
96. Kisielow M, Kisielow J, Capoferri-Sollami G, Karjalainen K: **Expression of lymphocyte activation gene 3 (LAG-3) on B cells is induced by T cells.** *Eur J Immunol* 2005, **35**:2081-2088.
97. Matsuzaki J, Gnjatich S, Mhawech-Fauceglia P, Beck A, Miller A, Tsuji T, Eppolito C, Qian F, Lele S, Shrikant P *et al.*: **Tumor-infiltrating NY-ESO-1-specific CD8⁺ T cells are negatively regulated by LAG-3 and PD-1 in human ovarian cancer.** *Proc Natl Acad Sci U S A* 2010, **107**:7875-7880.
98. Grosso JF, Goldberg MV, Getnet D, Bruno TC, Yen HR, Pyle KJ, Hipkiss E, Vignali DA, Pardoll DM, Drake CG: **Functionally distinct LAG-3 and PD-1 subsets on activated and chronically stimulated CD8 T cells.** *J Immunol* 2009, **182**:6659-6669.
99. Prigent P, El MS, Dreano M, Triebel F: **Lymphocyte activation gene-3 induces tumor regression and antitumor immune responses.** *Eur J Immunol* 1999, **29**:3867-3876.
100. Brignone C, Grygar C, Marcu M, Schakel K, Triebel F: **A soluble form of lymphocyte activation gene-3 (IMP321) induces activation of a large range of human effector cytotoxic cells.** *J Immunol* 2007, **179**:4202-4211.
101. Brignone C, Escudier B, Grygar C, Marcu M, Triebel F: **A phase I pharmacokinetic and biological correlative study of IMP321, a novel MHC class II agonist, in patients with advanced renal cell carcinoma.** *Clin Cancer Res* 2009, **15**:6225-6231.
102. Brignone C, Gutierrez M, Mefti F, Brain E, Jarcau R, Cvitkovic F, Boussetta N, Medioni J, Gligorov J, Grygar C *et al.*: **First-line chemioimmunotherapy in metastatic breast carcinoma: combination of paclitaxel and IMP321 (LAG-3lg) enhances immune responses and antitumor activity.** *J Transl Med* 2010, **8**:71.
103. Goldberg MV, Drake CG: **LAG-3 in cancer immunotherapy.** *Curr Top Microbiol Immunol* 2011, **344**:269-278.
104. Brewster AM, Hortobagyi GN, Broglio KR, Kau SW, Santa-Maria CA, Arun B, Buzdar AU, Booser DJ, Valero V, Bondy M *et al.*: **Residual risk of breast cancer recurrence 5 years after adjuvant therapy.** *J Natl Cancer Inst* 2008, **100**:1179-1183.
105. Chapoval AI, Ni J, Lau JS, Wilcox RA, Flies DB, Liu D, Dong H, Sica GL, Zhu G, Tamada K, Chen L: **B7-H3: a costimulatory molecule for T cell activation and IFN-gamma production.** *Nat Immunol* 2001, **2**:269-274.
106. Seliger B, Quandt D: **The expression, function, and clinical relevance of B7 family members in cancer.** *Cancer Immunol Immunother* 2012, **61**:1327-1341.
107. Sun Y, Wang Y, Zhao J, Gu M, Giscombe R, Lefvert AK, Wang X: **B7-H3 and B7-H4 expression in non-small-cell lung cancer.** *Lung Cancer* 2006, **53**:143-151.
108. Loos M, Hedderich DM, Ottenhausen M, Giese NA, Laschinger M, Esposito I, Kleeff J, Friess H: **Expression of the costimulatory molecule B7-H3 is associated with prolonged survival in human pancreatic cancer.** *BMC Cancer* 2009, **9**:463.
109. Zang X, Sullivan PS, Soslow RA, Waitz R, Reuter VE, Wilton A, Thaler HT, Arul M, Slovin SF, Wei J *et al.*: **Tumor associated endothelial expression of B7-H3 predicts survival in ovarian carcinomas.** *Mod Pathol* 2010, **23**:1104-1112.
110. Crispin PL, Sheinin Y, Roth TJ, Lohse CM, Kuntz SM, Frigola X, Thompson RH, Boorjian SA, Dong H, Leibovich BC *et al.*: **Tumor cell and tumor vasculature expression of B7-H3 predict survival in clear cell renal cell carcinoma.** *Clin Cancer Res* 2008, **14**:5150-5157.
111. Boorjian SA, Sheinin Y, Crispin PL, Farmer SA, Lohse CM, Kuntz SM, Leibovich BC, Kwon ED, Frank I: **T-cell coregulatory molecule expression in urothelial cell carcinoma: clinicopathologic correlations and association with survival.** *Clin Cancer Res* 2008, **14**:4800-4808.
112. Roth TJ, Sheinin Y, Lohse CM, Kuntz SM, Frigola X, Inman BA, Krambeck AE, McKenney ME, Karnes RJ, Blute ML *et al.*: **B7-H3 ligand expression by prostate cancer: a novel marker of**

- prognosis and potential target for therapy.** *Cancer Res* 2007, **67**:7893-7900.
113. Suh WK, Gajewska BU, Okada H, Gronski MA, Bertram EM, Dawicki W, Duncan GS, Bukczynski J, Plyte S, Elia A *et al.*: **The B7 family member B7-H3 preferentially down-regulates T helper type 1-mediated immune responses.** *Nat Immunol* 2003, **4**:899-906.
 114. Wang L, Kang FB, Shan BE: **B7-H3-mediated tumor immunology: friend or foe?** *Int J Cancer* 2014, **134**:2764-2771.
 115. Loos M, Hedderich DM, Friess H, Kleeff J: **B7-h3 and its role in antitumor immunity.** *Clin Dev Immunol* 2010, **2010**:683875.
 116. Yi KH, Chen L: **Fine tuning the immune response through B7-H3 and B7-H4.** *Immunol Rev* 2009, **229**:145-151.
 117. Loo D, Alderson RF, Chen FZ, Huang L, Zhang W, Gorlatov S, Burke S, Ciccarone V, Li H, Yang Y *et al.*: **Development of an Fc-enhanced anti-B7-H3 monoclonal antibody with potent antitumor activity.** *Clin Cancer Res* 2012, **18**:3834-3845.
 118. Sica GL, Choi IH, Zhu G, Tamada K, Wang SD, Tamura H, Chapoval AI, Flies DB, Bajorath J, Chen L: **B7-H4, a molecule of the B7 family, negatively regulates T cell immunity.** *Immunity* 2003, **18**:849-861.
 119. Prasad DV, Richards S, Mai XM, Dong C: **B7S1, a novel B7 family member that negatively regulates T cell activation.** *Immunity* 2003, **18**:863-873.
 120. Zang X, Loke P, Kim J, Murphy K, Waitz R, Allison JP: **B7x: a widely expressed B7 family member that inhibits T cell activation.** *Proc Natl Acad Sci U S A* 2003, **100**:10388-10392.
 121. Choi IH, Zhu G, Sica GL, Strome SE, Cheville JC, Lau JS, Zhu Y, Flies DB, Tamada K, Chen L: **Genomic organization and expression analysis of B7-H4, an immune inhibitory molecule of the B7 family.** *J Immunol* 2003, **171**:4650-4654.
 122. Tringler B, Liu W, Corral L, Torkko KC, Enomoto T, Davidson S, Lucia MS, Heinz DE, Papkoff J, Shroyer KR: **B7-H4 overexpression in ovarian tumors.** *Gynecol Oncol* 2006, **100**:44-52.
 123. Mugler KC, Singh M, Tringler B, Torkko KC, Liu W, Papkoff J, Shroyer KR: **B7-h4 expression in a range of breast pathology: correlation with tumor T-cell infiltration.** *Appl Immunohistochem Mol Morphol* 2007, **15**:363-370.
 124. Awadallah NS, Shroyer KR, Langer DA, Torkko KC, Chen YK, Bentz JS, Papkoff J, Liu W, Nash SR, Shah RJ: **Detection of B7-H4 and p53 in pancreatic cancer: potential role as a cytological diagnostic adjunct.** *Pancreas* 2008, **36**:200-206.
 125. Zang X, Thompson RH, Al-Ahmadie HA, Serio AM, Reuter VE, Eastham JA, Scardino PT, Sharma P, Allison JP: **B7-H3 and B7x are highly expressed in human prostate cancer and associated with disease spread and poor outcome.** *Proc Natl Acad Sci U S A* 2007, **104**:19458-19463.
 126. Krambeck AE, Thompson RH, Dong H, Lohse CM, Park ES, Kuntz SM, Leibovich BC, Blute ML, Cheville JC, Kwon ED: **B7-H4 expression in renal cell carcinoma and tumor vasculature: associations with cancer progression and survival.** *Proc Natl Acad Sci U S A* 2006, **103**:10391-10396.
 127. Yao Y, Wang X, Jin K, Zhu J, Wang Y, Xiong S, Mao Y, Zhou L: **B7-H4 is preferentially expressed in non-dividing brain tumor cells and in a subset of brain tumor stem-like cells.** *J Neurooncol* 2008, **89**:121-129.
 128. Jiang J, Zhu Y, Wu C, Shen Y, Wei W, Chen L, Zheng X, Sun J, Lu B, Zhang X: **Tumor expression of B7-H4 predicts poor survival of patients suffering from gastric cancer.** *Cancer Immunol Immunother* 2010, **59**:1707-1714.
 129. Kryczek I, Zou L, Rodriguez P, Zhu G, Wei S, Mottram P, Brumlik M, Cheng P, Curiel T, Myers L *et al.*: **B7-H4 expression identifies a novel suppressive macrophage population in human ovarian carcinoma.** *J Exp Med* 2006, **203**:871-881.
 130. Kryczek I, Wei S, Zou L, Zhu G, Mottram P, Xu H, Chen L, Zou W: **Cutting edge: induction of B7-H4 on APCs through IL-10: novel suppressive mode for regulatory T cells.** *J Immunol* 2006, **177**:40-44.
 131. Wang L, Rubinstein R, Lines JL, Wasiuk A, Ahonen C, Guo Y, Lu LF, Gondek D, Wang Y, Fava RA *et al.*: **VISTA, a novel mouse Ig superfamily ligand that negatively regulates T cell responses.** *J Exp Med* 2011, **208**:577-592.
- This article reports new member of immunoglobulin superfamily ligand, VISTA, which mainly expressed on hematopoietic cells. Its expression on APC inhibits T cell proliferation and cytokine production *in vitro*. This study also demonstrated the autoimmune phenomenon upon anti-VISTA treatment *in vivo* as well as impaired anti-tumor activity when it is over-expressed on tumor cells. It provides valid evidence of important role of VISTA in autoimmunity and immunosurveillance.
132. Flies DB, Wang S, Xu H, Chen L: **Cutting edge: a monoclonal antibody specific for the programmed death-1 homolog prevents graft-versus-host disease in mouse models.** *J Immunol* 2011, **187**:1537-1541.
 133. Norde WJ, Hobo W, van d V, Dolstra H: **Coinhibitory molecules in hematologic malignancies: targets for therapeutic intervention.** *Blood* 2012, **120**:728-736.
 134. Lines JL, Pantazi E, Mak J, Sempere LF, Wang L, O'Connell S, Ceeraz S, Suriawinata AA, Yan S, Ernstoff MS *et al.*: **VISTA is an immune checkpoint molecule for human T cells.** *Cancer Res* 2014, **74**:1924-1932.
 135. Le MI, Chen W, Lines JL, Day M, Li J, Sergeant P, Noelle RJ, Wang L: **VISTA regulates the development of protective antitumor immunity.** *Cancer Res* 2014, **74**:1933-1944.
- This article showed that the VISTA targeting monoclonal antibody treatment in mouse model demonstrated antitumor activity by which increasing number of tumor-specific T cells in the periphery and enhancing the infiltration, proliferation and effector function of tumor reactive T cells within tumor microenvironment.
136. Sakr MA, Takino T, Domoto T, Nakano H, Wong RW, Sasaki M, Nakanuma Y, Sato H: **GI24 enhances tumor invasiveness by regulating cell surface membrane-type 1 matrix metalloproteinase.** *Cancer Sci* 2010, **101**:2368-2374.
 137. Jung K, Choi I: **Emerging co-signaling networks in T cell immune regulation.** *Immune Netw* 2013, **13**:184-193.
 138. Beauchemin N, Draber P, Dveksler G, Gold P, Gray-Owen S, Grunert F, Hammarstrom S, Holmes KV, Karlsson A, Kuroki M *et al.*: **Redefined nomenclature for members of the carcinoembryonic antigen family.** *Exp Cell Res* 1999, **252**:243-249.
 139. Gray-Owen SD, Blumberg RS: **CEACAM1: contact-dependent control of immunity.** *Nat Rev Immunol* 2006, **6**:433-446.
 140. Kuespert K, Pils S, Hauck CR: **CEACAMs: their role in physiology and pathophysiology.** *Curr Opin Cell Biol* 2006, **18**:565-571.
 141. Gambichler T, Grothe S, Rotterdam S, Altmeyer P, Kreuter A: **Protein expression of carcinoembryonic antigen cell adhesion molecules in benign and malignant melanocytic skin lesions.** *Am J Clin Pathol* 2009, **131**:782-787.
 142. Thies A, Moll I, Berger J, Wagener C, Brummer J, Schulze HJ, Brunner G, Schumacher U: **CEACAM1 expression in cutaneous malignant melanoma predicts the development of metastatic disease.** *J Clin Oncol* 2002, **20**:2530-2536.
 143. Morales VM, Christ A, Watt SM, Kim HS, Johnson KW, Utku N, Texeira AM, Mizoguchi A, Mizoguchi E, Russell GJ *et al.*: **Regulation of human intestinal intraepithelial lymphocyte cytolytic function by biliary glycoprotein (CD66a).** *J Immunol* 1999, **163**:1363-1370.
 144. Markel G, Wolf D, Hanna J, Gazit R, Goldman-Wohl D, Lavy Y, Yagel S, Mandelboim O: **Pivotal role of CEACAM1 protein in the inhibition of activated decidual lymphocyte functions.** *J Clin Invest* 2002, **110**:943-953.
 145. Markel G, Lieberman N, Katz G, Arnon TI, Lotem M, Drize O, Blumberg RS, Bar-Haim E, Mader R, Eisenbach L *et al.*: **CD66a interactions between human melanoma and NK cells: a novel class I MHC-independent inhibitory mechanism of cytotoxicity.** *J Immunol* 2002, **168**:2803-2810.
 146. Huang YH, Zhu C, Kondo Y, Anderson AC, Gandhi A, Russell A, Dougan SK, Petersen BS, Melum E, Pertel T *et al.*: **CEACAM1**

- regulates TIM-3-mediated tolerance and exhaustion.** *Nature* 2015, **517**:386-390.
- This article demonstrates that TIM-3 is co-expressed and forms a heterodimer with CEACAM-1 which is another well known checkpoint inhibitory molecule expressed on activated T cell. It turns out that CEACAM-1 serves as a heterophilic ligand for TIM-3 that is required for its ability to inhibit T cell function. This study is highly relevant in translational point of view since TIM-3 targeting agent is currently under clinical development.
147. Ortenberg R, Sapir Y, Raz L, Hershkovitz L, Ben AA, Sapoznik S, Barshack I, Avivi C, Berkun Y, Besser MJ *et al.*: **Novel immunotherapy for malignant melanoma with a monoclonal antibody that blocks CEACAM1 homophilic interactions.** *Mol Cancer Ther* 2012, **11**:1300-1310.
 148. Watanabe N, Gavrieli M, Sedy JR, Yang J, Fallarino F, Loftin SK, Hurchla MA, Zimmerman N, Sim J, Zang X *et al.*: **BTLA is a lymphocyte inhibitory receptor with similarities to CTLA-4 and PD-1.** *Nat Immunol* 2003, **4**:670-679.
 149. Murphy TL, Murphy KM: **Slow down and survive: enigmatic immunoregulation by BTLA and HVEM.** *Annu Rev Immunol* 2010, **28**:389-411.
 150. Sedy JR, Gavrieli M, Potter KG, Hurchla MA, Lindsley RC, Hildner K, Scheu S, Pfeffer K, Ware CF, Murphy TL *et al.*: **B and T lymphocyte attenuator regulates T cell activation through interaction with herpesvirus entry mediator.** *Nat Immunol* 2005, **6**:90-98.
 151. Steinberg MW, Cheung TC, Ware CF: **The signaling networks of the herpesvirus entry mediator (TNFRSF14) in immune regulation.** *Immunol Rev* 2011, **244**:169-187.
 152. del Rio ML, Lucas CL, Buhler L, Rayat G, Rodriguez-Barbosa JL: **HVEM/LIGHT/BTLA/CD160 cosignaling pathways as targets for immune regulation.** *J Leukoc Biol* 2010, **87**:223-235.
 153. Cai G, Anumanthan A, Brown JA, Greenfield EA, Zhu B, Freeman GJ: **CD160 inhibits activation of human CD4⁺ T cells through interaction with herpesvirus entry mediator.** *Nat Immunol* 2008, **9**:176-185.
 154. Cai G, Freeman GJ: **The CD160, BTLA, LIGHT/HVEM pathway: a bidirectional switch regulating T-cell activation.** *Immunol Rev* 2009, **229**:244-258.
 155. M'Hidi H, Thibault ML, Chetaille B, Rey F, Bouadallah R, Nicolas R, Olive D, Xerri L: **High expression of the inhibitory receptor BTLA in T-follicular helper cells and in B-cell small lymphocytic lymphoma/chronic lymphocytic leukemia.** *Am J Clin Pathol* 2009, **132**:589-596.
 156. Pasero C, Speiser DE, Derre L, Olive D: **The HVEM network: new directions in targeting novel costimulatory/co-inhibitory molecules for cancer therapy.** *Curr Opin Pharmacol* 2012, **12**:478-485.
 157. Baitsch L, Baumgaertner P, Devereux E, Raghav SK, Legat A, Barba L, Wieckowski S, Bouzourene H, Deplancke B, Romero P *et al.*: **Exhaustion of tumor-specific CD8⁺ T cells in metastases from melanoma patients.** *J Clin Invest* 2011, **121**:2350-2360.
 158. Derre L, Rivals JP, Jandus C, Pastor S, Rimoldi D, Romero P, Michielin O, Olive D, Speiser DE: **BTLA mediates inhibition of human tumor-specific CD8⁺ T cells that can be partially reversed by vaccination.** *J Clin Invest* 2010, **120**:157-167.
 159. Fourcade J, Sun Z, Pagliano O, Guillaume P, Luescher IF, Sander C, Kirkwood JM, Olive D, Kuchroo V, Zarour HM: **CD8⁺ T cells specific for tumor antigens can be rendered dysfunctional by the tumor microenvironment through upregulation of the inhibitory receptors BTLA and PD-1.** *Cancer Res* 2012, **72**:887-896.
 160. Jenmalm MC, Cherwinski H, Bowman EP, Phillips JH, Sedgwick JD: **Regulation of myeloid cell function through the CD200 receptor.** *J Immunol* 2006, **176**:191-199.
 161. Minas K, Liversidge J: **Is the CD200/CD200 receptor interaction more than just a myeloid cell inhibitory signal?** *Crit Rev Immunol* 2006, **26**:213-230.
 162. Barclay AN, Clark MJ, McCaughan GW: **Neuronal/lymphoid membrane glycoprotein MRC OX-2 is a member of the immunoglobulin superfamily with a light-chain-like structure.** *Biochem Soc Symp* 1986, **51**:149-157.
 163. Snelgrove RJ, Goulding J, Didierlaurent AM, Lyonga D, Vekaria S, Edwards L, Gwyer E, Sedgwick JD, Barclay AN, Russell T: **A critical function for CD200 in lung immune homeostasis and the severity of influenza infection.** *Nat Immunol* 2008, **9**:1074-1083.
 164. Rygiel TP, Rijkers ES, de Ruiter T, Stolte EH, van der Valk M, Rimmelzwaan GF, Boon L, van Loon AM, Coenjaerts FE, Hoek RM *et al.*: **Lack of CD200 enhances pathological T cell responses during influenza infection.** *J Immunol* 2009, **183**:1990-1996.
 165. Tonks A, Hills R, White P, Rosie B, Mills KI, Burnett AK, Darley RL: **CD200 as a prognostic factor in acute myeloid leukaemia.** *Leukemia* 2007, **21**:566-568.
 166. Moreaux J, Hose D, Reme T, Jourdan E, Hundemer M, Legouffe E, Moine P, Bourin P, Moos M, Corre J *et al.*: **CD200 is a new prognostic factor in multiple myeloma.** *Blood* 2006, **108**:4194-4197.
 167. Siva A, Xin H, Qin F, Oltean D, Bowdish KS, Kretz-Rommel A: **Immune modulation by melanoma and ovarian tumor cells through expression of the immunosuppressive molecule CD200.** *Cancer Immunol Immunother* 2008, **57**:987-996.
 168. Coles SJ, Wang EC, Man S, Hills RK, Burnett AK, Tonks A, Darley RL: **CD200 expression suppresses natural killer cell function and directly inhibits patient anti-tumor response in acute myeloid leukemia.** *Leukemia* 2011, **25**:792-799.
 169. Gorczynski RM, Chen Z, Diao J, Khatri I, Wong K, Yu K, Behnke J: **Breast cancer cell CD200 expression regulates immune response to EMT6 tumor cells in mice.** *Breast Cancer Res Treat* 2010, **123**:405-415.
 170. de Visser KE: **Spontaneous immune responses to sporadic tumors: tumor-promoting, tumor-protective or both?** *Cancer Immunol Immunother* 2008, **57**:1531-1539.
 171. Stumpfova M, Ratner D, Desciak EB, Eliezi YD, Owens DM: **The immunosuppressive surface ligand CD200 augments the metastatic capacity of squamous cell carcinoma.** *Cancer Res* 2010, **70**:2962-2972.
 172. Rygiel TP, Meynard L: **CD200R signaling in tumor tolerance and inflammation: a tricky balance.** *Curr Opin Immunol* 2012, **24**:233-238.
- Excellent review on the roles of CD200-CD200R immunomodulatory pathway in tumor as well as inflammation. This review raises the concerns for potential adverse outcome of CD200 blockade in tumors from increased inflammation.
173. Joller N, Peters A, Anderson AC, Kuchroo VK: **Immune checkpoints in central nervous system autoimmunity.** *Immunol Rev* 2012, **248**:122-139.
 174. Yu X, Harden K, Gonzalez LC, Francesco S, Chiang E, Irving B, Tom I, Ivelja S, Refino CJ, Clark H *et al.*: **The surface protein TIGIT suppresses T cell activation by promoting the generation of mature immunoregulatory dendritic cells.** *Nat Immunol* 2009, **10**:48-57.
 175. Bottino C, Castriconi R, Pende D, Rivera P, Nanni M, Carnemolla B, Cantoni C, Grassi J, Marcenaro S, Reymond N *et al.*: **Identification of PVR (CD155) and Nectin-2 (CD112) as cell surface ligands for the human DNAM-1 (CD226) activating molecule.** *J Exp Med* 2003, **198**:557-567.
 176. Lozano E, Dominguez-Villar M, Kuchroo V, Hafler DA: **The TIGIT/CD226 axis regulates human T cell function.** *J Immunol* 2012, **188**:3869-3875.
 177. Hafler JP, Maier LM, Cooper JD, Plagnol V, Hinks A, Simmonds MJ, Stevens HE, Walker NM, Healy B, Howson JM *et al.*: **CD226 Gly307Ser association with multiple autoimmune diseases.** *Genes Immun* 2009, **10**:5-10.
 178. Maiti AK, Kim-Howard X, Viswanathan P, Guillen L, Qian X, Rojas-Villarraga A, Sun C, Canas C, Tobon GJ, Matsuda K *et al.*: **Non-synonymous variant (Gly307Ser) in CD226 is associated with susceptibility to multiple autoimmune diseases.** *Rheumatology (Oxford)* 2010, **49**:1239-1244.

179. Joller N, Hafler JP, Brynedaal B, Kassam N, Spoerl S, Levin SD, Sharpe AH, Kuchroo VK: **Cutting edge: TIGIT has T cell-intrinsic inhibitory functions.** *J Immunol* 2011, **186**:1338-1342.
180. Levin SD, Taft DW, Brandt CS, Bucher C, Howard ED, Chadwick EM, Johnston J, Hammond A, Bontadelli K, Ardourel D *et al.*: **Vstm3 is a member of the CD28 family and an important modulator of T-cell function.** *Eur J Immunol* 2011, **41**:902-915.
181. Curran MA, Montalvo W, Yagita H, Allison JP: **PD-1 and CTLA-4 combination blockade expands infiltrating T cells and reduces regulatory T and myeloid cells within B16 melanoma tumors.** *Proc Natl Acad Sci U S A* 2010, **107**:4275-4280.
- This is excellent preclinical model of combined checkpoint blockade of PD-1 and CTLA-4 which is in clinical development. It shows that combined blockade is more than twice as effective as either alone in promoting the rejection of B16 melanoma in conjunction with GM-CSF.
182. Intlekofer AM, Thompson CB: **At the bench: preclinical rationale for CTLA-4 and PD-1 blockade as cancer immunotherapy.** *J Leukoc Biol* 2013, **94**:25-39.
183. Wolchok JD, Kluger H, Callahan MK, Postow MA, Rizvi NA, Lesokhin AM, Segal NH, Ariyan CE, Gordon RA, Reed K *et al.*: **Nivolumab plus ipilimumab in advanced melanoma.** *N Engl J Med* 2013, **369**:122-133.
- The first clinical trial combining two immune checkpoint antibodies resulting in increased antitumor activity, but at the risk of increased autoimmune toxicities.
184. Sznol M, Kluger HM, Callahan MK, Postow MA, Gordon RA, Segal NH, Rizvi NA, Lesokhin AM, Atkins MB, Kirkwood JM *et al.*: **Survival, response duration, and activity by BRAF mutation (MT) status of nivolumab (NIVO, anti-PD-1, BMS-936558, ONO-4538) and ipilimumab (IPI) concurrent therapy in advanced melanoma (MEL).** *2014 ASCO Annual Meeting*. 2014. LBA9003 (abstract).
185. Duraiswamy J, Kaluza KM, Freeman GJ, Coukos G: **Dual blockade of PD-1 and CTLA-4 combined with tumor vaccine effectively restores T-cell rejection function in tumors.** *Cancer Res* 2013, **73**:3591-3603.
186. McArthur GA, Ribas A: **Targeting oncogenic drivers and the immune system in melanoma.** *J Clin Oncol* 2013, **31**:499-506.
187. Boni A, Cogdill AP, Dang P, Udayakumar D, Njauw CN, Sloss CM, Ferrone CR, Flaherty KT, Lawrence DP, Fisher DE *et al.*: **Selective BRAFV600E inhibition enhances T-cell recognition of melanoma without affecting lymphocyte function.** *Cancer Res* 2010, **70**:5213-5219.
- MAPK inhibitors are now standard of care for BRAF mutated patients and combined immunotherapy with MAPK inhibitors are being actively investigated. This paper describes that MAPK inhibitors increase T cell recognition of melanoma antigens, however, MEK inhibitor impairs T cell functionality whereas specific BRAF inhibitor does not. This study has important implication for ongoing and future studies for combined immunotherapy with MAPK inhibitors.
188. Koya RC, Mok S, Otte N, Blacketer KJ, Comin-Anduix B, Tumei PC, Minasyan A, Graham NA, Graeber TG, Chodon T *et al.*: **BRAF inhibitor vemurafenib improves the antitumor activity of adoptive cell immunotherapy.** *Cancer Res* 2012, **72**:3928-3937.
189. Liu C, Peng W, Xu C, Lou Y, Zhang M, Wargo JA, Chen JQ, Li HS, Watowich SS, Yang Y *et al.*: **BRAF inhibition increases tumor infiltration by T cells and enhances the antitumor activity of adoptive immunotherapy in mice.** *Clin Cancer Res* 2013, **19**:393-403.
190. Frederick DT, Piris A, Cogdill AP, Cooper ZA, Lezcano C, Ferrone CR, Mitra D, Boni A, Newton LP, Liu C *et al.*: **BRAF inhibition is associated with enhanced melanoma antigen expression and a more favorable tumor microenvironment in patients with metastatic melanoma.** *Clin Cancer Res* 2013, **19**:1225-1231.
191. Ribas A, Hodi FS, Callahan M, Konto C, Wolchok J: **Hepatotoxicity with combination of vemurafenib and ipilimumab.** *N Engl J Med* 2013, **368**:1365-1366.
192. Ott PA, Henry T, Baranda SJ, Frlita D, Manches O, Bogunovic D, Bhardwaj N: **Inhibition of both BRAF and MEK in BRAF(V600E) mutant melanoma restores compromised dendritic cell (DC) function while having differential direct effects on DC properties.** *Cancer Immunol Immunother* 2013, **62**:811-822.
193. Dubovsky JA, Beckwith KA, Natarajan G, Woyach JA, Jaglowski S, Zhong Y, Hessler JD, Liu TM, Chang BY, Larkin KM *et al.*: **Ibrutinib is an irreversible molecular inhibitor of ITK driving a Th1-selective pressure in T lymphocytes.** *Blood* 2013, **122**:2539-2549.
- Small molecule inhibitor, ibrutinib, is approved for chronic lymphocytic leukemia and mantle cell lymphoma and has good activity on activated B cell type diffuse large B cell lymphoma. It is a potent inhibitor of bruton tyrosine kinase (BTK), suggested to inhibit other important pathways. In this particular study, authors demonstrated that ibrutinib inhibits interleukin-2 inducible kinase (ITK) which promotes development of T cells toward a Th1 phenotype and results in improved outcome in infectious disease model. This study suggests that we may apply ibrutinib to variety types of malignancies.
194. Vargas L, Hamasy A, Nore BF, Smith CI: **Inhibitors of BTK and ITK: state of the new drugs for cancer, autoimmunity and inflammatory diseases.** *Scand J Immunol* 2013, **78**:130-139.
195. Ali K, Soond DR, Pineiro R, Hagemann T, Pearce W, Lim EL, Bouabe H, Scudamore CL, Hancox T, Maecker H *et al.*: **Inactivation of PI(3)K p110delta breaks regulatory T-cell-mediated immune tolerance to cancer.** *Nature* 2014, **510**:407-411.
- It was thought that phosphatidylinositol-3-kinase δ (PI3K δ) inhibition has only activity on hematological cancers. However, this study demonstrated that dominant Treg-cell dysfunction through PI3K δ inhibition augments T cell mediated anticancer immunity. This study also provided the evidence that PI3K δ inactivation suppressed the myeloid-derived suppressor cells. It would be very interesting to see whether PI3K δ inhibition synergizes with current immunotherapies, such as adoptive cell transfer and checkpoint blockade.
196. Kingwell K: **Anticancer drugs: horizons broaden for PI3Kdelta inhibitors.** *Nat Rev Drug Discov* 2014, **13**:573.