

Macroecological and macroevolutionary perspectives on the Cenozoic history of South American Mammals

Lucas Buffan

Table of contents

Foreword and Acknowledgements	i
Foreword	i
Acknowledgements	i
1 The unevenness of biodiversity: corner stone for the large-scale study of life	1
1.1 The heterogeneous organisation of life across space	1
1.2 From peaks to declines: temporally-heterogeneous biodiversity	2
1.3 Inter-clade heterogeneity	5
1.4 Methodological biases arising when studying biodiversity	7
1.4.1 Biodiversity shortfalls	7
1.4.2 Propagation of the caveats of the fossil record when studying biodiversity through time	7
2 Disentangling biodiversity patterns across space and time: the macro-eco-evolutionary perspective	8
2.1 Context and terminology	8
2.2 Macroevolution: evolution above the species level	8
2.2.1 Tempo of Macroevolution: how do rates change across time?	8
2.2.2 Mode of Macroevolution: what drives macroevolutionary rate variations?	9
2.3 Macro-eco-evolutionary understanding of the spatial organisation of life: insights from the Latitudinal Biodiversity Gradient	10
3 Analytical tools to study large-scale biodiversity patterns	12
3.1 Macroevolutionary approaches to study how clades thrive and go extinct	12
3.1.1 Phylogeny-based	12
3.1.2 Fossil occurrence-based	12
3.1.3 Towards integrating multiple sources of data	12
3.2 Integrating ecological processes with evolutionary forces: mechanistic eco-evolutionary simulations	12
4 A strange biodiversity on a strange land : the case of mammals in South America	13
4.1 South America across the Cenozoic: geographical and environmental context	13
4.1.1 Amazonia: a unique and dynamic tropical ecoregion	13
4.1.2 The rise of the Andes : emergence of one of the largest and highest mountain chains, and its consequences on biodiversity	13
4.1.3 Intermediate flooding events	13
4.1.4 South America as a playground for evolutionary biologists	13
4.2 ‘Splendid isolation’: the captivating history of South American mammals across the Cenozoic	13
4.2.1 History of mammal palaeontology in South America: a brief overview	14
4.2.2 First stratum: endemic radiations (Late Cretaceous–middle Eocene)	15

4.2.3	Second stratum: Trans-atlantic migrations and faunal ‘modernisation’ (late Eocene–Late Miocene)	18
4.2.4	Third stratum: Inter-american exchange, massive faunal disruptions, and establishment of the extant mammal biodiversity (latest Miocene–Present) . . .	20
4.3	Challenges for South American mammal palaeontology	20
5	Objectives of this doctoral work	23
5.1	Deciphering the dynamics of diversification of South American mammals and their likely drivers across the Cenozoic	23
5.1.1	Elucidating the impact of the Eocene–Oligocene Transition on the macroevolutionary dynamics of South American mammals	23
5.1.2	Did the evolutionary pathways towards increasing in body size and hypsodonty impact the diversification of chinchilloid rodents?	23
5.2	Assessing the role of palaeoclimate in the emergence of the asymmetry behind the Great American Biotic Interchange	23
	Chapter 1. Deciphering the trends and drivers of South American mammal large-scale evolution across the Cenozoic	24
	Chapter 2.1. The fate of South America’s endemic mammalian fauna in response to the most dramatic Cenozoic climate disruption	24
	Chapter 2.2. Continent-scale insights into the last 23 million years of South American mammalian evolution	24
	Chapter 2. A matter of size: what underpinned the rise and demise of the largest rodents that ever lived?	25
	Chapter 3. Palaeoclimate alone fails to explain the asymmetry of the Great American Biotic Interchange	26
	Discussion	27
	Limitations	27
	Terminology, definitions and other macroevolutionary enigmas	27
	Perspectives	27
	An almost closed system: founding migrations in the Palaeogene history of South American mammals	27
	Sustainable late Cenozoic connection to North America: what drove the asymmetry of the GABI?	27
	Future directions towards an improved understanding of the emergence and dynamics of South American ecosystems	28
	Concluding remarks	29
	Glossary	30
	Literature cited	32

Foreword and Acknowledgements

Foreword

Acknowledgements

1 The unevenness of biodiversity: corner stone for the large-scale study of life

The astonishing diversity of life on Earth has fascinated scientists for centuries. A study dating back to 15 years ago indeed estimated that around 8.7 million eukaryotic species are currently coexisting on our planet (Mora *et al.* 2011). This number would probably be significantly increased by accounting for the substantial improvements that have occurred in the field of systematics over the recent years, and expanded by orders of magnitude if prokaryotic species were incorporated. Yet, someone travelling to the Sahara desert would probably be less amazed by the surrounding diversity of life than someone travelling to the middle of the Amazonia, one of the most species-rich regions in the world (Raven *et al.* 2020). Another vertiginous data is that around 99.9% of the species that ever existed since the dawn of life on Earth, some 4 billion years ago, has gone extinct. It therefore stands out that, far from a fixed and smooth entity, biodiversity, here understood as species richness, presents a myriad of heterogeneities, that stimulated a vast quantity of research works in which this thesis happens to be nested.

1.1 The heterogeneous organisation of life across space

For more than two centuries, it has been recognised that species are not randomly distributed on Earth. In his pioneering “*Essay on the geography of plants*”, regarded as foundational for the science of Biogeography (see Glossary), Alexander von Humboldt emphasised the spatial nature of plant taxonomic assemblages (Humboldt & Bonpland 1807). Later in the 19th century, the scientific community acquired a uniquely integrated perspective on biodiversity fed by years of field observations, giving to the question of the spatial organisation of life on Earth a new dimension. Consequently, it has been proposed to subdivide the world into spatial units sharing similar fauna/flora, sometimes called ‘Homozoic Belts’, ‘Regions’, ‘Realms’, ‘Provinces’, ‘Areas’ or ‘*Gebiet*’ (German word for territory, area) (Burkill 1943). The history will mainly remember the efforts of Alfred R. Wallace in that respect, who relied on the subdivision made by Philip L. Sclater in 1874 based on mammals, as well as his own expertise, to establish a famous contouring on the earth’s ‘Zoogeographic Realms’ (Figure 1.1), two years after (Wallace 1876). Further efforts towards incorporating fossil evidence to better constrain the aforementioned subdivision have to be acknowledged, notably implused by Richard Lydekker (Lydekker 1896). Wallace’s subdivision has maintained in the field since its publication, and only received minor updates in the last decade (Holt *et al.* 2013), illustrating the degree of knowledge advancement of the English naturalist, in a context where Charles Darwin’s famous Theory of Evolution – that, actually, Wallace co-discovered in a totally independent manner, making such a theory one of the most striking examples of thought convergence in the history of sciences (Berry & Browne 2008) – still had a long way to go before being formally accepted by the broad scientific community.

Beyond its spatial organisation into units of shared taxonomic composition, biodiversity also



Figure 1.1: Biogeographic map based on faunal similarities within animal assemblages across space ('Zoogeographic Realms') proposed by Wallace (1876).

exhibits a striking latitudinal organisation, in which, species richness increases with decreasing latitude, known as the Latitudinal Biodiversity Gradient (Figure 1.2). Such a pattern was already recognised by Humboldt. According to him, low-latitude plant tropical assemblages would be essentially made up with a large number of different species per unit of area, each having a relatively low density ("isolated and sparse"). Conversely, higher-latitude temperate or cold assemblages would be characterised by a substantially lower species richness per unit of area, with a single dominant species being over-represented, "excluding [...] heterogeneous species" (Humboldt & Bonpland 1807). Half a century later, in his cornerstone book *"The Origin of Species"*, Darwin dedicated two chapters to the spatial distribution of species, recognising that many groups harbour a greater diversity in the tropics and seeking for potential mechanisms underlying this pattern (Darwin 1859). Today, though notable exceptions exist (e.g., pinnipeds, lagomorphs or several kinds of fungi), a latitudinal diversity gradient has been recognised among most taxonomic group (e.g., plants, insects, molluscs, vertebrates) (Hillebrand 2004), as depicted on Figure 1.2 for three vertebrate classes.

Hence, biodiversity is largely heterogeneous across space, raising questions that have been thoroughly investigated in the past, as to 'Why are there more species in the tropics than in the poles?' or, more generally, 'What constraints the distribution of species on our planet?'

1.2 From peaks to declines: temporally-heterogeneous biodiversity

The aforementioned complex spatial arrangement of life is in fact the legacy of hundreds of millions of years in which species have been fashioned by a wealth of forces – of which we intend to provide an overview in Section 2.2.2. The statement that species experience changes across time, and that these species-level temporal changes eventually scale up to multispecies assemblages making up biodiversity is now commonly accepted, mostly thanks to fossil data.




Figure 1.2: Extant land mammal (A), non-marine bird (B) and amphibian (C) species richness at the surface of the globe, expressed in number of species, with a 10 x 10km resolution. Data are from <https://biodiversitymapping.org/>

Being the only source of direct evidence of life in the past, fossils have indeed been of immense value to characterise the temporal dynamics of biodiversity, notably since they irrefutably document the occurrence of extinction in the history of life. It is however interesting to notice that the reasoning that makes us conclude that if, for instance, mammoths are found in the fossil record but not among extant assemblages, it is because they have gone extinct, has not always been as straightforward as it appears to us. Fossils have indeed puzzled eminent figures in the field of biology, such as Jean-Baptiste de Lamarck – to whom we incidentally owe the name ‘biology’. In the early 1800s, Lamarck tentatively formalised a theory where animals would be subject to a complexifying force driving body plans towards higher levels. According to his theory, the (animal) biosphere would thus represent a continuum where species would be unidirectionally connected between each others, which, importantly, leaves no room for extinction – or any sort of species richness variation across time – : every member of Lamarck’s gradual sequence towards increasingly-complex body plans would already exist (Lamarck 1815). However, Lamarck had to cope with the fact that the field of systematic palaeontology was emerging in parallel, notably impelled by his colleagues from the Muséum National d’Histoire Naturelle de Paris, including Georges Cuvier (Servais *et al.* 2012). Plenty of remains of animals trapped inside the rocks but also manifestly absent from the surface of the (known) world were thus unearthed and described, challenging the Lamarck’s model. To intend to make sense of this apparent inadequacy, Lamarck therefore formulated a famous *ad-hoc* hypothesis: the apparent absence of extant representatives of mammoths does not mean that mammoths have gone extinct. If they indeed locally disappeared from the places where they left fossil remains, they must have found refugia in a region of the globe unexplored by the occidental civilisation.

Markedly opposed to Lamarck’s theory of animal organisation following increasingly complex body plans, Cuvier was maybe one of the earliest scientists to propose that species assemblages could change across time. His rich fieldwork expertise led him to postulate that the history of the globe has been punctuated by abrupt ‘revolution’ events, which can be viewed as (abiotic) changes occurring within a given environment, in turn causing changes in the nature of the species present within this changing environment (Cuvier 1840). Two decades later, John Phillips, an English palaeontologist known as the father of the Phanerozoic eras (Palaeozoic, Mesozoic and Cenozoic)

compiled for the first time fossil occurrences of marine taxa from the British lithological record, that he used to produce the first continuous diversity-through-time curve, spanning the entire Phanerozoic (Phillips 1860). His work allowed him to discriminate what he called the Palaeozoic, Mesozoic and Caenozoic ‘lives’. These three ‘lives’ were separated by vast depressions, corresponding to the Permo–Triassic and Cretaceous–Palaeogene boundaries (Figure 1.3 A), that will be later identified as two amongst the five most violent extinction episodes that marked the history of life, also known as mass extinction crises (see Section 2.2.1). Phillips’ cornerstone compilation thus, for the first time, quantitatively demonstrated the heterogeneity of species richness across geological times.

A century later, the rise of quantitative palaeobiology (see Section 1.4.2) paved the way towards more accurate reconstructions of biodiversity across time. The publication, in the early 1950s, of the first edition of the *Treatise of Invertebrate Paleontology* provided an access to a vast quantity of marine invertebrate fossil data, that could be further used for such an achievement. In that framework, Norman Newell reconstructed diversity curves for about 16 different groups (Newell 1952). Again, the reconstructed curves displayed time-varying patterns. Newell noticed certain similarities in the reconstructed patterns between groups, that he suggested to be evidence for “common causes” underlying their large-scale evolutionary trends. A few decades later, with a higher-resolution curve, Valentine (1969) (Figure 1.3 B) proposed to “explain diversity trends in terms of ecological hierarchy” (Sepkoski 2012), constituting a significant advance towards integrating ecological and evolutionary processes (see Chapter 2).




Figure 1.3: Historical fossil-based reconstructions of marine diversity across geological times produced by Phillips (1860), using a marine invertebrate compilation from the British lithological record (A) and by Valentine (1969), using marine invertebrate record from the *Treatise of Invertebrate Paleontology* (B). Note that timescale in (A) has been reversed when compared (B), going back in time from left to right instead of right to left.

However, Valentine’s reconstructions raised some methodological criticism, especially expressed by David M. Raup, one of the leading figures of the ‘palaeobiological revolution’. Raup was indeed concerned about how the fossil record biases may affect Valentine’s reconstruction, importantly stressing that “the increase in the number of marine species across the Phanerozoic may be more apparent than real” (Raup 1972, see Section 1.4.2). Ten years of intense data compilation and methodological labour after, together with J. John Sepkoski Jr. (another major actor of the ‘palaeobiological revolution’) Raup came to the conclusion that marine biodiversity indeed experienced temporal variations, persisting even after accounting for biases inherent to the fossil record (Sep-

koski *et al.* 1981). A year after, Raup & Sepkoski (1982) characterised major temporal features in their diversity curve (Figure 1.4), now known as the ‘Big Five mass extinctions’ (that we will develop in further details in Section 2.2.1).

Nowadays, it makes no doubt that biodiversity is temporally heterogeneous. Clades originate, thrive and then possibly decline or even go extinct, sometimes fluctuating in species number in between. A core aim of macroevolution, in which this doctoral work is nested, is thus to elucidate, given a clade, how species diversity changes across time, and the factors underlying these changes (Section 2.2).




Figure 1.4: Temporal variations in marine animal biodiversity across the Phanerozoic. Colours indicate the three evolutionary faunas discriminated by Sepkoski (1984). Taxa that could not be assigned to any fauna are included in the clearest band. Arrows indicate the “Big Five” mass extinctions, as defined by Raup & Sepkoski (1982). Data are from the genus-level compendium released by Sepkoski (2002), binned across geological stages, accessed and displayed thanks to the R package `sepkoski` (Jones 2022). Geological timescale was added thanks to the `deeptime` R package (Gearty 2023).

1.3 Inter-clade heterogeneity

Another facet of the heterogeneity of biodiversity comes from the fact that some groups harbour an immensely larger species richness than others. This striking feature has intrigued evolutionary biologists since the times of Darwin. In a 1879 letter to Joseph Hooker, the renowned white-bearded English naturalist indeed famously termed the astonishing diversity of flowering plants (Angiosperma) and their apparent rapid burst in the fossil record as an “abominable mystery”. At present, with more than 300,000 described species, angiosperms encompass nearly 90% of the extant land plant diversity.

Over the past few decades, the rise of phylogenetics (Section 3.1.1) permitted to better constrain evolutionary relationships between lineages, thus highlighting inter-clade diversity disequilibria (Fig-

1.4 Methodological biases arising when studying biodiversity

1.4.1 Biodiversity shortfalls

Several caveats apply to the study of biodiversity patterns ‘from a distance’, the baseline of the macroecological perspective (Diniz-Filho 2023). Some of them are grouped into what is known as the ‘biodiversity shortfalls’, representing inherent limitations related to missing information in biodiversity data, thereby inevitably hampering accurate understanding of the modelled eco-evolutionary processes. Three biodiversity shortfalls are usually considered, though Hortal *et al.* (2015) in fact more finely discriminated seven shortfalls – that will not be extensively covered here. These three shortfalls have been named after three renowned figures of ecology, evolution and systematics.

The Linnean shortfall reflects to the taxonomic uncertainty, that is, the degree of certainty that one may have, after sampling two individuals at random in a community, to tell whether they belong to the same species or to two distinct species. Next, the Wallacean shortfall refers to the knowledge gaps in species distribution, *i.e.*, the fact that a species may be somewhere where we did not observe it; thus this place in particular would be excluded from upstream range map reconstructions of that species (sort of type-1 error applied to biodiversity data). Last, the Darwinian shortfall refers to the lack of knowledge of the evolutionary relationships between species.

1.4.2 Propagation of the caveats of the fossil record when studying biodiversity through time

When considering fossil data, the previously described biodiversity shortfalls take a new dimension.

Additional biases emerge as one intends to use fossil data to provide a temporal perspective on biodiversity reconstructions, merely coming from the inherent incompleteness of the fossil record.

Darwin himself recognised the imperfection of the fossil record, comparing the latter to a book written in different dialects with many missing pages, and with only a few lines preserved on the remaining pages Darwin (1859). This conception of the fossil record, too sparse to be accurately used to reconstruct evolutionary processes (“Darwin’s dilemma”, Sepkoski (2012)) was largely shared among the evolutionary biology community, and resulted in the marginalisation of palaeontology from the rest of the community during most of the 20th century. Roughly between 1970 and 1985, a massive change of paradigm occurred as some palaeontologists started to focus on large-scale analyses of evolutionary patterns using quantitative methods, known as the “palaeobiological revolution” Sepkoski (2012).

Early in the history of fossil-based reconstructions of diversity through time, scientists expressed concerns regarding sampling-related biases. In its cornerstone reconstruction, Phillips (1860) had the idea of controlling the richness estimates he obtained per bins with the thickness of the corresponding stratigraphic section. Diversity through time controversy: Raup (1972) and Sepkoski (2012)

Signor & Lipps effect Signor & Lipps (1982)

Close *et al.* (2020)

Thus, studying large-scale biodiversity requires embracing uncertainty!

2 Disentangling biodiversity patterns across space and time: the macro-eco-evolutionary perspective

2.1 Context and terminology

Key concepts and terminology

Speciation, extinction, diversification

Key relating process: biogeography.

2.2 Macroevolution: evolution above the species level

Macroevolution is usually understood as the study of evolutionary patterns occurring above the species level ([Hautmann 2020](#); [Saupe & Myers 2021](#)). Macroevolution has been classically approached under two non-mutually exclusive lenses ([Stanley 1979](#)), either focusing on (1) the diversification process (see Section 2.1 and [Glossary](#)), or (2) the evolutionary dynamics of phenotypes – as an echo to the term ‘diversification’, some authors refer to the process underlying morphological diversity as ‘disparification’ (*e.g.*, [Davis et al. 2016](#); [Nürk et al. 2019](#)). Unfortunately, despite their undeniable appeal, questions related to the dynamics of morphological evolution fall a bit outside the scope of this thesis. Thus, in this manuscript, emphasis will be placed on the tempo and mode of clade diversification.

Just like any other field of evolutionary biology ([Simpson 1944](#)), studying macroevolution is a matter of tempo (*i.e.*, the rate at which macroevolutionary changes occur, namely, speciation, extinction or morphological evolution rates) mode (*i.e.*, the forces underlying these changes) ([Saupe & Myers 2021](#)).

2.2.1 Tempo of Macroevolution: how do rates change across time?

How speciation and extinction rates vary through time mirrors the aforementioned changes in clade diversity across geological times (Section 1.2).

About evolutionary constraints: Raup ([1966](#))

- early acquisition of hovering flight and nectarivory among birds likely promoted the radiation of hummingbirds (Trochilidae) ([Mayr 2004](#))
- early acquisition of echolocation likely favoured the radiation of bats ([Hand et al. 2023](#))

- tooth loss and reappraisal in other diverse forms, believed to have leveraged the morphological constraints imposed by the tribosphenic geometry Vizcaíno (2009).
- flower, along with other traits related to physiology (faster pace of living) promoted angiosperm radiation, Darwin's 'second abominable mystery' (Benton *et al.* 2022; Sauquet *et al.* 2017)

Mass extinctions: Raup & Sepkoski (1982) Raup (1986), Sepkoski (2012) chap. 9

Biodiversity crisis. Parallel with the present-day

2.2.2 Mode of Macroevolution: what drives macroevolutionary rate variations?

It has long been established that biodiversity exhibits marked variations through time, that could not be explained by random fluctuations alone (Sepkoski *et al.* 1981). Since temporal variations of biodiversity can be either understood in terms of pace at which species originate or go extinct, this conclusion suggest that external factors may directly act on the speciation or the extinction processes. Seminal theories have been proposed to relate such environmental dynamics to the pace at which large-scale evolution occurs.

2.2.2.1 Abiotic factors: the 'Court Jester' hypothesis

Despite the relatively recent formulation of the 'Court Jester' model (Barnosky 2001), the idea that abiotic factors may affect diversification rates has been in the minds for a long time.

Palaeogeography, climate

2.2.2.2 Biotic factors: the 'Red Queen' hypothesis

A turning point in the appreciation of the macroevolutionary drivers underlying clade evolution arose from the ecological theory, in the second half of the 20th century, notably coming from the formalisation of the concept of ecological niche.

Concept of 'competitive exclusion' first developed in 1936 with the formalisation of the Lotka-Volterra model, and experiments using *Paramecium* (see Diniz-Filho (2023)).

Concept of 'character displacement' Pfenning & Pfenning (2009)

2.2.2.3 Traits

The idea that phenotypical traits can influence diversification has been around since the formulation of the concept of adaptive radiation, where the emergence of a phenotypical feature ('key innovation') confers to a clade the ability ("ecological opportunity") to radiate within a previously unexploited portion of the environment (see Simpson 1944). Traits may also mediate biotic interactions.

2.3 Macro-eco-evolutionary understanding of the spatial organisation of life: insights from the Latitudinal Biodiversity Gradient

The uneven species richness across the globe that we outlined in Section 1.1, and notably, the striking pattern of increasing species richness with decreasing latitude famously known as the ‘Latitudinal Biodiversity Gradient’ (LBG), can be understood under a macroevolutionary perspective involving ecological processes. Several hypotheses (*reviewed in Pontarp et al. 2019*) have been proposed to explain such a pattern, some of which imply the interplay between diversification rates and tropical environmental conditions (*Tietje et al. 2022*).

In the footsteps of Stebbins (1974), the ‘cradle’ hypothesis posits that tropical lineages would have larger speciation rates than extra-tropical lineages. Higher speciation rates for tropical lineages may be predicted from tropic’s greater environmental heterogeneity (*Stein et al. 2014*), increasing available niche space. Conversely, one can expect speciation rates to be lower outside the tropics, where climate seasonality is more pronounced. In addition, the ‘Metabolic Theory of Ecology’ (*Brown et al. 2004*) can also predict that tropical lineages would have larger speciation rates. According to the latter, the higher energy availability for the tropics (higher temperature) would locally increase tropical species’ metabolic rate (*Allen et al. 2002*), yielding larger mutation, and thus, speciation rates (*Allen et al. 2006; Lanfear et al. 2010*). Also, as summarised by *Mittelbach et al. (2007)*, tropical regions’ greater species richness has been suggested to enhance the importance of biotic interactions (*Schemske et al. 2009*). Consequently, since the ‘Red Queen’ hypothesis (Section 2.2.2) postulates that biotic interactions would enhance speciation, this would establish a positive feedback loop within the tropics, where increasing diversity would yield increasing speciation rates. This ‘cradle’ hypothesis is often contrasted with the ‘museum’ hypothesis, instead placing the emphasis on the fact that tropical lineages would have lower extinction rates. The ‘museum’ hypothesis may be a consequence of the ‘tropical stability’ hypothesis, assuming that the tropics have remained climatically stable across geological times, and that this stability would have made tropical lineages less prone to extinction (*Wallace 1878*). In addition, it might be expected that the aforementioned environmental heterogeneity offered by the tropics could buffer tropical lineages against eventual environmental changes.

A consensus emerged from late 2000s–2010s studies testing the extent to which diversification rates could explain the observed LBG, where the tropics may actually both act as a cradle and a museum of diversity – some works also emphasised the role of tropical-to-extratropical dispersal as a side cause behind the GABI (*e.g., Jablonski et al. 2006; Marshall 2006; Mittelbach et al. 2007; Moreau & Bell 2013; Rolland et al. 2014*). This view has been contrastingly supported in the recent literature. Fossil-based analyses of the South American Mammal fossil record supported this view, with tropical taxa showing elevated speciation rates and reduced extinction rates (*Buffan et al. 2025*). Nevertheless, joint analyses of fossil and phylogenetic data for several tetrapod clades, using cutting-edge methodological frameworks, provided support for elevated speciation rate outside the tropics as a likely mechanism for the LBG, thus instead favouring the ‘museum’ hypothesis (*Meseguer & Condamine 2020; Quintero et al. 2023*). Conversely, a recent study focusing on a massive sample of the seed plant diversity emphasised that no causal relationship could be established between diversification rates and species richness (*Tietje et al. 2022*). Given these elements, it seems like the study model, the primary source of data (*e.g., fossils vs. dated phylogenies*) and the methodology used to estimate diversification rates (see Section 3.1) affects upstream interpretations of the interplay between diversification rates and spatial patterns of species richness.

However, the notions of ‘cradle’ and ‘museum’ of biodiversity have recently been challenged by

Vasconcelos *et al.* (2022), emphasising a number of limitations for the way they are often considered in the literature. Among them, as several factors driving speciation, such as climatic stability/heterogeneity, may also drive extinction, they argue that it would sometimes make more sense to consider turnover rates (see [Glossary](#)) rather than speciation and extinction rates individually. Also, they point out that extant centres of biodiversity with a great concentration of ‘basal’ lineages are not necessarily centres of origin, thus challenging both the way previous works actually interpreted the notion of museum and cradle of diversity. The latter nuance echoes previous works of the same authors emphasising the need to include hidden states in models of character evolution, marking a turning point in macroevolution (*e.g.*, [Beaulieu & O’Meara 2016](#)).

Alternative explanations of the LBG not involving diversification rates have also been proposed. These either refer to species’ ecological limits (*e.g.*, carrying capacity, size of the available area, palaeogeography) or time left for species to diversify (assuming tropical stability would have left more time for species to diversify within than outside the tropics, and attested by fossil and phylogenetic evidence of lineage age distribution in and outside the tropics) ([Pontarp *et al.* 2019](#)). As stressed by [Mittelbach *et al.* \(2007\)](#), these three classes of hypotheses (diversification-, ecological- or time-dependent) are not necessarily mutually exclusive. Conversely, a fourth set of hypotheses emphasised the likely implication of pure contingency in shaping the LBG, notably through the use of null models ([Arita & Vázquez-Domínguez 2008](#)). Recently insights from eco-evolutionary simulations (see [Section 3.2](#)) evidenced that null models – where the modelled eco-evolutionary processes do not rely on any environmental constraint – could indeed match empirical LBG patterns, although generally worse at explaining empirical distribution data than model incorporating mechanistic constraints on species, function of their biotic and abiotic environment ([Hagen *et al.* 2021](#)).

3 Analytical tools to study large-scale biodiversity patterns

Means to study diversification

3.1 Macroevolutionary approaches to study how clades thrive and go extinct

Review from Benton ([2015](#))

3.1.1 Phylogeny-based

3.1.1.1 Overview

What is a phylogeny?

Phylogeny-based methods

3.1.1.2 Limitations

3.1.2 Fossil occurrence-based

3.1.2.1 Overview

DivDyn, PyRate

3.1.2.2 Limitations

3.1.3 Towards integrating multiple sources of data

TED

3.2 Integrating ecological processes with evolutionary forces: mechanistic eco-evolutionary simulations

4 A strange biodiversity on a strange land : the case of mammals in South America

South America has attracted the interest of natural scientists since the times of Humboldt, Darwin or Wallace.

In 1876, Wallace wrote that the Neotropical region, encompassing South America, Caribbean Islands and tropical North America “possesses more peculiar families of vertebrates and genera of birds and mammalia than any other region” ([Wallace 1876](#)).

Introduce Wallaces’ zoogeographical subregions for the Neotropics (Mexican, Brazilian, Antillean, Chilean)?

4.1 South America across the Cenozoic: geographical and environmental context

Start from SA ecoregions in the present day (plot map, and overlay with diversity of clades).

Palaeoenvironmental evolution: summarise & update Hoorn *et al.* ([2010](#))

Continental island framework

4.1.1 Amazonia: a unique and dynamic tropical ecoregion

4.1.2 The rise of the Andes : emergence of one of the largest and highest mountain chains, and its consequences on biodiversity

4.1.3 Intermediate flooding events

4.1.4 South America as a playground for evolutionary biologists

4.2 ‘Splendid isolation’: the captivating history of South American mammals across the Cenozoic

“the southern tree of life, its own tree and independent of the one born and raised in the opposite hemisphere” (Francisco Moreno, El Museo de La Plata: Rápida ojeada sobre su fundación y desarrollo. Revista del Museo de La Plata 1 (1890–1891): 27–55)

4.2.1 History of mammal palaeontology in South America: a brief overview

Some of the historical elements developed here were taken from Manias (2023).

South American mammal palaeontology sadly started as a colonisation by-product, as European explorators plundered Patagonian soils, intrigued by skeletal remains trapped in the rocks. In the late 18th century, some of the fossils brought back to Europe provoked curiosity among the leading figures of the emerging field of comparative anatomy. As a result, in 1796, Georges Cuvier described the famous *Megatherium americanum*, and proposed this six-meter-beast to be a sloth. A few decades later, in 1836, Charles Darwin, back from his five-years journey on board the HMS Beagle, brought back to England a wealth of fossil material (in particular from the region of Punta Alta, Patagonia, Figure 4.1). These fossil remains were given to the renowned palaeontologist Richard Owen, who described, in 1837, many emblematic fossil mammals from South America, including the car-sized armadillo *Glyptodon*, the giant sloth *Mylodon*, the giant camel-like native ungulate *Macrauchenia* or the rhino-sized native ungulate *Toxodon*.

The second half of the 19th century marked the beginning of South American mammal palaeontology made by scientists from South America, or more precisely, from the Republic of Argentina. At that time, natural sciences, as well as any other field contributing to sharpening knowledge on the country's own territory, were envisaged as nation-building, and the Southern part of the country (Pampas and Patagonia provinces) had long been recognised to overflow with fossil remains of extinct animals. Consequently, important efforts were devoted to collect and shed light on the country's archaeological and palaeontological patrimony, centralised within national museums, such as the "Museo Nacional de La Plata", headed by the prominent explorer and natural scientist Franciso Moreno in the 1870s.

South American mammal palaeontology owes a lot to the labour of the patriotic Argentinian palaeontologist Florentino Ameghino. Relying on fossil material mostly collected during field expeditions led by his brother Carlos, Florentino Ameghino produced a remarkable quantity of systematic palaeontological work, and his written productions are regarded as essential building-blocks for South American palaeomammalogy. These are probably best exemplified by his foundational monographies, such as "*Contribución al conocimiento de los mamíferos fósiles de la República Argentina*" (Ameghino 1889), where several hundreds of taxa from the Argentinian fossil record are described. Argentina's fossil richness, together with the aforementioned efforts to enlighten its natural patrimony, made that South American palaeomammalogy long remained essentially constrained to the country, hence the striking spatial clustering of the fossil occurrences in the Southern cone depicted on Figure 4.1. As recalled by Simpson (1940), in the first half of the 20th century, knowledge of Cenozoic mammal remains outside Argentina was very scarce. Many of the most significant discoveries of non-Argentinian mammal-bearing fossil localities (as those highlighted on Figure 4.1) have indeed occurred throughout the last 80 years.

In the 1930s, the young U.S. vertebrate palaeontologist George G. Simpson, later to become the renowned architect of the Modern Synthesis of Evolution (Futuyama 2015), carried out his first expeditions in South America, intrigued by the continent's peculiar fossils. Simpson further developed a strong interest towards understanding the evolution of South American mammals, that animated him all over his career (e.g., Simpson 1940, 1950, 1967; 1970). In 1980, he produced a cornerstone synthesis of South American mammalian evolution across the Cenozoic, that is still used as a reference today (Simpson 1980). In his book, he proposed that the tempo of South American mammalian evolution could be summarised by three temporally-distinct 'faunal strata', that will be briefly outlined in the subsequent sections (Figure 4.2).




Figure 4.1: Distribution of the South American terrestrial mammal fossil record across geological epochs (Holocene excluded), on top of the continent's present-day topography (Boschman 2021). Several keystone mammal-bearing localities have been highlighted.

4.2.2 First stratum: endemic radiations (Late Cretaceous–middle Eocene)

Simpson's first stratum characterises the early phase of South American mammal evolution in the aftermath of the continent's isolation (Figure 4.2). It is thought to have lasted from the Late Cretaceous/early Palaeocene to the late middle Eocene. Mammal at that time are often referred to as the continent's 'original inhabitants'. These would include endemic metatherians (Figure 4.3), native ungulates (Figure 4.4), xenarthrans (Figure 4.5), as well as a few remnant non-therian lineages (*e.g.*, Gondwanatheria).

The oldest Cenozoic South American land mammalian assemblages date back to the early Palaeocene. These have been recovered from the Punta Peligro (Argentina) and Tiupampa (Bolivia) localities (Figure 4.1), each leading to the recognition of the oldest intervals of the South American Land Mammal Age (SALMA) scale, namely, the Peligran and the Tiupampan (Figure 4.2). Biochronological evidence suggests the Tiupampan SALMA to be older than the Peligran SALMA, and corroborate an age interval of 64.5–64 Ma for the Tiupampan fauna (Gelfo *et al.* 2009). An additional occurrence of pre-Tiupampan therian mammal was recovered from the Grenier Farm locality (Figure 4.1), and was assigned to the extinct metatherian order Polydolopimorphia (Goin *et al.* 2006).

The faunal composition of the Tiupampan and Peligran assemblages delivered precious insights

into the early Cenozoic history of South American mammals. The great metatherian diversity (Figure 4.3), together with the presence of representatives from several ungulate-related lineages typically found in the Northern hemisphere (formerly grouped into the ‘condylarths’), suggested that South American mammalian assemblages received significant faunal supply from North America between the late Cretaceous and the early Palaeocene (Pascual 2006). Geological evidence corroborated the possibility for intermediate episodic connections between the two continents in that interval (Pascual 2006), possibly also involving the Caribbean islands (Viñola-López *et al.* 2022). As an echo to the name ‘Great American Biotic Interchange’ (see Section 4.2.4 and Chapter 3), this keystone biotic exchange for the architecture of Simpson’s first stratum was thus termed ‘First American Biotic Interchange’ (FABI) (Goin *et al.* 2012a).

Along with these therian lineages of North American affinity, non-therian lineages persisted in South America at least until the mid-Palaeogene, thus well after the Cretaceous–Palaeogene boundary (Goin *et al.* 2012b). The most extreme example of non-therian lineage persistence known to date is probably best exemplified by the enigmatic *Necrolestes*, from the Santa Cruz Formation (early Miocene, Santacrucian SALMA, see Figure 4.2), identified as a remnant of the clade Meridiolestida, a mammal group that thrived from the Late Jurassic up until the Late Cretaceous (Chimento *et al.* 2012; Rougier *et al.* 2012).

Younger assemblages characteristic of Simpson’s first stratum notably include the São José de Itaboraí (often abbreviated Itaboraí) fauna (Figure 4.1), one of the most diverse in the Palaeogene of South America. It led to the Itaboraian SALMA, approximately dating back to 53 Ma (Figure 4.2). The Itaboraian fauna is still dominated by metatherians (Goin *et al.* 2016). It also revealed that all the higher South American Native Ungulate (SANU) lineages were clearly differentiated at that time (Figure 4.4). More importantly, the Itaboraian fauna yielded the oldest known occurrence of armadillo (Xenarthra: Cingulata) (Bergqvist *et al.* 2004), thus documenting the appearance of Xenarthra in the fossil record (Figure 4.5). The late appearance of xenarthrans in the fossil record remains an enigma for South American mammal palaeontology, and has been referred to as the ‘xenarthran mystery’ (Goin *et al.* 2016). Indeed, as suggested by their name meaning “weird articulations”, xenarthrans, today represented by sloths, anteaters and armadillos, share very derived morphological characters that are not found in any other known eutherian mammal lineage. Molecular evidence suggests that they form a clade with the african group Afrotheria, called ‘Atlantogenata’, literally meaning “generated by the Atlantic ocean”, and thus, would have diverged from their sister-group following the break-up of Gondwana, some 120 Ma (Waddell *et al.* 1999). Why armadillos only appear in the fossil record *ca.* 53 Ma, and sloths in the Late Oligocene (Section 4.2.3), whereas their divergence with other South American eutherians largely predates the earliest known mammal assemblages found on the continent thus remains an open question.

Younger assemblages of the first stratum progressively record a turnover in the relative proportion of metatherian vs. eutherian mammals in the communities, with an increasing dominance of eutherians, that several authors related to the gradual Eocene climate cooling from *ca.* 50 Ma onwards (Croft 2016; Goin *et al.* 2016; Westerhold *et al.* 2020). Later in the Eocene, land mammal assemblages will know go through a series of gradual changes that will progressively mark the transition with the subsequent faunal stratum, detailed thereafter.




Figure 4.2: Integrative view of the Cenozoic of South America from the land mammal perspective. The South American Land Mammal Age (SALMA) scale used all over this thesis is shown, along with the corresponding Geological Time Scale (GTS) epochs, from the [International Chronostratigraphic Chart](#) (v2024/12) (Holocene not depicted). Boundaries between SALMAs were defined on the basis of the most up-to-date literature (*e.g.*, [Bellinzoni *et al.* 2025](#); [Dunn *et al.* 2013](#); [Gelfo *et al.* 2009](#); [Woodburne *et al.* 2014](#)), but note that many intervals lack absolute temporal calibration. Black spaces between SALMAs represent temporal gaps. Alongside are depicted several key biotic or abiotic upheaval in the Cenozoic history of South American mammals, as well as Simpson's three faunal strata ([Simpson 1980](#)). Silhouettes are from Phylopic (credit: Sarah Werning and Kai R. Caspar).

4.2.3 Second stratum: Trans-atlantic migrations and faunal ‘modernisation’ (late Eocene–Late Miocene)

Most specialists agree that major re-organisations occurred within South American mammal assemblages near the Eocene–Oligocene boundary (that will be more thoroughly covered in [Chapter 1](#)), marking the onset of Simpson’s second faunal stratum (Figure 4.2). The distinction between the second and the first strata lies in a three-part faunal ‘modernisation’ (Flynn & Wyss 1998): (1) emergence of morphological innovations among “original inhabitants”, (2) extinction of basal, ‘archaic’ lineages and (3) higher-level taxonomic origination within “original inhabitants”, alongside the arrival and radiation of hystricognathous rodents (Rodentia: Hystricognathi: Caviomorpha) and anthropoid primates (Primates: Anthropoidea) in South America (Figure 4.2).

The most notable morphological innovation that could be mentioned is probably the increase in cheek teeth crown height (*i.e.*, hypsodonty), occurring independently among several (at least six) groups of native ungulates (particularly notoungulates) from the middle Eocene onwards (Croft *et al.* 2020). The evolution towards hypsodonty in South America is particularly interesting as hypsodont cheek teeth started to be recorded 20 million years before the first reports of hypsodonty outside South America (García-López *et al.* 2019). As they diversified, freshly-arrived caviomorph rodents (see later in this section) also rapidly acquired hypsodont cheek teeth, again, way before any other place in the world (late Oligocene–Early Miocene), as exemplified by the superfamily Chinchilloidea – see [Chapter 2](#). The Eocene emergence of hypsodont cheek teeth and their Oligocene spread across lineages has long puzzled South American mammal palaeontologists (*e.g.*, Billet *et al.* 2009; Croft *et al.* 2003; Reguero *et al.* 2008, 2010). Hypsodont cheek teeth are indeed often considered as an indicator of an abrasive diet, typically, incorporating grasses, and thus, have often been related to landscape opening (Stebbins 1981). However, palaeoenvironmental evidence failed to support any concomitant grassland spreading in the regions where the first hypsodont lineages were recorded (*e.g.*, Kohn *et al.* 2004, 2015; Strömberg *et al.* 2013). Now, a consensus emerged in the community, as several lines of evidence instead suggested that the increase in cheek teeth crown height may have instead resulted from grit or volcanic dust consumption (Armella & Croft 2026; Damuth & Janis 2011; Jardine *et al.* 2012; Selkin *et al.* 2015).

Besides, by analysing two early Oligocene metatherian communities from Southern South America (one of Tinguirican age, the other being the La Cantera fauna, see Figure 4.2), Goin (2010) reported an important taxonomic turnover occurring among metatherians in the few million years that followed the Eocene–Oligocene transition (EOT). This geological boundary, that will be further explored in [Chapter 1](#), has been identified as the most intense Cenozoic climatic disruption, as global temperature cooled down by several degrees in a few hundred thousands of years (Coxall & Pearson 2007; Liu *et al.* 2009; Westerhold *et al.* 2020). The EOT was related to a major faunal turnover among mammalian communities from the Northern hemisphere (Meng & McKenna 1998; Stehlin 1909; Stucky 2014). As an echo to the french expression ‘Grande Coupure’ (Stehlin 1909) used to characterise the mammalian turnover that occurred among European mammal assemblages, Goin (2010) thus derived the expression ‘Bisagra Patagónica’ (*Patagonian hinge*) to qualify the faunal turnover that they characterised.

Next, as we briefly evoked, Simpson’s second stratum is also characterised by the abrupt apparition of hystricognathous rodents and anthropoid primates in the South American fossil record (Antoine *et al.* 2012; Bond *et al.* 2015; Marivaux *et al.* 2023; Seiffert *et al.* 2020). Explaining such an emergence has been one of the most important challenges for South American mammal palaeontology, given that, contray to xenarthrans (Section 4.2.2), it made no doubt that these groups were

the result of a much more recent faunal supply (Simpson 1950). After several decades of intense debate, combined morphological (Hoffstetter 1972) and molecular (*e.g.*, Poux *et al.* 2006) evidence provided strong support for the “rafting hypothesis”, in which both groups would descend from Afro-Asian ancestors that would have crossed the Atlantic in the Palaeogene (Figure 4.2).

Despite its wide acceptance, the timing of this fascinating trans-Atlantic journey remains under debate. Seiffert *et al.* (2020) proposed that the global sea-water drop that followed the EOT, by reducing the distance between Africa and South America, could have increased the likelihood for a trans-oceanic dispersal (assuming the latter to be inversely proportional to the distance between the source and the receiving areas). Such a hypothesis finds support in that, according to a study conducted by the same authors, the oldest fossil evidence for rodents and primates in South America cannot be older than the early Oligocene (Campbell *et al.* 2021). Conversely, Marivaux *et al.* (2023) proposed that this trans-Atlantic dispersal may have instead occurred 5 million years earlier, near the Middle Eocene Climate Optimum (MECO) (Bohaty *et al.* 2009). This interval may have been accompanied by intense meteorological episodes in the tropics (Mulch *et al.* 2015), increasing the likelihood of riverbank break-ups, and thus, the formation of floating islands serving as a means for the large-scale dispersal of small-sized mammals, such as rodents or primates (Houle 1998). This hypothesis more adequately explains the high taxonomic diversity displayed by several known early Oligocene rodent communities (*e.g.*, Antoine *et al.* 2021; Bertrand *et al.* 2012 *though some assemblages from the former may be of earliest Eocene age*), that would be hard to achieve in the few hundred thousand years separating these assemblages from the EOT. Note that, given the paucity of the primate fossil record in the Palaeogene of South America – only four specimens assigned to three distantly-related lineages that notably, and contrary to rodents, are not recognised as stem representatives of the extant South American primate radiation, namely, Platyrrhini (Marivaux *et al.* 2023) – it is currently impossible to apply such a limitation to the primate scenario proposed by Seiffert *et al.* (2020).

Simpson’s second stratum persisted all the way through the Oligocene and the Miocene. Keystone late Oligocene localities, such as Salla (Bolivia) or the Deseado massif (Argentina) (Figure 4.1) provided unprecedented insight into South American mammal diversity at that time. In particular, they helped identify the first records of sloths (Pujos & De Iuliis 2007) as well as to better constraint the diversification of Notoungulata, Litopterna, Astrapoheria and the extinction of Pyrotheria near the Oligocene–Miocene boundary (Croft *et al.* 2020) (Figure 4.4). Then, from the early Miocene, sampling gets increasingly less temporally discontinuous (Figure 4.2), thus better constraining our knowledge of the evolution of mammals on the continent. In addition, molecular and fossil data agree that most of the crown extant South American mammal families that survived until the present diverged in the Miocene (Álvarez-Carretero *et al.* 2022). Thus, mammal diversity from the Late Oligocene to the Late Miocene was flourishing, and many invaluable sites or geological formations (*e.g.*, Santa Cruz Formation, La Venta, Quebrada Honda, ... see Figure 4.1 as well as Croft (2016)) yielded precious discoveries attesting that statement.




Figure 4.3: Order-level persistence of South American metatherians in the fossil record. Fading colours mean that the group has no known fossil record for the period, but should already exist. Photos were taken from iNaturalist (credit: Jorge Brito, Eduardo Silva Rodriguez, Thiago Chagas). *Wamradolops telloi* cranium comes from Stutz *et al.* (2026), reproduced with permission. The picture of the *Thylacosmilus atrox* cranium was taken by Claire Houck

4.2.4 Third stratum: Inter-american exchange, massive faunal disruptions, and establishment of the extant mammal biodiversity (latest Miocene–Present)

The flourishing diversity that was characteristic of Simpson’s second stratum was massively disrupted with the late Neogene onset of land mammal exchange across the Americas, known as the Great American Biotic Interchange (Webb 1985) (Figure 4.2).

Tropical North America: “the portion of the North American continent which belongs zoologically to South America” (Wallace 1876).

4.3 Challenges for South American mammal palaeontology

Despite being unique and invaluable, several challenges arise as we intend to work with the South American mammal fossil record, in addition to those previously mentioned arising while using fossils (Section 1.4.2).

Flynn & Wyss (1998) already pointed the need to fill the gap in tropical fossil record

Striking unevenness of the fossil record Ugarte *et al.* (2025). This unevenness is spatial (Figure 4.1) and temporal (Figure 4.2). South American mammal fossil record paradox: there is almost an inverse gradient in extant mammal diversity Figure 1.2 and the abundance of fossil remains




Figure 4.4: Higher-level persistence of South American Native Ungulates (SANUs) in the fossil record, adapted from Croft *et al.* (2020). Fading colours depict uncertainty in the age of earliest fossil. Silhouettes are from Phylopic (credit: Julián Bayona).

across the continent, mostly concentrated within Southern South American plains Figure 4.1.

Difficulty to sample, consequently, and particularly under low latitudes (tropical), fossil record is sparse

Colonisation palaeontology.

The South American Land Mammal Ages (SALMA) system: between gaps and uncertainties




Figure 4.5: Higher-level persistence of xenarthrans in the fossil record. Fading colours with question marks depict the ‘xenarthran mystery’, *i.e.*, that the group in question has no known fossil record for the period, but should already exist. Divergence ages between the main lineages were taken from Tejada *et al.* (2024). Vermilingua and Folivora altogether form the order Pilosa. Silhouettes are from Phylopic (credit: Julián Bayona and Sarah Werning).

5 Objectives of this doctoral work

5.1 Deciphering the dynamics of diversification of South American mammals and their likely drivers across the Cenozoic

Can we explain the emergence of key phenotypical features (hypsodonty, body size)?

5.1.1 Elucidating the impact of the Eocene–Oligocene Transition on the macroevolutionary dynamics of South American mammals

5.1.2 Did the evolutionary pathways towards increasing in body size and hypsodonty impact the diversification of chinchilloid rodents?

5.2 Assessing the role of palaeoclimate in the emergence of the asymmetry behind the Great American Biotic Interchange

Chapter 1. Deciphering the trends and drivers of South American mammal large-scale evolution across the Cenozoic

Chapter 2.1. The fate of South America's endemic mammalian fauna in response to the most dramatic Cenozoic climate disruption

Chapter 2.2. Continent-scale insights into the last 23 million years of South American mammalian evolution

Chapter 2. A matter of size: what underpinned the rise and demise of the largest rodents that ever lived?

Chapter 3. Palaeoclimate alone fails to explain the asymmetry of the Great American Biotic Interchange

Discussion

Limitations

Lack of time-continuous proxy for precipitations.

Lack of well-constrained phylogenetic framework, as many of the endemic Cenozoic South American mammal fauna has gone extinct

Terminology, definitions and other macroevolutionary enigmas

Defining a species?

Inferring diversity/diversification patterns from the fossil record: Signor & Lipps effect Signor & Lipps ([1982](#))

Perspectives

An almost closed system: founding migrations in the Palaeogene history of South American mammals

Hystricognath rodents & Platyrrhine primates. Introduce the debate. We lack fossil evidence for the earliest comers, oldest occurrences in Western Amazonia whereas most likely come from Afro-Asian ancestors, so should have arrived by the East. Polyphyletic/several crossing events? Summarised in Marivaux *et al.* ([2023](#))

A recent quantitative study suggested that the diversification of rodents and primates negatively affected the diversity of the endemic groups ([Pino *et al.* 2025](#)).

There is still some work to do to accurately elucidate continent-scale macroevolutionary dynamics of Caviomorph rodents (others than Chinchilloids, a contribution of this thesis): set the scene for GIGARAT

Sustainable late Cenozoic connection to North America: what drove the asymmetry of the GABI?

Set the scene for IMBALANCE

How about neutral theory?

Future directions towards an improved understanding of the emergence and dynamics of South American ecosystems

DIACHRONANDES, PHENOMENA

Concluding remarks

Glossary

This section intends to provide simple definitions to key notions and concepts to which this thesis refers. Note that some are complex and could be the subject of entire research works. Thus, these definitions do not aim to be as exhaustive as possible, but rather, to improve the reader's understanding of this manuscript.

Adaptive radiation. Rapid *diversification* of a group taking advantage of a previously unexploited ecological dimension of the environment, generally spurred by a *key innovation*. This concept has been proposed by Simpson (1944).

Angiosperm Terrestrial Revolution (ATR). Suite of major global-scale ecosystem reshaping that followed the radiation of flowering plants (angiosperms) in the Early Cretaceous (*ca.* 140 Ma). The term was introduced by Benton *et al.* (2022).

Biogeography. Discipline aiming to describe and understand the spatial distribution of species.

Clade. Given a multi-species sample, a clade is an exhaustive subsample of all species sharing a common ancestry. Clades are referred to as *monophyletic*.

Diversification. The diversification process summarises the net evolutionary changes in a *clade* across time, that is, the gains (modelled by the *speciation* process) minus the losses (modelled by the *extinction* process). As such, it represents the process generating clade's diversity. The intensity of the diversification process (called diversification rate, often noted $d(t)$ for a given time t) is thus the difference between the rates of speciation and extinction, leading to $d(t) = \lambda(t) - \mu(t)$.

Ecological niche. The concept of ecological niche may be understood in different ways. In macroecology, the niche is more often viewed from a perspective of the species' requirements, therefore following the framework proposed by Grinnell (1917) and later Hutchinson (1957). Considering a landscape made up with n ecological variables (*e.g.*, climate, topography, abundance of other species, abundance of trophic resources, ...), the ecological niche of a given species theoretically represents the portion of the landscape where the species occurs. As such, it can be viewed as an n -hyperpolygon encompassing the species' tolerance range following each variable (*e.g.*, 15–25°C, 0–1000m, ...). Other definitions exist, such as the Eltonian niche Elton (1927), more centered on species' impact on the environment.

Extinction. Process of disappearance of all representatives of a given lineage. In *macroevolutionary modelling*, extinction rates (expressed in event per unit of time) are often noted $\mu(t)$.

Latitudinal Biodiversity Gradient (LBG). Pattern of increasing species diversity with decreasing latitude (*i.e.*, within a given unit of the globe's surface, the number of species within it increases as we get closer to the equator), found in a wide range of group.

Speciation. Origination of a new species from an ancestral lineage. There are several known types of speciation. Speciation is called allopatric when a physical barrier (*e.g.*, a mountain, a river) subdivides an ancestral population into several sub-populations, thus interrupting gene flow and driving to the split in as many species as initial subsets by divergence accumulation. Conversely,

a the barrier to gene flow arises within a population where individuals remain in connexion (*e.g.*, behavioural barrier), people refer to *sympatric* speciation. When speciation arises from a founder effect, *i.e.*, colonisation of a new area (*e.g.*, an island), one refers to *peripatry*. A particular case of peripatry, when the colonised area is adjacent to the range of the initial species, is called *parapatry*. Given these several modes, modelling speciation can be done in different ways, defined in this glossary (*cladogenesis*, *anagenesis*, *budding*). In *macroevolutionary modelling*, speciation rates through time, expressed in number of events per unit of time, are often noted $\lambda(t)$

Trait.

Literature cited

- Allen, A.P., Brown, J.H. & Gillooly, J.F. (2002). [Global biodiversity, biochemical kinetics, and the energetic-equivalence rule](#). *Science*, 297, 1545–1548.
- Allen, A.P., Gillooly, J.F., Savage, V.M. & Brown, J.H. (2006). [Kinetic effects of temperature on rates of genetic divergence and speciation](#). *Proceedings of the National Academy of Sciences*, 103, 9130–9135.
- Álvarez-Carretero, S., Tamuri, A.U., Battini, M., Nascimento, F.F., Carlisle, E., Asher, R.J., *et al.* (2022). [A species-level timeline of mammal evolution integrating phylogenomic data](#). *Nature*, 602, 263–267.
- Ameghino, F. (1889). *Contribución al conocimiento de los mamíferos fósiles de la república argentina*. Buenos Aires, Argentina.
- Antoine, P.-O., Marivaux, L., Croft, D.A., Billet, G., Ganerød, M., Jaramillo, C., *et al.* (2012). [Middle eocene rodents from peruvian amazonia reveal the pattern and timing of caviomorph origins and biogeography](#). *Proceedings of the Royal Society B: Biological Sciences*, 279, 1319–1326.
- Antoine, P.-O., Yans, J., Castillo, A.A., Stutz, N., Abello, M.A., Adnet, S., *et al.* (2021). [Biotic community and landscape changes around the Eocene–Oligocene transition at Shapaja, Peruvian Amazonia: Regional or global drivers?](#) *Global and Planetary Change*, 202, 103512.
- Arita, H.T. & Vázquez-Domínguez, E. (2008). [The tropics: cradle, museum or casino? A dynamic null model for latitudinal gradients of species diversity](#). *Ecology Letters*, 11, 653–663.
- Armella, M.A. & Croft, D.A. (2026). [Pass on the grass? The unexpected “last supper” of hypselodont pachyrukhinae \(notoungulata, mammalia\) from the late neogene of northwestern argentina](#). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 687, 113585.
- Barnosky, A.D. (2001). [Distinguishing the effects of the red queen and court jester on miocene mammal evolution in the northern rocky mountains](#). *Journal of Vertebrate Paleontology*, 21, 172–185.
- Beaulieu, J.M. & O’Meara, B.C. (2016). [Detecting hidden diversification shifts in models of trait-dependent speciation and extinction](#). *Systematic Biology*, 65, 583–601.
- Bellinzoni, J., Bonini, R., García-Morato, S., Gómez, G.N., Steffan, P., Marín-Monfort, M.D., *et al.* (2025). [New mammal assemblage from last interglacial in argentine pampas: Debating biostratigraphic and biochronological reliability](#). *Quaternary Science Reviews*, 367, 109511.
- Benton, M.J. (2015). [Exploring macroevolution using modern and fossil data](#). *Proceedings of the Royal Society B: Biological Sciences*, 282, 20150569.
- Benton, M.J., Wilf, P. & Sauquet, H. (2022). [The Angiosperm Terrestrial Revolution and the origins of modern biodiversity](#). *New Phytologist*, 233, 2017–2035.
- Bergqvist, L.P., Abrantes, É.A.L. & Avilla, L.S. (2004). [The Xenarthra \(Mammalia\) of São José de Itaboraí Basin \(upper Paleocene, Itaboraian\), Rio de Janeiro, Brazil](#). *Geodiversitas*, 26, 323–337.
- Berry, A. & Browne, J. (2008). [The other beetle-hunter](#). *Nature*, 453, 1188–1190.
- Bertrand, O.C., Flynn, J.J., Croft, D.A. & Wyss, A.R. (2012). [Two new taxa \(caviomorpha, rodentia\) from the early oligocene tinguiririca fauna \(chile\)](#). *American Museum Novitates*, 2012, 1–36.

- Billet, G., Blondel, C. & Muizon, C. de. (2009). [Dental microwear analysis of notoungulates \(mammalia\) from salla \(late oligocene, bolivia\) and discussion on their precocious hypsodonty](#). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 274, 114–124.
- Bohaty, S.M., Zachos, J.C., Florindo, F. & Delaney, M.L. (2009). [Coupled greenhouse warming and deep-sea acidification in the middle Eocene](#). *Paleoceanography*, 24.
- Bond, M., Tejedor, M.F., Campbell, K.E., Chornogubsky, L., Novo, N. & Goin, F. (2015). [Eocene primates of South America and the African origins of New World monkeys](#). *Nature*, 520, 538–541.
- Boschman, L.M. (2021). [Andean mountain building since the late cretaceous: A paleoelevation reconstruction](#). *Earth-Science Reviews*, 220, 103640.
- Brown, J.H., Gillooly, J.F., Allen, A.P., Savage, V.M. & West, G.B. (2004). [Toward a Metabolic Theory of Ecology](#). *Ecology*, 85, 1771–1789.
- Buffan, L., Condamine, F.L., Stutz, N.S., Pujos, F., Antoine, P.-O. & Marivaux, L. (2025). [The fate of south america's endemic mammalian fauna in response to the most dramatic cenozoic climate disruption](#). *Proceedings of the National Academy of Sciences*, 122, e2419520122.
- Burkill, I.H. (1943). [The biogeographic division of the indo-australian archipelago. 2. A history of the divisions which have been proposed](#). *Proceedings Linnean Society London*, 154, 127–138.
- Campbell, K.E., O'Sullivan, P.B., Fleagle, J.G., Vries, D. de & Seiffert, E.R. (2021). [An early oligocene age for the oldest known monkeys and rodents of south america](#). *Proceedings of the National Academy of Sciences*, 118, e2105956118.
- Chimento, N.R., Agnolin, F.L. & Novas, F.E. (2012). The Patagonian fossil mammal Necrolestes: a Neogene survivor of Dryolestidae. *Revista del Museo Argentino de Ciencias Naturales*.
- Close, R.A., Benson, R.B.J., Alroy, J., Carrano, M.T., Cleary, T.J., Dunne, E.M., *et al.* (2020). [The apparent exponential radiation of phanerozoic land vertebrates is an artefact of spatial sampling biases](#). *Proceedings of the Royal Society B: Biological Sciences*, 287, 20200372.
- Coxall, H.K. & Pearson, P.N. (2007). [The eocene–oligocene transition](#). In: (eds. Williams, M., Haywood, A.M., Gregory, F.J. & Schmidt, D.N.). Geological Society of London, pp. 351–387.
- Croft, D.A. (2016). *Horned armadillos and rafting monkeys: The fascinating fossil mammals of south america*. Indiana University Press.
- Croft, D.A., Croft, D.A., Bond, M., Flynn, J.J., Reguero, M. & Wyss, A.R. (2003). [Large archaeohyracids \(typotheria, notoungulata\) from central chile and patagonia : Including a revision of archaeotypotherium](#). Field Museum of Natural History, Chicago, Ill.
- Croft, D.A., Gelfo, J.N. & López, G.M. (2020). [Splendid innovation: The extinct south american native ungulates](#). *Annual Review of Earth and Planetary Sciences*, 48, 259–290.
- Cuvier, G. (1840). Discours sur les révolutions de la surface du globe.
- Damuth, J. & Janis, C.M. (2011). [On the relationship between hypsodonty and feeding ecology in ungulate mammals, and its utility in palaeoecology](#). *Biological Reviews*, 86, 733–758.
- Darwin, C. (1859). *On the origin of species by means of natural selection, or the preservation of favoured races in the struggle for life*. John Murray. London.
- Davis, A.M., Unmack, P.J., Vari, R.P. & Betancur-R., R. (2016). [Herbivory Promotes Dental Disparification and Macroevolutionary Dynamics in Grunters \(Teleostei: Terapontidae\), a Freshwater Adaptive Radiation](#). *The American Naturalist*, 187, 320–333.
- Diniz-Filho, J.A.F. (2023). [The macroecological perspective](#). Springer Cham.
- Dunn, R.E., Madden, R.H., Kohn, M.J., Schmitz, M.D., Strömberg, C.A.E., Carlini, A.A., *et al.* (2013). [A new chronology for middle eocene–early miocene south american land mammal ages](#). *GSA Bulletin*, 125, 539–555.
- Elton, C. (1927). *Animal ecology*. Sidgwick and Jackson. London.
- Flynn, J.J. & Wyss, A.R. (1998). [Recent advances in south american mammalian paleontology](#). *Trends in Ecology & Evolution*, 13, 449–454.

- Futuyma, D.J. (2015). [Can Modern Evolutionary Theory Explain Macroevolution?](#) In: (eds. Serrelli, E. & Gontier, N.). Springer International Publishing, Cham, pp. 29–85.
- García-López, D.A., Gelfo, J.N., Scanferla, A., Babot, J., Ferro, A. & Saade, L.S. (2019). [The pre-oligocene diversity of hypsodont typotherians \(mammalia, notoungulata\) in northwestern argentina.](#) *Ameghiniana*, 57, 117–131.
- Gearty, W. (2023). [Deeptime: Plotting tools for anyone working in deep time.](#) *R package version 1.0.1*.
- Gelfo, J.N., Goin, F.J., Woodburne, M.O. & Muizon, C.D. (2009). [Biochronological Relationships of the Earliest South American Paleogene Mammalian Faunas.](#) *Palaeontology*, 52, 251–269.
- Goin, F.J. (2010). [Goin FJ, et al. \(2010\) middle tertiary marsupials from central patagonia \(early oligocene of gran barranca\): Understanding south america’s grande coupure.](#) In: *The Paleontology of Gran Barranca*. Cambridge Univ Press, New York, 448 pp. Pp. 71-107.
- Goin, F.J., Gelfo, J.N., Chornogubsky, L., Woodburne, M.O. & Martin, T. (2012a). [3. Origins, radiations, and distribution of south american mammals: From greenhouse to icehouse worlds.](#) In: *Bones, clones, and biomes: The history and geography of recent neotropical mammals* (eds. Patterson, B.D. & Costa, L.P.). University of Chicago Press, pp. 20–50.
- Goin, F.J., Pascual, R., Tejedor, M.F., Gelfo, J.N., Woodburne, M.O., Case, J.A., et al. (2006). [The earliest tertiary therian mammal from south america.](#) *Journal of Vertebrate Paleontology*, 26, 505–510.
- Goin, F.J., Tejedor, M.F., Chornogubsky, L., López, G.M., Gelfo, J.N., Bond, M., et al. (2012b). [Persistence of a mesozoic, non-therian mammalian lineage \(gondwanatheria\) in the mid-paleogene of patagonia.](#) *Naturwissenschaften*, 99, 449–463.
- Goin, F.J., Woodburne, M.O., Zimicz, A.N., Martin, G.M. & Chornogubsky, L. (2016). [A brief history of south american metatherians: Evolutionary contexts and intercontinental dispersals.](#) Springer Dordrecht.
- Grinnell, J. (1917). [The niche-relationships of the california thrasher.](#) *The Auk*, 34, 427–433.
- Hagen, O., Flück, B., Fopp, F., Cabral, J.S., Hartig, F., Pontarp, M., et al. (2021). [gen3sis: A general engine for eco-evolutionary simulations of the processes that shape Earth’s biodiversity.](#) *PLOS Biology*, 19, e3001340.
- Hand, S.J., Maugoust, J., Beck, R.M.D. & Orliac, M.J. (2023). [A 50-million-year-old, three-dimensionally preserved bat skull supports an early origin for modern echolocation.](#) *Current Biology*, 33, 4624–4640.e21.
- Hautmann, M. (2020). [What is macroevolution?](#) *Palaeontology*, 63, 1–11.
- Hévin, N.M.-C. & Condamine, F.L. (2024). [Bayesian Total-Evidence Dating Reshapes the Age and Historical Biogeography of Parnassiinae.](#) *Bulletin of the Society of Systematic Biologists*, 3.
- Hillebrand, H. (2004). On the generality of the latitudinal diversity gradient. *The American Naturalist*, 163, 192211.
- Hoffstetter, R. (1972). [Relationships, Origins, and History of the Ceboid Monkeys and Caviomorph Rodents: A Modern Reinterpretation.](#) In: (eds. Dobzhansky, T., Hecht, M.K. & Steere, W.C.). Springer US, New York, NY, pp. 323–347.
- Holt, B.G., Lessard, J.-P., Borregaard, M.K., Fritz, S.A., Araújo, M.B., Dimitrov, D., et al. (2013). [An update of wallace’s zoogeographic regions of the world.](#) *Science*, 339, 74–78.
- Hoorn, C., Wesselingh, F.P., Steege, H. ter, Bermudez, M.A., Mora, A., Sevink, J., et al. (2010). [Amazonia through time: Andean uplift, climate change, landscape evolution, and biodiversity.](#) *Science*, 330, 927–931.
- Hortal, J., Bello, F. de, Diniz-Filho, J.A.F., Lewinsohn, T.M., Lobo, J.M. & Ladle, R.J. (2015). [Seven Shortfalls that Beset Large-Scale Knowledge of Biodiversity.](#) *Annual Review of Ecology, Evolution, and Systematics*, 46, 523–549.

- Houle, A. (1998). Floating islands: A mode of long-distance dispersal for small and medium- sized terrestrial vertebrates. *Diversity and Distributions*, 4, 201–216.
- Humboldt, A. von & Bonpland, A. (1807). *Essai sur la géographie des plantes; accompagne d'un tableau physique des régions équinoxiales*. Paris: Fr. Schoell.
- Hutchinson, G.E. (1957). Cold spring harb symp quant biol. pp. 415–427.
- Jablonski, D., Roy, K. & Valentine, J.W. (2006). Out of the tropics: Evolutionary dynamics of the latitudinal diversity gradient. *Science*, 314, 102–106.
- Jardine, P.E., Janis, C.M., Sahney, S. & Benton, M.J. (2012). Grit not grass: Concordant patterns of early origin of hypsodonty in great plains ungulates and glires. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 365–366, 1–10.
- Jones, L.A. (2022). *Sepkoski: Sepkoski's fossil marine animal genera compendium*.
- Kohn, M.J., Josef, J.A., Madden, R., Kay, R., Vucetich, G. & Carlini, A.A. (2004). Climate stability across the eocene-oligocene transition, southern argentina. *Geology*, 32, 621–624.
- Kohn, M.J., Strömberg, C.A.E., Madden, R.H., Dunn, R.E., Evans, S., Palacios, A., *et al.* (2015). Quasi-static eocene–oligocene climate in patagonia promotes slow faunal evolution and mid-cenozoic global cooling. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 435, 24–37.
- Lamarck, J.-B.P.A. de M. de. (1815). *Histoire naturelle des animaux sans vertèbres*. Paris.
- Lanfear, R., Ho, S.Y.W., Love, D. & Bromham, L. (2010). Mutation rate is linked to diversification in birds. *Proceedings of the National Academy of Sciences*, 107, 20423–20428.
- Liu, Z., Pagani, M., Zinniker, D., DeConto, R., Huber, M., Brinkhuis, H., *et al.* (2009). Global cooling during the eocene-oligocene climate transition. *Science*, 323, 1187–1190.
- Lydekker, R. (1896). *A geographical history of mammals*. Cambridge University Press.
- Manias, C. (2023). *The age of mammals: Nature, development, and paleontology in the long nineteenth century*. University of Pittsburgh Press.
- Marion, A.F.P., Condamine, F.L. & Guinot, G. (2024). Sequential trait evolution did not drive deep-time diversification in sharks. *Evolution*, qpae070.
- Marivaux, L., Negri, F.R., Antoine, P.-O., Stutz, N.S., Condamine, F.L., Kerber, L., *et al.* (2023). An eosimiid primate of south asian affinities in the paleogene of western amazonia and the origin of new world monkeys. *Proceedings of the National Academy of Sciences*, 120, e2301338120.
- Marshall, C.R. (2006). Fossil record reveals tropics as cradle and museum. *Science*, 314, 66–67.
- Mayr, G. (2004). Old world fossil record of modern-type hummingbirds. *Science*, 304, 861–864.
- Meng, J. & McKenna, M.C. (1998). Faunal turnovers of Palaeogene mammals from the Mongolian Plateau. *Nature*, 394, 364–367.
- Meseguer, A.S. & Condamine, F.L. (2020). Ancient tropical extinctions at high latitudes contributed to the latitudinal diversity gradient. *Evolution*, 74, 1966–1987.
- Mittelbach, G.G., Schemske, D.W., Cornell, H.V., Allen, A.P., Brown, J.M., Bush, M.B., *et al.* (2007). Evolution and the latitudinal diversity gradient: speciation, extinction and biogeography. *Ecology Letters*, 10, 315–331.
- Mora, C., Tittensor, D.P., Adl, S., Simpson, A.G.B. & Worm, B. (2011). How Many Species Are There on Earth and in the Ocean? *PLOS Biology*, 9, e1001127.
- Moreau, C.S. & Bell, C.D. (2013). Testing the museum versus cradle tropical biological diversity hypothesis: Phylogeny, diversification, and ancestral biogeographic range evolution of the ants. *Evolution*, 67, 2240–2257.
- Mulch, A., Chamberlain, C.P., Cosca, M.A., Teyssier, C., Methner, K., Hren, M.T., *et al.* (2015). Rapid change in high-elevation precipitation patterns of western North America during the Middle Eocene Climatic Optimum (MECO). *American Journal of Science*, 315, 317–336.
- Newell, N.D. (1952). Periodicity in invertebrate evolution. *Journal of Paleontology*, 26, 371–385.
- Nürk, N.M., Atchison, G.W. & Hughes, C.E. (2019). Island woodiness underpins accelerated dis-

- parification in plant radiations. *New Phytologist*, 224, 518–531.
- Pascual, R. (2006). Evolution and geography: the biogeographic history of South American land mammals. *Annals of the Missouri Botanical Garden*, 93, 209–230.
- Pfenning, K.S. & Pfenning, D.W. (2009). Character Displacement: Ecological And Reproductive Responses To A Common Evolutionary Problem. *The Quarterly Review of Biology*, 84.
- Phillips, J. (1860). *Life on the earth; its origin and succession*. MacMillan; co, Cambridge; London.
- Pino, K., Rodriguez-Serrano, E., Carrillo, J.D., Cooper, R.B. & Silvestro, D. (2025). Splendid isolation with migration: Diversity dynamics of South American mammals.
- Pontarp, M., Bunnefeld, L., Cabral, J.S., Etienne, R.S., Fritz, S.A., Gillespie, R., *et al.* (2019). The latitudinal diversity gradient: Novel understanding through mechanistic eco-evolutionary models. *Trends in Ecology & Evolution*, 34, 211–223.
- Poux, C., Chevret, P., Huchon, D., Jong, W.W. de & Douzery, E.J.P. (2006). Arrival and diversification of caviomorph rodents and platyrrhine primates in south america. *Systematic Biology*, 55, 228–244.
- Pujos, F. & De Iuliis, G. (2007). Late oligocene megatherioidea fauna (mammalia: Xenarthra) from salla-luribay (bolivia): New data on basal sloth radiation and cingulata-tardigrada split. *Journal of Vertebrate Paleontology*, 27, 132–144.
- Quintero, I., Landis, M.J., Jetz, W. & Morlon, H. (2023). The build-up of the present-day tropical diversity of tetrapods. *Proceedings of the National Academy of Sciences*, 120, e2220672120.
- Raup, D.M. (1966). Geometric analysis of shell coiling: General problems. *Journal of Paleontology*, 40, 1178–1190.
- Raup, D.M. (1972). Taxonomic diversity during the phanerozoic. *Science*, 177, 1065–1071.
- Raup, D.M. (1986). Biological extinction in earth history. *Science*, 231, 1528–1533.
- Raup, D.M. & Sepkoski, J.J. (1982). Mass extinctions in the marine fossil record. *Science*, 215, 1501–1503.
- Raven, P.H., Gereau, R.E., Phillipson, P.B., Chatelain, C., Jenkins, C.N. & Ulloa Ulloa, C. (2020). The distribution of biodiversity richness in the tropics. *Science Advances*, 6, eabc6228.
- Reguero, M.A., Candela, A.M. & Cassini, G.H. (2010). Hypsodonty and body size in rodent-like notoungulates. Cambridge University Press, pp. 358–367.
- Reguero, M.A., Croft, D.C., López, G.M. & Alonso, R.N. (2008). Eocene archaeohyracids (mammalia: Notoungulata: Hegetotheria) from the puna, northwest argentina. *Journal of South American Earth Sciences*, 26, 225–233.
- Rolland, J., Condamine, F.L., Jiguet, F. & Morlon, H. (2014). Faster Speciation and Reduced Extinction in the Tropics Contribute to the Mammalian Latitudinal Diversity Gradient. *PLOS Biology*, 12, e1001775.
- Rougier, G.W., Wible, J.R., Beck, R.M.D. & Apesteguía, S. (2012). The miocene mammal necrolestes demonstrates the survival of a mesozoic nontherian lineage into the late cenozoic of south america. *Proceedings of the National Academy of Sciences*, 109, 20053–20058.
- Saupe, E.E. & Myers, C.E. (2021). *Macroevolution*. Springer, Cham, pp. 149–167.
- Sauquet, H., Balthazar, M. von, Magallón, S., Doyle, J.A., Endress, P.K., Bailes, E.J., *et al.* (2017). The ancestral flower of angiosperms and its early diversification. *Nature Communications*, 8, 16047.
- Schemske, D.W., Mittelbach, G.G., Cornell, H.V., Sobel, J.M. & Roy, K. (2009). Is There a Latitudinal Gradient in the Importance of Biotic Interactions? *Annual Review of Ecology, Evolution, and Systematics*, 40, 245–269.
- Seiffert, E.R., Tejedor, M.F., Fleagle, J.G., Novo, N.M., Cornejo, F.M., Bond, M., *et al.* (2020). A parapithecoid stem anthropoid of african origin in the paleogene of south america. *Science*, 368, 194–197.

- Selkin, P.A., Strömberg, C.A.E., Dunn, R., Kohn, M.J., Carlini, A.A., Davies-Vollum, K.S., *et al.* (2015). [Climate, dust, and fire across the eocene-oligocene transition, patagonia](#). *Geology*, 43, 567–570.
- Sepkoski, D. (2012). *Rereading the Fossil Record: The Growth of Paleobiology as an Evolutionary Discipline*. University of Chicago Press, Chicago, IL.
- Sepkoski, J.J. (1984). [A kinetic model of Phanerozoic taxonomic diversity. III. Post-Paleozoic families and mass extinctions](#). *Paleobiology*, 10, 246–267.
- Sepkoski, J.J. (2002). [A compendium of fossil marine animal genera](#). *Bulletins of American Paleontology*, 363, 1–560.
- Sepkoski, J.J., Bambach, R.K., Raup, D.M. & Valentine, J.W. (1981). [Phanerozoic marine diversity and the fossil record](#). *Nature*, 293, 435–437.
- Servais, T., Antoine, P.-O., Danielan, T., Lefebvre, B. & Meyer-Berthaud, B. (2012). Paleontology in france: 200 years in the footsteps of cuvier and lamarck. *Palaeontologia Electronica*, 15.
- Signor, P.W.,I. & Lipps, J.H. (1982). [Sampling bias, gradual extinction patterns and catastrophes in the fossil record](#). *Geological Society of America Special papers*, 0.
- Simpson, G.G. (1940). [Review of the mammal-bearing tertiary of south america](#). *Proceedings of the American Philosophical Society*, 83, 649–709.
- Simpson, G.G. (1944). *Tempo and mode in evolution*. Columbia University Press.
- Simpson, G.G. (1950). [History of the fauna of latin america](#). *American Scientist*, 38, 361–389.
- Simpson, G.G. (1967). The beginning of the age of mammals in south america. Part II. *Bulletin of the American Museum of Natural History*, 137, 1–260.
- Simpson, G.G. (1970). [Mammals from the early cenozoic of chubut, argentina](#). *Breviora*, 360, 1–13.
- Simpson, G.G. (1980). *Splendid isolation : The curious history of south american mammals*. Yale University Press, New Haven.
- Stanley, S.M. (1979). *Macroevolution: Pattern and process*. W. H. Freeman, San Francisco.
- Stebbins, G.L. (1974). Trends of angiosperm phylogeny - the ecological basis of diversity. Cambridge, Mass, pp. 156–170.
- Stebbins, G.L. (1981). [Coevolution of grasses and herbivores](#). *Annals of the Missouri Botanical Garden*, 68, 75–86.
- Stehlin, H.G. (1909). Remarques sur les faunules de mammifères des couches éocènes et oligocènes du Bassin de Paris. *Bulletin de la Société Géologique de France*, 488–520.
- Stein, A., Gerstner, K. & Kreft, H. (2014). [Environmental heterogeneity as a universal driver of species richness across taxa, biomes and spatial scales](#). *Ecology Letters*, 17, 866–880.
- Strömberg, C.A.E., Dunn, R.E., Madden, R.H., Kohn, M.J. & Carlini, A.A. (2013). [Decoupling the spread of grasslands from the evolution of grazer-type herbivores in South America](#). *Nature Communications*, 4, 1478.
- Stucky, R.K. (2014). [24. Mammalian Faunas in North America of Bridgerian to Early Arikareean “Ages” \(Eocene and Oligocene\)](#). Princeton University Press, pp. 464–493.
- Stutz, N.S., Ribeiro, A.M., Charruault, A.-L., Boivin, M., Pujos, F., Salas-Gismondi, R., *et al.* (2026). [Hidden diversity of palaeogene metatherians: A new family of polydolopimorphian marsupials from peruvian amazonia](#). *Zoological Journal of the Linnean Society*, 206, zlag006.
- Tejada, J.V., Antoine, P.-O., Münch, P., Billet, G., Hautier, L., Delsuc, F., *et al.* (2024). [Bayesian total-evidence dating revisits sloth phylogeny and biogeography: A cautionary tale on morphological clock analyses](#). *Systematic Biology*, 73, 125–139.
- Tietje, M., Antonelli, A., Baker, W.J., Govaerts, R., Smith, S.A. & Eiserhardt, W.L. (2022). [Global variation in diversification rate and species richness are unlinked in plants](#). *Proceedings of the National Academy of Sciences*, 119, e2120662119.
- Ugarte, P.D. de S., Nascimento, J.C.S. & Pires, M.M. (2025). [Spatiotemporal variability in the south](#)

- american mammalian fossil record and its impact on macroevolutionary inference. *Frontiers in Mammal Science*, 3.
- Valentine, J.W. (1969). Patterns of taxonomic and ecological structure of the shelf benthos during phanerozoic time. *Palaeontology*, 12, 684–709.
- Vasconcelos, T., O’Meara, B.C. & Beaulieu, J.M. (2022). Retiring “cradles” and “museums” of biodiversity. *The American Naturalist*, 199, 194–205.
- Viñola-López, L.W., Cerda, I.A., Correa-Narvaez, J., Codorniú, L., Borges-Sellén, C.R., Arano-Ruiz, A.F., *et al.* (2022). New fossils shed light on the late cretaceous terrestrial community in the caribbean and the first american biotic interchange. *Cretaceous Research*, 130, 105067.
- Vizcaíno, S.F. (2009). The teeth of the “toothless”: novelties and key innovations in the evolution of xenarthrans (Mammalia, Xenarthra). *Paleobiology*, 35, 343–366.
- Waddell, P.J., Cao, Y., Hasegawa, M. & Mindell, D.P. (1999). Assessing the cretaceous superordinal divergence times within birds and placental mammals by using whole mitochondrial protein sequences and an extended statistical framework. *Systematic Biology*, 48, 119–137.
- Wallace, A.R. (1876). *The geographical distribution of animals. With a study of the relations of living and extinct faunas as elucidating the past changes of the earth’s surface*. Harper; brothers, New York.
- Wallace, A.R. (1878). *Tropical Nature, and Other Essays*. Macmillan; Company.
- Webb, S.D. (1985). Late cenozoic mammal dispersals between the americas.
- Westerhold, T., Marwan, N., Drury, A.J., Liebrand, D., Agnini, C., Anagnostou, E., *et al.* (2020). An astronomically dated record of earth’s climate and its predictability over the last 66 million years. *Science*, 369, 1383–1387.
- Woodburne, M.O., Goin, F.J., Bond, M., Carlini, A.A., Gelfo, J.N., López, G.M., *et al.* (2014). Paleogene Land Mammal Faunas of South America; a Response to Global Climatic Changes and Indigenous Floral Diversity. *Journal of Mammalian Evolution*, 21, 1–73.