

# Merge and Count

## Merge and count step.

- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.

$i = 6$



3	7	10	14	18	19
---	---	----	----	----	----



2	11	16	17	23	25
---	----	----	----	----	----

two sorted halves

--	--	--	--	--	--	--	--	--	--	--	--

auxiliary array

Total:

# Merge and Count

## Merge and count step.

- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.

$i = 6$



3	7	10	14	18	19
---	---	----	----	----	----



2	11	16	17	23	25
---	----	----	----	----	----

two sorted halves

6

2											
---	--	--	--	--	--	--	--	--	--	--	--

auxiliary array

Total: 6

# Merge and Count

## Merge and count step.

- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.

$i = 6$



3	7	10	14	18	19
---	---	----	----	----	----



2	11	16	17	23	25
---	----	----	----	----	----

two sorted halves

6

2											
---	--	--	--	--	--	--	--	--	--	--	--

auxiliary array

Total: 6

# Merge and Count

## Merge and count step.

- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.

$i = 6$



3	7	10	14	18	19
---	---	----	----	----	----



2	11	16	17	23	25
---	----	----	----	----	----

two sorted halves

6

2	3										
---	---	--	--	--	--	--	--	--	--	--	--


auxiliary array

Total: 6

# Merge and Count

## Merge and count step.


- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.



# Merge and Count

## Merge and count step.


- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.



# Merge and Count

## Merge and count step.


- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.



# Merge and Count

## Merge and count step.

- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.




Total: 6



# Merge and Count

## Merge and count step.


- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.



# Merge and Count

## Merge and count step.

- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.




Total: 6 + 3

# Merge and Count

## Merge and count step.

- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.




Total: 6 + 3

# Merge and Count

## Merge and count step.

- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.




Total: 6 + 3

# Merge and Count

## Merge and count step.

- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.




Total: 6 + 3

# Merge and Count

## Merge and count step.

- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.




Total:  $6 + 3 + 2$

# Merge and Count

## Merge and count step.

- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.




Total:  $6 + 3 + 2$

# Merge and Count

## Merge and count step.

- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.




Total:  $6 + 3 + 2 + 2$



# Merge and Count

## Merge and count step.

- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.




Total:  $6 + 3 + 2 + 2$

# Merge and Count

## Merge and count step.

- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.




Total:  $6 + 3 + 2 + 2$

# Merge and Count

## Merge and count step.

- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.




Total:  $6 + 3 + 2 + 2$

# Merge and Count

## Merge and count step.

- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.




Total:  $6 + 3 + 2 + 2$

## Merge and Count

Merge and count step.

- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.




Total:  $6 + 3 + 2 + 2$

# Merge and Count

## Merge and count step.

- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.




Total:  $6 + 3 + 2 + 2 + 0$

# Merge and Count

## Merge and count step.

- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.




Total:  $6 + 3 + 2 + 2 + 0$

# Merge and Count

## Merge and count step.

- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.




Total:  $6 + 3 + 2 + 2 + 0 + 0$



# Merge and Count

## Merge and count step.

- Given two sorted halves, count number of inversions where  $a_i$  and  $a_j$  are in different halves.
- Combine two sorted halves into sorted whole.



Total:  $6 + 3 + 2 + 2 + 0 + 0 = 13$