Statistics-S2 - 2012-January

Question 1

The time in minutes that Elaine takes to checkout at her local supermarket follows a continuous uniform distribution defined over the interval [3, 9].

Find

(a) Elaine's expected checkout time,

(1)

(b) the variance of the time taken to checkout at the supermarket,

(2)

(c) the probability that Elaine will take more than 7 minutes to checkout.

(2)

Given that Elaine has already spent 4 minutes at the checkout,

(d) find the probability that she will take a total of less than 6 minutes to checkout.

(3)

Question 2

David claims that the weather forecasts produced by local radio are no better than those achieved by tossing a fair coin and predicting rain if a head is obtained or no rain if a tail is obtained. He records the weather for 30 randomly selected days. The local radio forecast is correct on 21 of these days.

Test David's claim at the 5% level of significance.

State your hypotheses clearly.

(7)

Question 3

The	The probability of a telesales representative making a sale on a customer call is 0.15		
Fin	d the probability that		
(a)	no sales are made in 10 calls,		
		(2)	
(b)	more than 3 sales are made in 20 calls.	(2)	
Rep	presentatives are required to achieve a mean of at least 5 sales each day.		
(c)	Find the least number of calls each day a representative should make to achieve the	nis	
	requirement.	(2)	
(d)	Calculate the least number of calls that need to be made by a representative for t		
	probability of at least 1 sale to exceed 0.95	(3)	
	Question 4		
Aw	vebsite receives hits at a rate of 300 per hour.		
	vebsite receives hits at a rate of 300 per hour. State a distribution that is suitable to model the number of hits obtained duri a 1 minute interval.	ng	
	State a distribution that is suitable to model the number of hits obtained duri a 1 minute interval.	ng (1)	
(a)	State a distribution that is suitable to model the number of hits obtained duri a 1 minute interval. State two reasons for your answer to part (a).		
(a) (b)	State a distribution that is suitable to model the number of hits obtained duri a 1 minute interval. State two reasons for your answer to part (a).	(1)	
(a) (b) Fine	State a distribution that is suitable to model the number of hits obtained duri a 1 minute interval. State two reasons for your answer to part (a).	(1)	
(a) (b) Fine	State a distribution that is suitable to model the number of hits obtained duri a 1 minute interval. State two reasons for your answer to part (a). d the probability of 10 hits in a given minute,	(1)	
(a) (b) Fine (c)	State a distribution that is suitable to model the number of hits obtained duri a 1 minute interval. State two reasons for your answer to part (a). d the probability of 10 hits in a given minute, at least 15 hits in 2 minutes.	(1) (2) (3)	
(a) (b) Fine (c) (d)	State a distribution that is suitable to model the number of hits obtained duri a 1 minute interval. State two reasons for your answer to part (a). d the probability of 10 hits in a given minute, at least 15 hits in 2 minutes.	(1)	
(a) (b) Fine (c) (d)	State a distribution that is suitable to model the number of hits obtained duri a 1 minute interval. State two reasons for your answer to part (a). d the probability of 10 hits in a given minute, at least 15 hits in 2 minutes.	(1) (2) (3)	
(a) (b) Fine (c) (d) The	State a distribution that is suitable to model the number of hits obtained duri a 1 minute interval. State two reasons for your answer to part (a). d the probability of 10 hits in a given minute, at least 15 hits in 2 minutes.	(1) (2) (3)	

Question 5

The probability of an electrical component being defective is 0.075. The component is supplied in boxes of 120.

(a) Using a suitable approximation, estimate the probability that there are more than 3 defective components in a box.

(5)

A retailer buys 2 boxes of components.

(b) Estimate the probability that there are at least 4 defective components in each box.

(2)

Question 6

A random variable X has probability density function given by

$$f(x) = \begin{cases} \frac{1}{2} & 0 \le x < 1 \\ x - \frac{1}{2} & 1 \le x \le k \\ 0 & \text{otherwise} \end{cases}$$

where k is a positive constant.

(a) Sketch the graph of f(x).

(2)

(b) Show that
$$k = \frac{1}{2}(1+\sqrt{5})$$
.

(4)

(c) Define fully the cumulative distribution function F(x).

(6)

(d) Find P(0.5 < X < 1.5).

(2)

(e) Write down the median of X and the mode of X.

(2)

(f) Describe the skewness of the distribution of X. Give a reason for your answer.

(2)

Question 7

- (a) Explain briefly what you understand by
 - (i) a critical region of a test statistic,
 - (ii) the level of significance of a hypothesis test.

(2)

- (b) An estate agent has been selling houses at a rate of 8 per month. She believes that the rate of sales will decrease in the next month.
 - Using a 5% level of significance, find the critical region for a one tailed test of the hypothesis that the rate of sales will decrease from 8 per month.
 - (ii) Write down the actual significance level of the test in part (b)(i).

(3)

The estate agent is surprised to find that she actually sold 13 houses in the next month. She now claims that this is evidence of an increase in the rate of sales per month.

(c) Test the estate agent's claim at the 5% level of significance. State your hypotheses clearly.

(5)