

Core-Maths-C4 - 2010-June

Question 1




Figure 1

Figure 1 shows part of the curve with equation $y = \sqrt{0.75 + \cos^2 x}$. The finite region R , shown shaded in Figure 1, is bounded by the curve, the y -axis, the x -axis and the line with equation $x = \frac{\pi}{3}$.

- (a) Complete the table with values of y corresponding to $x = \frac{\pi}{6}$ and $x = \frac{\pi}{4}$.

x	0	$\frac{\pi}{12}$	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$
y	1.3229	1.2973			1

(2)

- (b) Use the trapezium rule

- (i) with the values of y at $x = 0$, $x = \frac{\pi}{6}$ and $x = \frac{\pi}{3}$ to find an estimate of the area of R .

Give your answer to 3 decimal places.

- (ii) with the values of y at $x = 0$, $x = \frac{\pi}{12}$, $x = \frac{\pi}{6}$, $x = \frac{\pi}{4}$ and $x = \frac{\pi}{3}$ to find a further estimate of the area of R . Give your answer to 3 decimal places.

(6)

Question 2

Using the substitution $u = \cos x + 1$, or otherwise, show that

$$\int_0^{\frac{\pi}{2}} e^{\cos x + 1} \sin x \, dx = e(e - 1)$$

(6)

Question 3

A curve C has equation

$$2^x + y^2 = 2xy$$

Find the exact value of $\frac{dy}{dx}$ at the point on C with coordinates $(3, 2)$.

(7)

Question 4

A curve C has parametric equations

$$x = \sin^2 t, \quad y = 2 \tan t, \quad 0 \leq t < \frac{\pi}{2}$$

(a) Find $\frac{dy}{dx}$ in terms of t .

(4)

The tangent to C at the point where $t = \frac{\pi}{3}$ cuts the x -axis at the point P .

(b) Find the x -coordinate of P .

(6)

Question 5

$$\frac{2x^2 + 5x - 10}{(x-1)(x+2)} \equiv A + \frac{B}{x-1} + \frac{C}{x+2}$$

- (a) Find the values of the constants A , B and C .

(4)

- (b) Hence, or otherwise, expand $\frac{2x^2 + 5x - 10}{(x-1)(x+2)}$ in ascending powers of x , as far as the term in x^2 . Give each coefficient as a simplified fraction.

(7)

Question 6

$$f(\theta) = 4 \cos^2 \theta - 3 \sin^2 \theta$$

- (a) Show that $f(\theta) = \frac{1}{2} + \frac{7}{2} \cos 2\theta$.

(3)

- (b) Hence, using calculus, find the exact value of $\int_0^{\frac{\pi}{2}} \theta f(\theta) \, d\theta$.

(7)

Question 7

The line l_1 has equation $\mathbf{r} = \begin{pmatrix} 2 \\ 3 \\ -4 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$, where λ is a scalar parameter.

The line l_2 has equation $\mathbf{r} = \begin{pmatrix} 0 \\ 9 \\ -3 \end{pmatrix} + \mu \begin{pmatrix} 5 \\ 0 \\ 2 \end{pmatrix}$, where μ is a scalar parameter.

Given that l_1 and l_2 meet at the point C , find

(a) the coordinates of C .

(3)

The point A is the point on l_1 where $\lambda = 0$ and the point B is the point on l_2 where $\mu = -1$.

(b) Find the size of the angle ACB . Give your answer in degrees to 2 decimal places.

(4)

(c) Hence, or otherwise, find the area of the triangle ABC .

(5)

Question 8




Figure 2

Figure 2 shows a cylindrical water tank. The diameter of a circular cross-section of the tank is 6 m. Water is flowing into the tank at a constant rate of $0.48\pi \text{ m}^3 \text{ min}^{-1}$. At time t minutes, the depth of the water in the tank is h metres. There is a tap at a point T at the bottom of the tank. When the tap is open, water leaves the tank at a rate of $0.6\pi h \text{ m}^3 \text{ min}^{-1}$.

- (a) Show that t minutes after the tap has been opened

$$75 \frac{dh}{dt} = (4 - 5h) \quad (5)$$

When $t = 0$, $h = 0.2$

- (b) Find the value of t when $h = 0.5$

(6)
