

Session 1: Vector spaces

Optimization and Computational Linear Algebra for Data Science

Léo Miolane

Contents

1. Subspaces
2. Linear dependency
3. Properties of the dimension
4. Coordinates
5. Why do we care about all these things ?

Application to data science

Questions ?

Questions ?

Questions ?

Subspaces

What are the subspaces of \mathbb{R}^2 ?

The span is always a subspace

Proposition

Let $x_1, \dots, x_k \in V$. Then, $\text{Span}(x_1, \dots, x_k)$ is a subspace of V .

Linear dependency

A useful lemma

Lemma

Let $v_1, \dots, v_n \in V$ and let $x_1, \dots, x_k \in \text{Span}(v_1, \dots, v_n)$.
Then, if $k > n$, x_1, \dots, x_k are linearly dependent.

Abuse of language: Instead of saying « x_1, \dots, x_k are linearly dependent», we should have said «the family (x_1, \dots, x_k) is linearly dependent».

Basis, dimension

The dimension is well defined!

Theorem

If V admits a basis (v_1, \dots, v_n) , then every basis of V has also n vectors. We say that V has dimension n and write $\dim(V) = n$.

Proof.



Properties of the dimension

Proposition

Let V be a vector space that has dimension $\dim(V) = n$. Then

- Any family of vectors of V that are linearly independent contains at most n vectors.

i.e. if $x_1, \dots, x_k \in V$ are linearly independent, then $k \leq n$.

- Any family of vectors of V that spans V contains at least n vectors.

i.e. if $x_1, \dots, x_k \in V$ are such that $\text{Span}(x_1, \dots, x_k) = V$, then $k \geq n$.

Proof.



Properties of the dimension

Proposition

Let V be a vector space that has dimension $\dim(V) = n$. Then

- Any family of vectors of V that are linearly independent contains at most n vectors.

i.e. if $x_1, \dots, x_k \in V$ are linearly independent, then $k \leq n$.

- Any family of vectors of V that spans V contains at least n vectors.

i.e. if $x_1, \dots, x_k \in V$ are such that $\text{Span}(x_1, \dots, x_k) = V$, then $k \geq n$.

Proof.



Properties of the dimension

Proposition

Let V be a vector space of dimension n and let $x_1, \dots, x_n \in V$.

1. If x_1, \dots, x_n are linearly independent, then (x_1, \dots, x_n) is a basis of V .
2. If $\text{Span}(x_1, \dots, x_n) = V$, then (x_1, \dots, x_n) is a basis of V .

Very useful to show that a family of vector forms a basis!

Proof.



An inequality

Proposition

Let U and V be two subspaces of \mathbb{R}^n . Assume that $U \subset V$. Then

$$\dim(U) \leq \dim(V) \leq n.$$

If **moreover** $\dim(U) = \dim(V)$, then $U = V$.

A bit of vocabulary

Definition

Let S be a subspace of \mathbb{R}^n .

- ❖ We call S a *line* if $\dim(S) = 1$.
- ❖ We call S a *hyperplane* if $\dim(S) = n - 1$.

Coordinates

Coordinates of a vector in a basis

Definition

If (v_1, \dots, v_n) is a basis of V , then for every $x \in V$ there exists a unique vector $(\alpha_1, \dots, \alpha_n) \in \mathbb{R}^n$ such that

$$x = \alpha_1 v_1 + \dots + \alpha_n v_n.$$

We say that $(\alpha_1, \dots, \alpha_n)$ are the coordinates of x in the basis (v_1, \dots, v_n) .

Proof.



Exercise

1. Show that the vectors $v_1 = (1, 1)$ and $v_2 = (1, -1)$ form a basis of \mathbb{R}^2 .
2. Express the coordinates of $u = (x, y)$ in the basis (v_1, v_2) in terms of x and y .

Exercise

1. Show that the vectors $v_1 = (1, 1)$ and $v_2 = (1, -1)$ form a basis of \mathbb{R}^2 .
2. Express the coordinates of $u = (x, y)$ in the basis (v_1, v_2) in terms of x and y .

Why do we care about this ?

Application to image compression

✚ Image = grid of pixels