

# Optimization and Computational Linear Algebra for Data Science

## Homework 3: Rank

Due on September 27, 2020

- 
- Unless otherwise stated, all answers must be mathematically justified.
  - Partial answers will be graded.
  - Your submission has to be uploaded to Gradescope. Indicate Gradescope the page on which each problem is written.
  - You can work in groups but each student must write his/her own solution based on his/her own understanding of the problem. Please list on your submission the students you work with for the homework (this will not affect your grade).
  - Problems with a  $(\star)$  are extra credit, they will not (directly) contribute to your score of this homework. However, for every 4 extra credit questions successfully answered your lowest homework score get replaced by a perfect score.
  - If you have any questions, feel free to contact me ([lm4271@nyu.edu](mailto:lm4271@nyu.edu)) or to stop at the office hours.
- 

**Problem 3.1** (2 points). Let  $A \in \mathbb{R}^{n \times n}$ .

- (a) Show that if  $A = \alpha \text{Id}_n$  for some  $\alpha \in \mathbb{R}$ , then for all  $B \in \mathbb{R}^{n \times n}$  we have  $AB = BA$ .
- (b) Conversely, show that if for all  $B \in \mathbb{R}^{n \times n}$  we have  $AB = BA$ , then there exists  $\alpha \in \mathbb{R}$  such that  $A = \alpha \text{Id}_n$ .

**Problem 3.2** (3 points). Let  $M \in \mathbb{R}^{n \times m}$  and  $r = \text{rank}(M)$ . Show that there exists  $A \in \mathbb{R}^{n \times r}$  and  $B \in \mathbb{R}^{r \times m}$  such that  $M = AB$ .

**Problem 3.3** (3 points). Let  $A \in \mathbb{R}^{n \times m}$ .

- (a) Let  $M \in \mathbb{R}^{m \times m}$  be an invertible matrix. Show that
$$\text{rank}(AM) = \text{rank}(A).$$
- (b) Let  $M \in \mathbb{R}^{n \times n}$  be an invertible matrix. Show that
$$\text{rank}(MA) = \text{rank}(A).$$

**Problem 3.4** (2 points). The trace  $\text{Tr}(M)$  of a  $k \times k$  matrix  $M$  is defined as the sum of its diagonal coefficients, i.e.

$$\text{Tr}(M) = \sum_{i=1}^k M_{i,i}.$$

- (a) Let  $A \in \mathbb{R}^{n \times m}$  and  $B \in \mathbb{R}^{m \times n}$ . Show that  $\text{Tr}(AB) = \text{Tr}(BA)$ .
- (b) For  $A, B, C \in \mathbb{R}^{n \times n}$ , do we have  $\text{Tr}(ABC) = \text{Tr}(CAB) = \text{Tr}(ACB)$  ?

**Problem 3.5**  $(\star)$ . Let  $A \in \mathbb{R}^{10 \times 10}$  such that  $A^{2020} = \underbrace{A \times A \times \cdots \times A}_{2020 \text{ times}} = 0$ . Show that  $A^{10} = 0$ .

