

Session 9: Convex functions

Optimization and Computational Linear Algebra for Data Science

Contents

1. Recap of the videos
2. Convex sets and convex functions
3. Convex functions and derivatives
4. Jensen's inequality

Optimization

In machine learning, we often have to minimize functions

$$f(\theta) = \text{Loss}(\text{data}, \text{model}_\theta) \quad \text{with respect to} \quad \theta \in \mathbb{R}^n.$$

- ❖ For $n = 1, 2$, one could plot f to find the minimizer.
- ❖ This is intractable for larger dimension.

We will

- ❖ focus on convex cost functions f .
- ❖ study gradient descent algorithms to minimize f .

Convex vs non-convex

Convex

Non-convex

Gradient/Hessian

For $f : \mathbb{R}^n \rightarrow \mathbb{R}$:

▣ Gradient at $x \in \mathbb{R}^n$:

$$\nabla f(x) = \begin{pmatrix} \frac{\partial f}{\partial x_1}(x) \\ \vdots \\ \frac{\partial f}{\partial x_n}(x) \end{pmatrix} \in \mathbb{R}^n$$

▣ Hessian at $x \in \mathbb{R}^n$:

$$H_f(x) = \begin{pmatrix} \frac{\partial^2 f}{\partial x_1^2}(x) & \frac{\partial^2 f}{\partial x_1 \partial x_2}(x) & \cdots & \frac{\partial^2 f}{\partial x_1 \partial x_n}(x) \\ \frac{\partial^2 f}{\partial x_2 \partial x_1}(x) & \frac{\partial^2 f}{\partial x_2^2}(x) & \cdots & \frac{\partial^2 f}{\partial x_2 \partial x_n}(x) \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1}(x) & \frac{\partial^2 f}{\partial x_n \partial x_2}(x) & \cdots & \frac{\partial^2 f}{\partial x_n^2}(x) \end{pmatrix} \in \mathbb{R}^{n \times n}$$

Taylor's formulas

Let $x \in \mathbb{R}^n$. Heuristically, for $h \in \mathbb{R}^n$ "small", we have

$$f(x+h) \simeq f(x) + \langle \nabla f(x), h \rangle.$$

Taylor's formulas

Let $x \in \mathbb{R}^n$. Heuristically, for $h \in \mathbb{R}^n$ "small", we have

$$f(x+h) \simeq f(x) + \langle \nabla f(x), h \rangle + \frac{1}{2} h^\top H_f(x) h.$$

Convex sets

Convex set

Definition

A set $S \subset \mathbb{R}^n$ is called a convex set if for all $x, y \in C$ and all $\alpha \in [0, 1]$,

$$\alpha x + (1 - \alpha)y \in C.$$

Exercise

1. Show that any subspace S of \mathbb{R}^n is convex.
2. Let $\|\cdot\|$ be a (arbitrary) norm and $r \geq 0$. Show that the "ball" of radius r :

$$B(r) = \{x \in \mathbb{R}^n \mid \|x\| \leq r\}$$

is convex.

Convex functions

Convex / concave functions

Definition

A function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is convex if for all $x, y \in \mathbb{R}^n$ and all $\alpha \in [0, 1]$,

$$f(\alpha x + (1 - \alpha)y) \leq \alpha f(x) + (1 - \alpha)f(y). \quad (1)$$

Convex / concave functions

Definition

A function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is convex if for all $x, y \in \mathbb{R}^n$ and all $\alpha \in [0, 1]$,

$$f(\alpha x + (1 - \alpha)y) \leq \alpha f(x) + (1 - \alpha)f(y). \quad (1)$$

- ❑ We say that f is *strictly convex* if there is strict inequality in (1) whenever $x \neq y$ and $\alpha \in (0, 1)$.
- ❑ A function f is called *concave* if $-f$ is convex.

Exercise

1. Show that any linear map $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is convex and concave.
2. Show that a norm $\| \cdot \|$ is convex.
3. Show that the sum of two convex functions is also a convex function.

Convex functions and derivatives

Convex functions vs their tangents

Proposition

A differentiable function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is convex if and only if for all $x, y \in \mathbb{R}^n$

$$f(y) \geq f(x) + \langle \nabla f(x), (y - x) \rangle.$$

Proof

Proof

Proof

Minimizers of a convex function

Corollary

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a differentiable convex function and $x \in \mathbb{R}^n$.
Then

$$x \text{ is a minimizer of } f \iff \nabla f(x) = 0.$$

Hessian of convex function

Proposition

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a twice-differentiable function. Then f is convex if and only if for all $x \in \mathbb{R}^n$, $H_f(x)$ is positive semi-definite.

Hessian of convex function

Proposition

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a twice-differentiable function. Then f is convex if and only if for all $x \in \mathbb{R}^n$, $H_f(x)$ is positive semi-definite.

Jensen's inequality

Jensen's inequality

Theorem

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function. Then for all $x_1, \dots, x_k \in \mathbb{R}^n$ and all $\alpha_1, \dots, \alpha_k \geq 0$ such that $\sum_{i=1}^k \alpha_i = 1$ we have

$$f\left(\sum_{i=1}^k \alpha_i x_i\right) \leq \sum_{i=1}^k \alpha_i f(x_i).$$

More generally, if X is a random variable that takes value in \mathbb{R}^n we have

$$f(\mathbb{E}[X]) \leq \mathbb{E}[f(X)].$$

Example: entropy

Example: entropy

Questions?

Questions?