

Mathematics Extension 1 11–12 (2024)

curriculum.nsw.edu.au

Generated Apr 2026

Downloaded April 2026

© 2026 NSW Education Standards Authority

NESA acknowledges Traditional Owners and Custodians of Country throughout NSW, and pays respect to Elders past and present. NESA recognises Aboriginal Peoples' continuing Cultures and Connections to lands, waters, skies and Community.

The documents on the NSW Education Standards Authority (NESA) website and the NSW Curriculum website contain material prepared by NESA for and on behalf of the Crown in right of the State of New South Wales. The material is protected by Crown copyright.

These websites hold the only official and up-to-date versions of the documents available on the internet. Any other copies of these documents, or parts of these documents, that may be found elsewhere on the internet might not be current and are not authorised. You cannot rely on copies from any other source.

All rights are reserved. No part of the material may be:

- reproduced in Australia or in any other country by any process, electronic or otherwise, in any material form
- transmitted to any other person or stored electronically in any form without the written permission of NESA except as permitted by the *Copyright Act 1968* (Cth).

When you access the material, you agree:

- to use the material for research or study, criticism or review, reporting news and parody or satire
- to use the material for information purposes only
- not to modify the material or any part of the material without the written permission of NESA
- to reproduce a single copy for personal bona fide study use only and not to reproduce any major extract or the entire material without the permission of NESA
- to include this copyright notice in any copy made
- to acknowledge that NESA is the source of the material.

The documents may include third-party copyright material such as photos, diagrams, quotations, cartoons and artworks. This material is protected by Australian and international copyright laws and may not be reproduced or transmitted in any format without the copyright owner's permission. Unauthorised reproduction, transmission or commercial use of such copyright material may result in prosecution.

NESA has made all reasonable attempts to locate the owners of third-party copyright material. NESA invites anyone from whom permission has not been sought to contact the [Copyright Officer](#).

Special arrangements applying to the NSW Curriculum Reform

As part of the NSW Curriculum Reform process, NESA grants a limited non-exclusive licence to:

- teachers employed in NSW government schools and registered non-government schools
- representatives from peak school education bodies, such as the NSW Department of Education, Catholic Schools NSW, the Association of Independent Schools of NSW and TAFE NSW
- parents of children registered for home schooling

to use, modify and adapt the NSW syllabuses for **non-commercial educational use only**. The adaptation must not have the effect of bringing NESA into disrepute.

Note: The above arrangements do not apply to private/home tutoring companies, professional learning service providers, publishers, and other organisations.

For more information on the above or for **commercial use or any other purpose**, please contact the [Copyright Officer](#) for permission.

Email: copyright@nesa.nsw.edu.au

Table of contents

Table of contents	4
Mathematics Extension 1 11–12 (2024)	5
Implementation from 2026	5
Overview	5
Syllabus overview	5
Course description	7
Rationale	10
Aim	11
Table of outcomes	12
Outcomes and content for Year 11	13
Further work with functions	13
Polynomials.....	15
Further trigonometry.....	17
Permutations and combinations.....	18
The binomial theorem	20
Outcomes and content for Year 12	21
Proof by mathematical induction	21
Introduction to vectors	22
Inverse trigonometric functions	25
Further calculus skills.....	27
Further applications of calculus.....	28
The binomial distribution and sampling distribution of the mean	30

Mathematics Extension 1 11–12 (2024)

Implementation from 2026

The new Mathematics Extension 1 11–12 Syllabus (2024) is to be implemented from 2026 and will replace the [Mathematics Extension 1 Stage 6 Syllabus \(2017\)](#).

2026, Term 1

- Start teaching the new syllabus for Year 11
- Start implementing the new Year 11 school-based assessment requirements
- Continue to teach the [Mathematics Extension 1 Stage 6 Syllabus \(2017\)](#) for Year 12

2026, Term 4

- Start teaching the new syllabus for Year 12
- Start implementing the new Year 12 school-based assessment requirements

2027

- First HSC examination for the new syllabus

Overview

Syllabus overview

Organisation of Mathematics Extension 1 11–12

The organisation of outcomes and content for Mathematics Extension 1 11–12 highlights the important role Working mathematically plays across all areas of mathematics and reflects the strengthened connections between concepts. Working mathematically has been embedded in the outcomes and content of the syllabus.

Mathematics Extension 1 outcomes and their related content are organised into 7 areas of study:

- Functions
- Proof
- Vectors
- Trigonometric functions
- Combinatorics
- Calculus
- Statistical analysis.

Figure 1 gives a broad overview of the organisation of content, including the focus areas to be completed in each area of study.




Figure 1: The organisation of Mathematics Extension 1 11–12

Image long description: Seven rows represent the areas of study for Mathematics Extension 1 11–12. Each row is divided in half, showing the focus areas for Year 11 and Year 12. The Functions area of study includes Year 11 focus areas Further work with functions and Polynomials. The Proof area of study includes the Year 12 focus area Proof by mathematical induction. The Vectors area of study includes the Year 12 focus area Introduction to vectors. The Trigonometric functions area of study includes the Year 11 focus area Further trigonometry, and the Year 12 focus area Inverse trigonometric functions. The Combinatorics area of study includes Year 11 focus areas Permutations and combinations, and The binomial theorem. The Calculus area of study includes the Year 12 focus areas Further calculus skills and Further applications of calculus. The Statistical analysis area of study includes the Year 12 focus area The binomial distribution and the sampling distribution of the mean. All content is surrounded by a box labelled with the phrase, ‘Working mathematically through communicating reasoning, understanding and fluency, and problem solving’.

Protocols for collaborating with Aboriginal and Torres Strait Islander Communities

NESA is committed to working in partnership with Aboriginal Communities and supporting teachers, schools and schooling sectors to improve educational outcomes for young people.

It is important to respect appropriate ways of interacting with Aboriginal Communities and Cultural material when teachers plan, program and implement learning experiences that focus on Aboriginal and Torres Strait Islander Priorities.

Indigenous Cultural and Intellectual Property (ICIP) protocols need to be followed. Aboriginal and Torres Strait Islander Peoples’ ICIP protocols include Cultural Knowledges, Cultural Expression and Cultural Property and documentation of Aboriginal and Torres Strait Islander Peoples’ Identities and lived experiences. It is important to recognise the diversity and complexity of different Cultural groups in NSW, as protocols may differ between local Aboriginal Communities.

Teachers should work in partnership with Elders, parents, Community members, Cultural Knowledge Holders, or a local, regional or state Aboriginal Education Consultative Group. It is important to respect Elders and the roles of men and women. Local Aboriginal Peoples should be

invited to share their Cultural Knowledges with students and staff when engaging with Aboriginal histories and Cultural Practices.

Course description

Mathematics Extension 1 focuses on the development of students mathematical arguments and proofs, and use of mathematical models. The course allows students to develop a thorough knowledge and understanding of and competence in further aspects of mathematics as an extension of the Mathematics Advanced 11–12 course.

What students learn

Through the study of Mathematics Extension 1, students:

- develop thorough knowledge, understanding and skills in Working mathematically and in communicating concisely and precisely
- develop rigorous mathematical arguments and proofs, and use mathematical models extensively
- develop awareness of the interconnected nature of mathematics, its beauty and its functionality
- gain an appropriate mathematical background for future pathways that may involve mathematics and its applications.

Course structure and requirements

Year 11 course structure

Area of study	Focus area
Functions	Further work with functions Polynomials
Trigonometric functions	Further trigonometry
Combinatorics	Permutations and combinations The binomial theorem

Year 12 course structure

Area of study	Focus area
Proof	Proof by mathematical induction
Vectors	Introduction to vectors
Trigonometric functions	Inverse trigonometric functions
Calculus	Further calculus skills Further applications of calculus
Statistical analysis	The binomial distribution and sampling distribution of the mean

Safety and risk management

Schools are required to ensure they follow [safety and risk management](#) in delivering the *Mathematics Extension 1 11–12 Syllabus*.

Course enrolment details

Mathematics Extension 1 Year 11

- Course number: 11250
- Course hours: 60
- Course units: 1
- Enrolment type: Elective
- Endorsement type: Board developed

Exclusions

- Mathematics Standard (Year 11, 2 units): 11236
- Mathematics Life Skills (Year 11, 2 units): 17680

Prerequisites

- Nil

Corequisites

- Mathematics Advanced (Year 11, 2 units): 11255

Mathematics Extension 1 Year 12

- Course number: 15250
- Course hours: 60
- Course units: 1
- Enrolment type: Elective
- Endorsement type: Board developed

Exclusions

- Mathematics Standard 1 (Year 12, 2 units): 15231
- Mathematics Standard 1 (examination) (Year 12, 2 units): 15232
- Mathematics Standard 2 (Year 12, 2 units): 15236

- Mathematics Life Skills (Year 12, 2 units): 17680

Prerequisites

- Mathematics Advanced (Year 11, 2 units): 11255
- Mathematics Extension 1 (Year 11, 1 unit) 11250

Corequisites

- Mathematics Advanced (Year 12, 2 units): 15255

HSC information

Information about [curriculum requirements](#) for the HSC are available on [Assessment Certification Examination \(ACE\)](#).

Rationale

Mathematics in the Stage 6 Curriculum

Mathematical ideas have evolved and continue to develop across cultures and have been practised in Australia by Aboriginal and Torres Strait Islander Peoples for thousands of years.

The utility of mathematics extends from counting and measuring to explaining real-world phenomena and its inherent beauty. It has evolved in sophisticated ways to become the language used to describe and understand much of the modern world.

The creative application of mathematical concepts is integral to scientific and technological advancement. The symbolic nature of mathematics provides a precise means of communication.

The Mathematics Stage 6 syllabuses are designed to offer students opportunities to think mathematically. Mathematical thinking is supported through questioning, communicating, reasoning, reflecting and the use of appropriate technology, and is encouraged through providing students with opportunities to generalise, challenge, find connections and to think critically and creatively. Using mathematical thinking, students apply their knowledge and skills to deepen their understanding of the world.

Mathematics Extension 1

Mathematics Advanced, Mathematics Extension 1 and Mathematics Extension 2 form a continuum to provide students with opportunities to acquire knowledge, understanding and skills in mathematical concepts at progressively higher levels and with applications in an increasing number of contexts. Mathematics Advanced, Mathematics Extension 1 and Mathematics Extension 2 provide students with opportunities to learn about the interconnected nature of mathematics, its beauty and its functionality. The concepts and techniques of differential and integral calculus are the basis of the courses.

Mathematics Extension 1 enables students to develop a thorough knowledge and understanding of and competence in further aspects of mathematics as an extension of the Mathematics Advanced course. The course provides opportunities to develop rigorous mathematical arguments and proofs, and to use mathematical models more extensively.

Mathematics Extension 1 supports students in tertiary study in mathematics and related fields.

Aim

The aim of Mathematics Extension 1 is for students to extend their knowledge and understanding of Working mathematically from Mathematics Advanced and Stage 5, further their understanding of the relationship between real-world problems and mathematical models, make connections within mathematics, and enhance their skills in using the language of mathematics to communicate in a concise and systematic manner.

Table of outcomes

MAO-WM-01 Working mathematically

develops understanding and fluency in mathematics through exploring and connecting mathematical concepts, choosing and applying mathematical techniques to solve problems, and communicating their thinking and reasoning coherently and clearly

This outcome is aligned to all content in each Stage.

Year 11	Year 12
ME1-11-01 solves problems involving inequalities, functions and their inverses, graphical relationships between functions, and parametric equations	ME1-12-01 uses mathematical induction to prove results involving sums and divisibility
ME1-11-02 applies the remainder and factor theorem and sums and products of zeroes to solve problems involving polynomials	ME1-12-02 operates with 2D and 3D vectors and uses 2D vectors to solve problems involving motion in two dimensions
ME1-11-03 solves problems in three dimensions using trigonometry and simplifies expressions, proves results and solves problems involving compound angles using trigonometric identities	ME1-12-03 solves problems involving inverse trigonometric functions
ME1-11-04 uses permutations and combinations to solve problems involving counting, ordering and probability	ME1-12-04 selects and applies differentiation and integration techniques to solve problems
ME1-11-05 uses the binomial theorem to solve problems and prove identities	ME1-12-05 applies calculus to solve problems involving polynomials, further rates of change, areas and volumes and differential equations
	ME1-12-06 solves problems involving binomial distributions, sampling distribution of the mean and the central limit theorem

Outcomes and content for Year 11

Further work with functions

Outcomes

A student:

- develops understanding and fluency in mathematics through exploring and connecting mathematical concepts, choosing and applying mathematical techniques to solve problems, and communicating their thinking and reasoning coherently and clearly **MAO-WM-01**
- solves problems involving inequalities, functions and their inverses, graphical relationships between functions, and parametric equations **ME1-11-01**

Content

Graphical relationships

- Examine the relationship between the [graph](#) of $y = f(x)$ and the graph of $y = \frac{1}{f(x)}$ using graphing applications, and graph $y = \frac{1}{f(x)}$ given $y = f(x)$ in algebraic or graphical form, identifying any vertical and horizontal [asymptotes](#) of $y = f(x)$ and $y = \frac{1}{f(x)}$
- Graph $y = \sec x$, $y = \operatorname{cosec} x$ and $y = \cot x$ in both [radians](#) and degrees, identifying key properties including asymptotes, [period](#), [domain](#), [range](#) and symmetry, and compare each graph with the graph of its [reciprocal](#)
- Examine the relationship between the graph of $y = f(x)$ and the graphs of $y = |f(x)|$ and $y = f(|x|)$ using graphing applications, and graph $y = |f(x)|$ and $y = f(|x|)$ given $y = f(x)$ in algebraic or graphical form
- Examine the relationship between the graphs of $y = f(x)$ and $y = g(x)$ and the graphs of $y = f(x) + g(x)$ and $y = f(x) - g(x)$ using graphing applications, and graph $y = f(x) + g(x)$ and $y = f(x) - g(x)$ given $y = f(x)$ and $y = g(x)$ in algebraic or graphical form
- Determine the domains and ranges of the sum and difference of functions where possible, and, if appropriate, verify them using a graphing application
- Apply knowledge of graphical relationships to solve problems involving graphs of functions, justifying conclusions in the context of the problem where appropriate

Inverse functions

- Describe a [function](#) as [one-to-one](#) if every [element](#) in the range of the function corresponds to exactly one element of the domain
- Establish that the [reflection](#) of a point in the line $y = x$ reverses the coordinates of the point
- Define an [inverse function](#) f^{-1} informally as a function that reverses or undoes the effect of the function f
- Recognise that inverse functions exist for one-to-one functions
- Determine the [equation](#) for the inverse function $f^{-1}(x)$ of a given one-to-one function $y = f(x)$ by interchanging the variables x and y in $y = f(x)$
- Compare the graphs of a function and its inverse function using graphing applications, and recognise that the two graphs are reflections of each other in the line $y = x$
- Establish that the domain of $f^{-1}(x)$ is the range of $f(x)$, and the range of $f^{-1}(x)$ is the domain of $f(x)$, and use this relationship to solve problems

- Graph the inverse function of a given one-to-one function
- Explain that the reflection in the line $y = x$ exchanges horizontal and vertical lines and recognise that the horizontal line test can therefore be applied to the graph of $y = f(x)$ to determine whether its reflection in the line $y = x$ is a function
- Apply restrictions to the domain of a function, if it is not one-to-one, to obtain an inverse function
- Define formally $f^{-1}(x)$ to be the inverse function of $f(x)$ if the relationships $f(f^{-1}(x)) = x$ and $f^{-1}(f(x)) = x$ hold, and use this definition to solve problems
- Solve problems based on the relationship between a function and its inverse function using algebraic and graphical techniques, including determining the points of intersection of a function and its inverse, where they exist

Parametric form of a function or relation

- Recognise that a curve may be represented by two [parametric equations](#) that give x and y as functions of a [parameter](#) and that the curve may also have a [Cartesian equation](#) in x and y
- Express [linear](#) and [quadratic functions](#) and [circles](#) in parametric form
- Convert linear and quadratic functions and circles from parametric form to Cartesian form
- Graph linear and quadratic functions and circles expressed in parametric form

Inequalities

- Solve cubic inequalities where the cubic is expressed as a product of [linear](#) factors
- Solve inequalities involving rational [expressions](#) with [variables](#) in the [denominator](#)
- Solve [absolute value](#) inequalities of the form $|ax + b| \geq k$, $|ax + b| \leq k$, $|ax + b| < k$, $|ax + b| > k$, where a , b and k are [constants](#), using algebraic and graphical methods or the characterisation of $|x|$ as the [distance](#) of x from the origin

Polynomials

Outcomes

A student:

- develops understanding and fluency in mathematics through exploring and connecting mathematical concepts, choosing and applying mathematical techniques to solve problems, and communicating their thinking and reasoning coherently and clearly **MAO-WM-01**
- applies the remainder and factor theorem and sums and products of zeroes to solve problems involving polynomials **ME1-11-02**

Content

Language and graphs of polynomials

- Define a [polynomial](#) function $P(x)$ of [degree](#) n , where n is a non-negative [integer](#), to be a function that can be expressed in the form $P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_2 x^2 + a_1 x + a_0$, for real $a_n, \dots, a_0$ and $a_n \neq 0$
- Define the [leading term](#) of $P(x)$ to be the term of highest degree and define the [leading coefficient](#) of $P(x)$ to be the coefficient of the leading term
- Define the constant term of $P(x)$ to be a_0
- Define a polynomial to be [monic](#) if its leading coefficient is 1
- Define the [zero polynomial](#) to be $P(x) = 0$ to be the polynomial with all the coefficients [equal](#) to zero and recognise that the zero polynomial has no leading term, no leading coefficient, no degree and constant term 0
- Determine the degree of $P(x) + Q(x)$ when two non-zero polynomials $P(x)$ and $Q(x)$, of degrees n and m respectively, are added
- Explain how the leading coefficient and the degree determine whether $y \rightarrow \infty$ or $y \rightarrow -\infty$ as $x \rightarrow \infty$ and as $x \rightarrow -\infty$
- Define the zeroes of $P(x)$ to be the numbers α such that $P(\alpha) = 0$, and define the [roots](#) of the polynomial equation $P(x) = 0$ to be its solutions, and recognise that every [real number](#) is a zero of the zero polynomial
- Define α as a repeated zero or multiple zero of a non-zero polynomial $P(x) = (x - \alpha)Q(x)$ when $(x - \alpha)$ is a factor of $Q(x)$, and α as a single zero of $P(x)$ when $(x - \alpha)$ is not a factor of $Q(x)$, for $Q(x) \neq 0$
- Define α as a zero of $P(x)$ of [multiplicity](#) m if $P(x) = (x - \alpha)^m Q(x)$, where m is a positive integer and $Q(\alpha) \neq 0$
- State the multiplicity of each root of a polynomial equation given in factored form
- Find the zeroes of a polynomial that is expressed as a product of linear factors, determine their multiplicity and graph the polynomial

Remainder and factor theorems

- Examine the process of division of polynomials by comparing with the process of division with remainders for [whole numbers](#), and use the terms [dividend](#), [divisor](#), [quotient](#) and [remainder](#)
- Express in the form $P(x) = A(x)Q(x) + R(x)$ the result of dividing $P(x)$ by a divisor $A(x)$, that is not the zero polynomial, with quotient $Q(x)$ and remainder $R(x)$, and explain why either $R(x) = 0$ or $\deg R(x) < \deg A(x)$
- Express the result of the division also in the form $\frac{P(x)}{A(x)} = Q(x) + \frac{R(x)}{A(x)}$

- Explain why division by $x - \alpha$ yields $P(x) = (x - \alpha)Q(x) + r$, where r is a constant, and why $x - \alpha$ is a factor if and only if $r = 0$
- Prove and apply the [remainder theorem](#) for polynomials: when $P(x)$ is divided by $x - \alpha$, the remainder is $P(\alpha)$, and solve related polynomial problems
- Prove and apply the [factor theorem](#) for polynomials: $P(\alpha) = 0$ if and only if $x - \alpha$ is a factor of $P(x)$, and solve related polynomial problems
- Use the factor theorem to find all factors of $P(x)$ of the form $x - \alpha$, where α is an integer, and use division to find the remaining factor of the polynomial

Sums and products of zeroes of polynomials

- Prove that if a quadratic $P(x) = ax^2 + bx + c$ has zeroes α and β then $\alpha + \beta = -\frac{b}{a}$, the sum of the zeroes, and $\alpha\beta = \frac{c}{a}$, the product of the zeroes
- Prove that if a cubic $P(x) = ax^3 + bx^2 + cx + d$ has three zeroes α, β, γ , then $\alpha + \beta + \gamma = -\frac{b}{a}$, $\alpha\beta + \beta\gamma + \gamma\alpha = \frac{c}{a}$ and $\alpha\beta\gamma = -\frac{d}{a}$
- Prove that if a quartic $P(x) = ax^4 + bx^3 + cx^2 + dx + e$ has four zeroes $\alpha, \beta, \gamma, \delta$ then $\alpha + \beta + \gamma + \delta = -\frac{b}{a}$, $\alpha\beta + \alpha\gamma + \alpha\delta + \beta\gamma + \beta\delta + \gamma\delta = \frac{c}{a}$, $\alpha\beta\gamma + \beta\gamma\delta + \gamma\delta\alpha + \delta\alpha\beta = -\frac{d}{a}$ and $\alpha\beta\gamma\delta = \frac{e}{a}$
- Use the formulas for the sums and products of zeroes to solve problems involving zeroes and coefficients of quadratic, cubic and quartic polynomials

Further trigonometry

Outcomes

A student:

- develops understanding and fluency in mathematics through exploring and connecting mathematical concepts, choosing and applying mathematical techniques to solve problems, and communicating their thinking and reasoning coherently and clearly **MAO-WM-01**
- solves problems in three dimensions using trigonometry and simplifies expressions, proves results and solves problems involving compound angles using trigonometric identities **ME1-11-03**

Content

Trigonometry in three dimensions

- Interpret information about a 3-dimensional context given in diagrammatical form
- Interpret information about a 3-dimensional context given in written form
- Apply trigonometry to solve problems in three dimensions where a diagram is given

Further trigonometric identities

- Derive the sum and difference expansions for the trigonometric functions $\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$, $\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$ and $\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$
- Use sum and difference expansion formulas to solve problems and prove results
- Derive the double [angle](#) formulas $\sin 2A = 2\sin A \cos A$, $\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$ and $\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$
- Use the double angle formulas to solve problems and prove results

Further trigonometric equations

- Solve trigonometric equations involving factorisation and/or [substitution](#) of trigonometric identities over restricted domains
- Examine the representations of $a \cos x + b \sin x$ as $R \cos(x \pm \alpha)$ or $R \sin(x \pm \alpha)$ with and without graphing applications, where $a \neq 0$ and $b \neq 0$
- Apply the representations of $a \cos x + b \sin x$ as $R \cos(x \pm \alpha)$ or $R \sin(x \pm \alpha)$ to graph functions, solve equations of the form $a \cos x + b \sin x = c$ over restricted domains, and [model](#) and solve related problems
- Interpret solutions of trigonometric equations graphically with and without graphing applications
- Model and solve practical problems using trigonometric equations and analyse their solutions in a variety of contexts

Permutations and combinations

Outcomes

A student:

- develops understanding and fluency in mathematics through exploring and connecting mathematical concepts, choosing and applying mathematical techniques to solve problems, and communicating their thinking and reasoning coherently and clearly **MAO-WM-01**
- uses permutations and combinations to solve problems involving counting, ordering and probability **ME1-11-04**

Content

- Use the notation $n!$ (read as n [factorial](#)), where $n! = n(n-1)(n-2) \times \dots \times 3 \times 2 \times 1$ for positive integers n
- Use $n! = n \times (n-1)!$ and the convention $0! = 1$ in calculations and to simplify [algebraic expressions](#) involving factorials
- Establish and use the [multiplication principle](#): if a selection can be made in two stages, where there are m choices for the first stage and n choices for the second stage then there are $m \times n$ choices for the selection
- Apply the multiplication principle to explain why the number of ways of ordering n [distinct](#) objects in a straight line is $n!$
- Define a [permutation](#) as an ordered selection of some or all objects from a [set](#) of distinct objects
- Use the notation nP_r to represent an ordered selection of r objects from n distinct objects and observe that ${}^nP_n = n!$ and ${}^nP_0 = 1$
- Use the multiplication principle to establish that the number of ordered selections of r objects from n distinct objects is $n(n-1)(n-2) \times \dots \times (n-r+1)$ and show that
$${}^nP_r = n(n-1)(n-2) \times \dots \times (n-r+1) = \frac{n!}{(n-r)!}$$
- Solve problems involving permutations, including situations where the objects are not all distinct
- Solve problems involving permutations with restrictions on the placement of one or more objects
- Explain why the number of ways to arrange n distinct objects in a circle is $(n-1)!$
- Solve problems involving circular arrangements of distinct objects with or without restrictions on the placement of one or more objects
- Define a [combination](#) and use the notation nC_r or $\binom{n}{r}$ to represent the number of ways of selecting a [subset](#) of r objects from n distinct objects, where order is not important
- Establish and use the formula ${}^nC_r = \frac{n!}{r!(n-r)!}$
- Show that ${}^nC_n = {}^nC_0 = 1$ and ${}^nC_1 = {}^nC_{n-1} = n$
- Show that ${}^nC_r = {}^nC_{n-r}$, for $0 \leq r \leq n$ by selecting r objects from n distinct objects for inclusions and $n-r$ objects from n distinct objects for exclusion
- Prove ${}^nC_r = {}^{n-1}C_{r-1} + {}^{n-1}C_r$ for $1 \leq r \leq n-1$ algebraically and using combinatorial arguments
- Solve problems involving combinations with or without restrictions on the selection of one or more objects

- Solve problems involving both permutations and combinations, including problems which require consideration of cases
- Solve [probability](#) problems involving permutations and combinations

The binomial theorem

Outcomes

A student:

- develops understanding and fluency in mathematics through exploring and connecting mathematical concepts, choosing and applying mathematical techniques to solve problems, and communicating their thinking and reasoning coherently and clearly **MAO-WM-01**
- uses the binomial theorem to solve problems and prove identities **ME1-11-05**

Content

- Recognise that a [binomial](#) is an expression with two terms, and that a [binomial expansion](#) is an expansion of a power of a binomial
- Examine the symmetry formed by the coefficients of decreasing powers of x in the expansion of $(x + y)^n$ for $n = 0, 1, 2, 3, 4, 5$ and arrange the coefficients into [Pascal's triangle](#)
- Recognise the equivalence between the coefficient of $x^{n-r}y^r$ in the expansion of $(x + y)^n$ and nC_r , when n is a positive integer
- Use patterns and symmetry in Pascal's triangle to confirm the identities ${}^nC_r = {}^{n-1}C_{r-1} + {}^{n-1}C_r$ for $1 \leq r \leq n - 1$ and ${}^nC_r = {}^nC_{n-r}$ for $0 \leq r \leq n$
- Derive the [binomial theorem](#):
 $(x + y)^n = {}^nC_0x^n + {}^nC_1x^{n-1}y + {}^nC_2x^{n-2}y^2 + \dots + {}^nC_{n-1}xy^{n-1} + {}^nC_ny^n$, when n is a positive integer
- Apply the binomial theorem to expand and simplify expressions of the form $(x + y)^n$
- Use the binomial theorem to determine the coefficient of a term with a specific power or the constant term in a binomial expansion
- Use the identities ${}^nC_0 = 1$, ${}^nC_n = 1$, ${}^nC_r = {}^{n-1}C_{r-1} + {}^{n-1}C_r$ for $1 \leq r \leq n - 1$ and ${}^nC_r = {}^nC_{n-r}$ for $0 \leq r \leq n$ to simplify expressions involving the [binomial coefficients](#)
- Prove identities involving binomial coefficients in binomial expansions by substituting values, comparing coefficients or applying a combinatorial argument to a specified context
- Apply given or proven identities involving binomial coefficients to prove further identities, without the use of calculus

Outcomes and content for Year 12

Proof by mathematical induction

Outcomes

A student:

- develops understanding and fluency in mathematics through exploring and connecting mathematical concepts, choosing and applying mathematical techniques to solve problems, and communicating their thinking and reasoning coherently and clearly **MAO-WM-01**
- uses mathematical induction to prove results involving sums and divisibility **ME1-12-01**

Content

- Examine the nature of inductive [proof](#), including the [statement](#) to be proved, the base case and the inductive step
- Prove results for sums using [mathematical induction](#)
- Prove divisibility results using mathematical induction
- Identify errors in false ‘proofs by induction’, such as cases where only one of the two required steps of a proof by induction is true

Introduction to vectors

Outcomes

A student:

- develops understanding and fluency in mathematics through exploring and connecting mathematical concepts, choosing and applying mathematical techniques to solve problems, and communicating their thinking and reasoning coherently and clearly **MAO-WM-01**
- operates with 2D and 3D vectors and uses 2D vectors to solve problems involving motion in two dimensions **ME1-12-02**

Content

Vector representation and notation

- Define a [vector](#) as a quantity having both [magnitude](#) and direction
- Associate vectors with [directed line segments](#) and recognise that a vector may have many directed line segments associated with it
- Identify and use notation for vectors in both two dimensions and three dimensions, including $\mathbf{a}$, $\vec{a}$ and $\overrightarrow{AB}$, where $\overrightarrow{AB}$ is the vector with magnitude and direction those of the directed line segment from A to B
- Use notations $|\mathbf{a}|$, $|\vec{a}|$ and $|\overrightarrow{AB}|$ to represent the magnitude of a vector
- Describe a [position vector](#) as a vector with its tail at the origin
- Represent vectors graphically with and without graphing applications
- Recognise and use the fact that two vectors are [equal](#) if they have the same magnitude and direction to solve problems

Introduction to 2D and 3D vectors

- Use [Cartesian coordinates](#) to represent points in 2-dimensional (2D) and 3-dimensional (3D) space with and without graphing applications
- Use the [midpoint](#) and distance formulas in two dimensions and three dimensions
- Identify that, in three dimensions, all points on the xy -plane have a z -coordinate of 0, and deduce similar properties for points on the xz and yz -planes and thus the equations of the three [coordinate planes](#)
- Define the [zero vector](#) $\mathbf{0}$, written as $\vec{0}$, as the vector with zero magnitude and no direction
- Define [unit vectors](#) as vectors of magnitude 1
- Recognise and use $\hat{\mathbf{a}}$ and $\vec{\hat{\mathbf{a}}}$ as the notation for the unit vector in the direction of $\mathbf{a}$
- Define the standard [perpendicular](#) unit vectors $\mathbf{i}$ and $\mathbf{j}$ in two dimensions and $\mathbf{i}$, $\mathbf{j}$ and $\mathbf{k}$ in three dimensions
- Express 2D vectors in [component form](#), $x\mathbf{i} + y\mathbf{j}$; as an ordered pair, (x,y) ; and in [column vector notation](#) $\begin{pmatrix} x \\ y \end{pmatrix}$
- Recognise $\overrightarrow{AB} = \begin{pmatrix} u-x \\ v-y \end{pmatrix}$ as the vector associated with the directed line segment from the point $A(x,y)$ to $B(u,v)$ in two dimensions

- Express 3D vectors in component form, $x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$; as an ordered triple, (x,y,z) ; and in column vector notation $\begin{pmatrix} x \\ y \\ z \end{pmatrix}$
- Recognise $\overrightarrow{AB} = \begin{pmatrix} u-x \\ v-y \\ w-z \end{pmatrix}$ as the vector associated with the directed line segment from $A(x,y,z)$ to $B(u,v,w)$ in three dimensions

Operating with vectors

- Define a [scalar](#) as a real number that is used to multiply a vector
- Represent geometrically a scalar multiple of a vector in two dimensions and three dimensions with and without graphing applications
- Perform multiplication of a vector by a scalar algebraically in component form
- Establish and identify $\mathbf{a} = k\mathbf{b}$, for a non-zero scalar k , as a condition for two non-zero vectors $\mathbf{a}$ and $\mathbf{b}$ to be [parallel](#) to each other and determine with justification if two vectors are parallel to one another
- Identify $\begin{pmatrix} b \\ -a \end{pmatrix}$ and $\begin{pmatrix} -b \\ a \end{pmatrix}$ as vectors perpendicular to $\begin{pmatrix} a \\ b \end{pmatrix}$ and with equal magnitude
- Perform addition and subtraction of vectors algebraically in component form, and verify, with and without graphing applications, that geometrically these are obtained using the [triangle law](#) or the [parallelogram law](#)
- Establish and calculate the magnitude of a vector using $|x\mathbf{i} + y\mathbf{j}| = \sqrt{x^2 + y^2}$ for 2D vectors and $|x\mathbf{i} + y\mathbf{j} + z\mathbf{k}| = \sqrt{x^2 + y^2 + z^2}$ for 3D vectors
- Use the magnitude of a vector to find the unit vector $\hat{\mathbf{a}} = \frac{1}{|\mathbf{a}|}\mathbf{a}$ in two dimensions and three dimensions

Further operations with vectors

- Define $\mathbf{a} \cdot \mathbf{b} = x_1x_2 + y_1y_2$ as the [scalar \(dot\) product](#) of vectors $\mathbf{a} = x_1\mathbf{i} + y_1\mathbf{j}$ and $\mathbf{b} = x_2\mathbf{i} + y_2\mathbf{j}$ and use the scalar product to solve problems
- Define $\mathbf{a} \cdot \mathbf{b} = x_1x_2 + y_1y_2 + z_1z_2$ as the scalar product of vectors $\mathbf{a} = x_1\mathbf{i} + y_1\mathbf{j} + z_1\mathbf{k}$ and $\mathbf{b} = x_2\mathbf{i} + y_2\mathbf{j} + z_2\mathbf{k}$ and use the scalar product to solve problems
- Use $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}|\cos \theta$ as a geometric expression of the scalar product of non-zero vectors $\mathbf{a}$ and $\mathbf{b}$ in two dimensions and three dimensions, where θ is the angle between the vectors and $0^\circ \leq \theta \leq 180^\circ$.
- Verify the equivalence of $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}|\cos \theta$ with the algebraic definition of the scalar product, $\mathbf{a} \cdot \mathbf{b} = x_1x_2 + y_1y_2$ for two dimensions and $\mathbf{a} \cdot \mathbf{b} = x_1x_2 + y_1y_2 + z_1z_2$ for three dimensions
- Derive and use the property $\mathbf{a} \cdot \mathbf{a} = |\mathbf{a}|^2$ to establish the scalar product definition of the magnitude of a vector ($|\mathbf{a}| = \sqrt{\mathbf{a} \cdot \mathbf{a}}$) in two dimensions and three dimensions
- Calculate the angle between two non-zero vectors $\mathbf{a}$ and $\mathbf{b}$, in both two dimensions and three dimensions, using the scalar product by deriving and applying the relationship $\cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|} = (\hat{\mathbf{a}} \cdot \hat{\mathbf{b}})$
- Establish $\mathbf{a} \cdot \mathbf{b} = 0$ as a condition for two non-zero vectors $\mathbf{a}$ and $\mathbf{b}$ to be perpendicular to each other and use it to determine if two vectors are perpendicular
- Establish $\mathbf{a} \cdot \mathbf{b} = \pm |\mathbf{a}||\mathbf{b}|$ as another way to determine if two non-zero vectors $\mathbf{a}$ and $\mathbf{b}$ are parallel

- Define the [projection](#) of a vector $\mathbf{a}$ onto a vector $\mathbf{b}$, denoted by $\text{proj}_{\mathbf{b}}\mathbf{a}$, to be the vector [component](#) of $\mathbf{a}$ in the direction of vector $\mathbf{b}$
- Examine the proof of the formula $\text{proj}_{\mathbf{b}}\mathbf{a} = \left(\frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{b}|^2}\right)\mathbf{b} = (\mathbf{a} \cdot \hat{\mathbf{b}})\hat{\mathbf{b}}$ and use the formula to solve problems
- Determine that the component of a vector $\mathbf{a}$ perpendicular to another vector $\mathbf{b}$ is $\mathbf{a} - \text{proj}_{\mathbf{b}}\mathbf{a}$

Motion in vector form in two dimensions

- Describe the position of an object at a point in 2D space using a vector
- Describe the changing positions of an object by expressing its vector as a function of time using $\mathbf{r}(t) = x(t)\mathbf{i} + y(t)\mathbf{j}$, or $\mathbf{r} = x\mathbf{i} + y\mathbf{j}$ where x and y are functions of time t
- Recognise that $x(t)$ and $y(t)$ form a pair of parametric equations for the path of the object
- Find the Cartesian equation of the path of an object, where the path is a straight line, [parabola](#) or circle
- Express the change in an object's position between two points as a [displacement vector](#) and recognise the magnitude of the displacement vector as the distance between the two points
- Solve motion problems involving constant [velocity](#) using vectors
- Solve relative velocity problems involving constant crosswind/cross-current using vector diagrams, and describe the direction of a vector where required
- Find the velocity vector $\mathbf{v} = \dot{x}\mathbf{i} + \dot{y}\mathbf{j}$ and [acceleration vector](#) $\mathbf{a} = \ddot{x}\mathbf{i} + \ddot{y}\mathbf{j}$ of an object using differential calculus
- Find the position vector and the velocity vector of an object using integral calculus given its acceleration vector
- Solve motion problems involving non-constant velocity using vectors

Projectile motion

- Recognise that the [gravitational force](#) on a [mass](#) may be regarded as a constant acting in a downwards direction when the motion of the object is restricted to a small region near the Earth's surface
- Model and analyse a [projectile's](#) path where the projectile is a point and air resistance is negligible, subject to only [acceleration](#) due to gravity, assuming that the projectile is moving close to the Earth's surface
- Represent the motion of a projectile using vectors
- Recognise that the horizontal and vertical components of the motion of a projectile can be represented by horizontal and vertical vectors
- Derive and use the equations of motion of a projectile in vector form by splitting 2D motion into horizontal and vertical components to solve problems on projectiles
- Find the Cartesian equation of the path of a projectile using parametric equations for the horizontal and vertical components of the displacement vector
- Determine features of the flight of a projectile, including time of flight, maximum height, range, instantaneous velocity and impact velocity
- Solve problems relating to the path of a projectile in which the initial velocity and/or angle of projection may be unknown, in a variety of contexts

Inverse trigonometric functions

Outcomes

A student:

- develops understanding and fluency in mathematics through exploring and connecting mathematical concepts, choosing and applying mathematical techniques to solve problems, and communicating their thinking and reasoning coherently and clearly **MAO-WM-01**
- solves problems involving inverse trigonometric functions **ME1-12-03**

Content

Definitions of inverse trigonometric functions

- Graph $y = \sin x$, $y = \cos x$ and $y = \tan x$, for $-2\pi \leq x \leq 2\pi$ using graphing applications and recognise that these three functions fail the horizontal line test
- Examine possible domain restrictions of $y = \sin x$, $y = \cos x$ and $y = \tan x$ to obtain the corresponding inverse functions using graphing applications
- Define $\sin^{-1}x$ or $\arcsin x$ to be the inverse function of $y = \sin x$ restricted to $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$, and determine the domain and range of $y = \sin^{-1}x$
- Define $\cos^{-1}x$ or $\arccos x$ to be the inverse function of $y = \cos x$ restricted to $0 \leq x \leq \pi$, and determine the domain and range of $y = \cos^{-1}x$
- Define $\tan^{-1}x$ or $\arctan x$ to be the inverse function of $y = \tan x$ restricted to $-\frac{\pi}{2} < x < \frac{\pi}{2}$, and determine the domain and range of $y = \tan^{-1}x$
- Evaluate and simplify expressions, and prove results using the definitions of $\sin^{-1}x$, $\cos^{-1}x$ and $\tan^{-1}x$

Graphs of inverse trigonometric functions

- Graph $y = \sin x$ for $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$, then use a reflection in $y = x$ to graph $y = \sin^{-1}x$ with and without graphing applications
- Graph $y = \cos x$ for $0 \leq x \leq \pi$, then use a reflection in $y = x$ to graph $y = \cos^{-1}x$ with and without graphing applications
- Graph $y = \tan x$ for $-\frac{\pi}{2} < x < \frac{\pi}{2}$, then use a reflection in $y = x$ to graph $y = \tan^{-1}x$ with and without graphing applications
- Classify $y = \sin^{-1}x$, $y = \cos^{-1}x$ and $y = \tan^{-1}x$ as odd, even, or neither odd nor even
- Examine the properties $\sin^{-1}(-x) = -\sin^{-1}x$, $\cos^{-1}(-x) = \pi - \cos^{-1}x$, $\tan^{-1}(-x) = -\tan^{-1}x$ and $\cos^{-1}x + \sin^{-1}x = \frac{\pi}{2}$ using graphing applications and prove them algebraically
- Use the properties $\sin^{-1}(-x) = -\sin^{-1}x$, $\cos^{-1}(-x) = \pi - \cos^{-1}x$, $\tan^{-1}(-x) = -\tan^{-1}x$ and $\cos^{-1}x + \sin^{-1}x = \frac{\pi}{2}$ to solve problems, simplify expressions and prove results
- Graph the [composite functions](#) $y = \sin(\sin^{-1}x)$, $y = \cos(\cos^{-1}x)$ and $y = \tan(\tan^{-1}x)$ by considering their domains and ranges
- Examine the graphs of the composite functions $y = \sin(\sin^{-1}x)$, $y = \cos(\cos^{-1}x)$ and $y = \tan(\tan^{-1}x)$, establish the values of x for which the formulas $\sin(\sin^{-1}x) = x$, $\cos(\cos^{-1}x) = x$ and $\tan(\tan^{-1}x) = x$ are true, and apply the formulas to solve problems, simplify expressions and prove results

- Graph the composite functions $y = \sin^{-1}(\sin x)$, $y = \cos^{-1}(\cos x)$ and $y = \tan^{-1}(\tan x)$ by considering their domains and ranges
- Examine the graphs of the composite functions $y = \sin^{-1}(\sin x)$, $y = \cos^{-1}(\cos x)$ and $y = \tan^{-1}(\tan x)$, establish the values of x for which the formulas $\sin^{-1}(\sin x) = x$, $\cos^{-1}(\cos x) = x$ and $\tan^{-1}(\tan x) = x$ are true, and apply the formulas to solve problems, simplify expressions and prove results
- Use graphing applications to explore reflections, [translations](#) and [dilations](#) of the functions $f(x) = \sin^{-1} x$, $f(x) = \cos^{-1} x$ and $f(x) = \tan^{-1} x$, and confirm that the principles of [transformations](#) hold for the inverse trigonometric functions
- Apply the principles of transformations to inverse trigonometric functions to determine the function rule and graph the function, finding the domain and range of the transformed graph and determining any [intercepts](#) and asymptotes where appropriate

Further calculus skills

Outcomes

A student:

- develops understanding and fluency in mathematics through exploring and connecting mathematical concepts, choosing and applying mathematical techniques to solve problems, and communicating their thinking and reasoning coherently and clearly **MAO-WM-01**
- selects and applies differentiation and integration techniques to solve problems **ME1-12-04**

Content

Further derivatives of functions

- Find the [derivative](#) of a function defined parametrically using the [chain rule](#)
- Solve problems involving derivatives of functions defined parametrically
- Verify using the chain rule that the derivative of the inverse function is the reciprocal of the derivative of the function, evaluated at the value of the inverse function, that is
$$(f^{-1})'(x) = \frac{1}{f'(f^{-1}(x))}$$
- Solve problems involving derivatives of inverse functions
- Examine the proofs of the derivatives of $\sin^{-1} x$, $\cos^{-1} x$ and $\tan^{-1} x$
- Use the chain rule to show that $\frac{d}{dx}[\sin^{-1} f(x)] = \frac{f'(x)}{\sqrt{1-[f(x)]^2}}$,
$$\frac{d}{dx}[\cos^{-1} f(x)] = -\frac{f'(x)}{\sqrt{1-[f(x)]^2}} = -\frac{d}{dx}[\sin^{-1} f(x)] \text{ and } \frac{d}{dx}[\tan^{-1} f(x)] = \frac{f'(x)}{1+[f(x)]^2}$$
 and apply the results to solve problems involving derivatives of $\sin^{-1} f(x)$, $\cos^{-1} f(x)$ and $\tan^{-1} f(x)$
- Apply the product, [quotient](#) and chain rules to find derivatives of functions of the form $f(x)g(x)$, $\frac{f(x)}{g(x)}$ and $f(g(x))$ where $f(x)$ and $g(x)$ are any of the functions covered in the scope of the *Mathematics Advanced 11–12 Syllabus* (2024) or inverse trigonometric functions and solve related problems

Techniques of integration

- Find indefinite and definite integrals involving expressions of the form $\frac{1}{\sqrt{a^2-x^2}}$ or $\frac{a}{a^2+x^2}$
- Use [integration](#) by substitution to evaluate definite and [indefinite integrals](#) given the substitution, where the substitution is expressed as a function of the variable of integration or where the variable of integration is the subject of the substitution
- Prove and use the identities $\sin^2 nx = \frac{1}{2}(1 - \cos 2nx)$ and $\cos^2 nx = \frac{1}{2}(1 + \cos 2nx)$ to find integrals involving $\int \sin^2 nx \, dx$ and $\int \cos^2 nx \, dx$

Further applications of calculus

Outcomes

A student:

- develops understanding and fluency in mathematics through exploring and connecting mathematical concepts, choosing and applying mathematical techniques to solve problems, and communicating their thinking and reasoning coherently and clearly **MAO-WM-01**
- applies calculus to solve problems involving polynomials, further rates of change, areas and volumes and differential equations **ME1-12-05**

Content

Multiplicity of zeroes of polynomial functions

- Use the [product rule](#) to prove that if $P(x)$ has a zero, α , of multiplicity $m > 1$, then α is a zero of $P'(x)$ of multiplicity $m - 1$, and use the result to determine the multiplicity of a discovered zero of $P(x)$ and solve related polynomial problems
- Prove that if α is a zero of multiplicity m , meeting the x -axis at $A(\alpha, 0)$, then if $m = 1$, the curve crosses the x -axis at A at an [acute](#) or [obtuse angle](#); if $m > 1$ is even, the curve is [tangent](#) to the x -axis at A and does not cross it; if $m > 1$ is odd, the curve has a horizontal inflection at A
- Graph a polynomial function in factored form, identifying its [turning points](#) and [points of inflection](#) if possible, and explore its behaviour as $x \rightarrow \infty$ and $x \rightarrow -\infty$, verifying the shape of the graph using graphing applications

Further rates of change

- Develop models in contexts where a [rate of change](#) of a function can be expressed as a rate of change of a composition of two functions, so that the chain rule can be applied
- Solve problems involving related rates of change using the chain rule, given the required formulas for problems relating to area, surface area or [volume](#)
- Describe and examine the graphs of practical situations where the rate of change of a quantity is proportional to the amount $Q - P$ by which the quantity Q exceeds some fixed value P
- Explain that if the rate of change of a quantity Q over time t is proportional to the difference $Q - P$ at any instant, then this may be represented by the equation $\frac{dQ}{dt} = k(Q - P)$, where k is a constant
- Verify by substitution that the function $Q = P + Ae^{kt}$, where A is a constant, satisfies the relationship $\frac{dQ}{dt} = k(Q - P)$, and that $Q = P$ in the case where $A = 0$
- Graph the function $Q = P + Ae^{kt}$, where $k > 0$ and $k < 0$ and $A > 0$ and $A < 0$, with and without graphing applications, and identify any asymptotes
- Use $\frac{dQ}{dt} = k(Q - P)$, $Q = P + Ae^{kt}$ and the graph of $Q = P + Ae^{kt}$ for $t \geq 0$, where $k > 0$ or $k < 0$ and $A > 0$ or $A < 0$, to model and solve problems where a limiting value of Q exists, including Newton's Law of Cooling and ecosystems with a natural carrying capacity, and justify conclusions in the context of the problem

Areas between curves and volumes of solids of revolution

- Calculate areas of regions between curves determined by functions in both real-life and abstract contexts

- Examine a solid of revolution whose boundary is formed by rotating an [arc](#) of a function about the x -axis or y -axis with and without graphing applications
- Calculate the volume of a solid of revolution formed by rotating a region in the plane about the x -axis or y -axis in both real-life and abstract contexts
- Calculate the volume of a solid of revolution formed by rotating the region between two curves about either the x -axis or y -axis in both real-life and abstract contexts

Differential equations

- Define a [differential equation](#) as an equation involving an unknown function and one or more of its derivatives
- Define and identify the order of a differential equation as the order of the highest derivative contained within the equation
- Recognise that a solution to a first order differential equations is a function, and that there may be [infinitely](#) many functions that are solutions to a given first order differential equation
- Recognise the solutions to differential equations in the context of [slope fields](#), and that slope fields are useful in determining the behaviour of solutions when the differential equation cannot be easily solved
- Recognise that a unique solution of a differential equation can be determined when sufficient initial conditions are given, and refer to a problem involving a differential equation and initial conditions as an initial value problem (IVP)
- Graph solutions to first order differential equations given a slope field and identify the unique solution curve that satisfies a set of initial conditions
- Explore problems given a slope field representing a practical context and justify conclusions
- Form a slope field for a first order differential equation using graphing applications
- Recognise the features of a slope field corresponding to a first order differential equation and vice versa
- Solve first order differential equations of the form $\frac{dy}{dx} = f(x)$
- Solve first order differential equations of the form $\frac{dy}{dx} = g(y)$, where possible expressing the solution as a function with y as the subject
- Recognise and solve the first order differential equations for exponential growth and decay: $\frac{dQ}{dt} = kQ$ and $\frac{dQ}{dt} = k(Q - P)$
- Solve first order differential equations of the form $\frac{dy}{dx} = f(x)g(y)$ using separation of variables, where possible expressing the solution as a function with y as the subject
- Graph solutions of first order differential equations using graphing applications and examine the behaviour of solutions for different values of the constant of integration and initial conditions
- Solve differential equations of the form $\frac{dP}{dt} = kP\left(1 - \frac{P}{C}\right)$ for some constants k and C , given the appropriate decomposition into [partial fractions](#), to obtain the [logistic function](#)
- Model and solve differential equations in practical scenarios including in chemistry, biology and economics

The binomial distribution and sampling distribution of the mean

Outcomes

A student:

- develops understanding and fluency in mathematics through exploring and connecting mathematical concepts, choosing and applying mathematical techniques to solve problems, and communicating their thinking and reasoning coherently and clearly **MAO-WM-01**
- solves problems involving binomial distributions, sampling distribution of the mean and the central limit theorem **ME1-12-06**

Content

Bernoulli distributions

- Define a [Bernoulli random variable](#) as a model for two-[outcome](#) situations, referred to as success and failure
- Model a [Bernoulli trial](#) using a [random variable](#), X , where the probability of success, $X = 1$, is p and the probability of failure, $X = 0$, is $q = 1 - p$
- Define a [Bernoulli distribution](#) as the [discrete probability distribution](#) of a Bernoulli random variable
- Establish and use the formulae $E(X) = \mu = p$ to find the [mean](#), and $\text{Var}(X) = p(1 - p)$ to find the [variance](#) of the Bernoulli distribution with parameter p
- Identify contexts suitable to be modelled by Bernoulli random variables
- Solve practical problems involving Bernoulli random variables

Binomial distributions

- Define a [binomial random variable](#), X , as the number of 'successes' in n independent and identically distributed Bernoulli trials, with the same probability of success p in each [trial](#), where n is a fixed [finite](#) number
- Identify contexts suitable to be modelled using [binomial distributions](#)
- Define a binomial experiment as a fixed number of Bernoulli trials
- Use the notation $X \sim \text{Bin}(n, p)$ to represent a binomial random variable, X , where n is the number of trials and probability of success is p
- Recognise the significance of nC_r as the number of ways in which an outcome with r successes can occur in n trials
- Use the formula $P(X = r) = {}^nC_r p^r (1 - p)^{n-r}$ to find the probability of r successes in a binomial experiment involving n trials
- Calculate the expected frequencies of the various possible outcomes from a binomial experiment, where the [expected frequency](#) of an outcome happening r times in n trials is $n \times P(X = r)$
- Use the formulae $E(X) = \mu = np$ to find the mean, and $\text{Var}(X) = np(1 - p)$ to find the variance when $X \sim \text{Bin}(n, p)$
- Solve practical problems involving binomial distributions and binomial probabilities with and without online computational applications, excluding the normal approximation to the binomial distribution

Sampling distribution of the mean and the central limit theorem

- Define a statistical [population](#) as the entire group of people or objects about which information is sought
- Define a [sample](#) as a selection of people or objects drawn from a population
- Recognise that a statistic taken from a sample will be a good approximation if the sample is random and the sample size is large enough, typically $n \geq 30$
- Recognise that, in general, the distribution of a population and a statistic, such as its mean, are unknown
- Recognise that it is likely that the distribution of a random sample will resemble the distribution of the population when the sample size, n , is large enough, typically $n \geq 30$
- Define the [sample mean](#) for a sample $X_1, X_2, \dots, X_n$ taken from a population as $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$
- Recognise that the sample means obtained from repeated sampling may be different, even if all the samples are of the same size
- Define, for a given sample size n , the [sampling distribution](#) of the mean to be the distribution of the sample means of all samples of size n
- Recognise that sample mean, $\bar{X}$, is a random [variable](#) that estimates the population mean, μ , when the size of the sample is large enough and consider its usefulness in making predictions in a variety of contexts including politics, finance, agriculture and biology
- State the central limit theorem as: for a population with mean μ and variance σ^2 , provided the sample size n is large enough ($n \geq 30$), the sampling distribution is approximately normal with mean μ and variance $\frac{\sigma^2}{n}$; that is, $\bar{X}$ is approximately $N\left(\mu, \frac{\sigma^2}{n}\right)$
- Recognise the significance of the central limit theorem for populations that are not necessarily normally distributed; that, irrespective of the population distribution, for a sufficiently large sample size, the sampling distribution of the mean is approximately normal
- Use the formulas $E(\bar{X}) = \mu$ and $\text{Var}(\bar{X}) = \frac{\sigma^2}{n}$ as the mean and the variance of the sampling distribution of the mean for samples of size n drawn from a population with mean μ and variance σ^2
- Examine the effect of the sample size on the variance of sample means with digital tools
- Apply the central limit theorem to estimate the probability that the sample mean lies within given bounds