

1 Basic notions

1.2 Graph isomorphism

Def 1.6. Let $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ be graphs. An isomorphism $\phi : G_1 \rightarrow G_2$ is a bijection (a one-to-one correspondence) from V_1 to V_2 such that $(u, v) \in E_1$ iff $(\phi(u), \phi(v)) \in E_2$. We say G_1 is isomorphic to G_2 if there is an isomorphism between them.

Rem 1.8. Isomorphism is an equivalence relation of graphs. (reflexive, symmetric, transitive)

Def 1.9. An unlabelled gr. is an isomorphism class of graphs.

1.3 The adjacency and incidence matrices

Let $[n] = \{1, \dots, n\}$.

Def 1.10. Let $G = (V, E)$ be a gr. with $V = [n]$. The adjacency matrix $A = A(G)$ is the gr. with $V = [n]$. The adjacency matrix $A = A(G)$ is the $n \times n$ symmetric matrix defined by

$$a_{ij} = \begin{cases} 1 & \text{if } (i, j) \in E \\ 0 & \text{otherwise} \end{cases}$$

Def 1.13. Let $G = (V, E)$ be a gr. with $V = \{v_1, \dots, v_n\}$ and $E = \{e_1, \dots, e_m\}$. Then the incidence matrix $B = B(G)$ of G is the $n \times m$ matrix defined by

$$b_{ij} = \begin{cases} 1 & \text{if } v_i \in e_j \\ 0 & \text{otherwise} \end{cases}$$

Rem 1.15. Every column of B has $|e| = 2$ entries 1.

1.4 Degree

Fact 1. For any G on the vx set $[n]$ with adjacency and incidence matrices A and B , we have $BB^T = D + A$, where

$$D = \begin{pmatrix} d(1) & 0 & 0 \\ 0 & \ddots & 0 \\ 0 & 0 & d(n) \end{pmatrix}$$

Def 1.20. A G is d -regular iff all vxs have degree d .

Prop 1.22. For every $G = (V, E)$, $\sum_{v \in G} d(G) = 2|E|$

Cor 1.23. Every gr. has an even # vxs of odd degree.

1.5 Subgraphs

Def 1.24. A gr. $H = (U, F)$ is a subgr. of a gr. $G = (V, E)$ if $U \subseteq V$ and $F \subseteq E$. If $U = V$ then H is called spanning.

Def 1.25. Given $G = (V, E)$ and $U \subseteq V (U \neq \emptyset)$, let $G[U]$

denote the gr. with vx set U and edge set $E(G[U]) = \{e \in E(G) : e \subseteq U\}$. (We include all the edges of G which have both endpoints in U). Then $G[U]$ is called the subgr. of G induced by U .

1.7 Walks, paths and cycles

Def 1.29. A walk in G is a sequence of vxs v_0, v_1, \dots, v_k , and a sequence of edges $(v_i, v_{i+1}) \in E(G)$. A walk is a path if all v_i are distinct. If for such a path with $k \geq 2$, (v_0, v_k) is also an edge in G , then $v_0, v_1, \dots, v_k, v_0$ is a cycle. For multigraphs, we also consider loops and pairs of multiple edges to be cycles.

Def 1.30. The length of a path, cycle or walk is the # edges in it.

Prop 1.32. Every walk from u to v in G contains a path between u and v .

Prop 1.33. Every G with minimum degree $\delta \geq 2$ contains a path of length δ and a cycle of length at least $\delta + 1$.

Rem 1.34. Note that we have also proved that a gr. with minimum degree $\delta \geq 2$ contains cycles of at least $\delta - 1$

different lengths. This fact, and the statement of Proposition 1.32, are both tight, to see this, consider the complete gr. $G = K_{\delta+1}$.

1.8 Connectivity

Def 1.35. A G is conn. if for all pairs $u, v \in G$, there is a path in G from u to v .

Note that it suffices for there to be a walk from u to v , by Proposition 1.31.

Def 1.37. A (conn.) component of G is a conn. subgr. that is maximal by inclusion. We say G is conn. iff it has one conn. component.

Prop 1.39. A gr. with n vxs and m edges has at least $n - m$ conn. components.

1.9 Graph operations and parameters

Def 1.40. Given $G = (V, E)$, the complement \overline{G} of G has the same vx set V and $(u, v) \in E(\overline{G})$ iff $(u, v) \notin E(G)$.

Def 1.42. A clique in G is a complete subgr. in G . An independent set is an empty induced subgr. in G .

Not 1.44. Let $\omega(G)$ denote the # vxs in a maximum-size

clique in G , let $\alpha(G)$ denote the # vxs in a maximum-size independent set in G .

Claim 1.45. A vx set $U \subseteq V(G)$ is a clique iff $U \subseteq V(\overline{G})$ is an independent set.

Cor 1.46. We have $\omega(G) = \alpha(\overline{G})$ and $\alpha(G) = \omega(\overline{G})$.

2 Trees

2.1 Trees

Def 2.1. A gr. having no cycle is acyclic. A forest is an acyclic graph, a tree is a conn. acyclic graph. A leaf is a vx of degree 1.

Lem 2.3. Every finite tree with at least two vxs has at least two leaves. Deleting a leaf from an n -vx tree produces a tree with $n - 1$ vxs.

2.2 Equivalent definitions of trees

Thm 2.4. For an n -vx simple G (with $n \geq 1$), the following are equivalent (and characterize the trees with n vxs). (a) G is conn. and has no cycles. (b) G is conn. and has $n - 1$ edges. (c) G has $n - 1$ edges and no cycles. (d) For every pair $u, v \in V(G)$, there is exactly one u, v -path in G .

Def 2.5. An edge of a gr. is a cut-edge if its deletion disconnects the graph.

Lem 2.6. An edge contained in a cycle is not a cut-edge.

Cor 2.8. • Every conn. gr. on n vxs has at least $n - 1$ edges and contains a spanning tree,

- Every edge of a tree is a cut-edge,
- Adding an edge to a tree creates exactly one cycle.

2.3 Cayley's formula

Thm 2.11 (Cayley's Formula). There are n^{n-2} trees with vx set $[n]$.

Def 2.12 (Prüfer code). Let T be a tree on an ordered set S of n vxs. To compute the Prüfer sequence $f(T)$, iteratively delete the leaf with the smallest label and append the label of its neighbour to the sequence. After $n - 2$ iterations a single edge remains and we have produced a sequence $f(T)$ of length $n - 2$.

Prop 2.14. For an ordered n -element set S , the Prüfer code f is a bijection between the trees with vx set S and the sequences in S^{n-2} .

3 Connectivity

3.1 Vertex connectivity

Def 3.1. A vx cut in a conn. gr. $G = (V, E)$ is a set $S \subseteq V$ such that $G \setminus S := G[V \setminus S]$ has more than one conn. component. A cut vx is a vx v such that $\{v\}$ is a cut.

Def 3.2. G is called k -conn. if $|V(G)| > k$ and if $G \setminus X$ is conn. for every set $X \subseteq V$ with $|X| < k$. In other words, no two vxs of G are separated by fewer than k other vxs. Every (non-empty) gr. is 0-conn. and the 1-conn. graphs are precisely the non-trivial conn. graphs. The greatest integer k such that G is k -conn. is the connectivity $\kappa(G)$ of G . 17

$$G = K_n : \kappa(G) = n - 1$$

$G = K_{m,n}, m \leq n : \kappa(G) = m$. Indeed, let G have bipartition $A \cup B$, with $|A| = m$ and $|B| = n$. Deleting A disconnects the graph. On the other hand, deleting $S \subset V$ with $|S| < m$ leaves both $A \setminus S$ and $B \setminus S$ non-empty and any $a \in A \setminus S$ is conn. to any $b \in B \setminus S$. Hence $G \setminus S$ is conn.

Prop 3.3. For every G , $\kappa(G) \leq \delta(G)$.

Rem 3.4. High minimum degree does not imply connectivity. Consider two disjoint copies of K_n .

Thm 3.5 (Mader 1972). Every gr. of average degree at least $4k$ has a k -conn. subgraph.

3.2 Edge connectivity

Def 3.6. A disconnecting set of edges is a set $F \subseteq E(G)$ such that $G \setminus F$ has more than one component. Given $S, T \subset V(G)$, the notation $[S, T]$ specifies the set of edges having one endpoint in S and the other in T . An edge cut is an edge set of the form $[S, S]$, where S is a non-empty proper subset of $V(G)$. A gr. is k -edge-conn. if every disconnecting set has at least k edges. The edge-connectivity of G , written $\kappa'(G)$, is the minimum size of a disconnecting set. One edge disconnecting G is called a bridge. $G = K_n : \kappa'(G) = n - 1$.

Rem 3.8. An edge cut is a disconnecting set but not the other way around. However,

every minimal disconnecting set is a cut.

Thm 3.9. $\kappa(G) \leq \kappa'(G) \leq \delta(G)$.

3.3 Blocks

Def 3.10. A block of a G is a maximal conn. subgr. of G that has no cut-vx. If G itself is conn. and has no cut-vx, then G is a block.

Rem 3.12. If a block B has at least three vxs, then B is 2-conn. If an edge is a block of G then it is a cut-edge of G .

Prop 3.13. Two blocks in a gr. share at most one vx.

Def 3.14. The block gr. of a G is a bipartite gr. H in which one partite set consists of the cut-vxs of G , and the other has a vx b_i for each block B_i of G . We include (v, b_i) as an edge of H iff $v \in B_i$.

Prop 3.16. The block gr. of a conn. gr. is a tree.

3.4 2-conn. graphs

Def 3.17. Two paths are internally disjoint if neither contains a non-endpoint vx of the other. We denote the length of the shortest path

from u to v (the distance from u to v) by $d(u, v)$.

Thm 3.18 (Whitney 1932). A G having at least three vxs is 2-conn. if and only if each pair $u, v \in V(G)$ is conn. by a pair of internally disjoint u, v -paths in G .

Cor 3.19. G is 2-conn. and $|G| \geq 3$ iff every two vxs in G lie on a common cycle.

3.5 Menger's Thm

Def 3.20. Let $A, B \subseteq V$. An $A - B$ path is a path with one endpoint in A , the other endpoint in B , and all interior vxs outside of $A \cup B$. Any vx in $A - B$ is a trivial $A - B$ path.

If $X \subseteq V$ (or $X \subseteq E$) is such that every $A - B$ path in G contains a vx (or an edge) from X , we say that X separates the sets A and B in G . This implies in particular that $A \cap B \subseteq X$.

Thm 3.21 (Menger 1927). Let $G = (V, E)$ be a gr. and let $S, T \subseteq V$. Then the maximum # vx-disjoint $S - T$ paths is equal to the minimum size of an $S - T$ separating vx set.

Cor 3.22. For $S \subseteq V$ and $v \in V \setminus S$, the minimum # vxs distinct from v separating v from S in G is equal to the maximum # paths forming an $v - S$ fan in G . (that is, the maximum # $\{v\} - S$ paths which are disjoint except at v).

Def 3.23. The line gr. of G , written $L(G)$, is the gr. whose vxs are the edges of G , with $(e, f) \in E(L(G))$ when $e = (u, v)$ and $f = (v, w)$ in G (i.e. when e and f share a vx).

Cor 3.25. Let u and v be two distinct vxs of G .

1. If $(u, v) \notin E$, then the minimum # vxs different from u, v separating u from v in G is equal to the maximum # internally vx-disjoint $u - v$ paths in G .
2. The minimum # edges separating u from v in G is equal to the maximum # edge-disjoint $u - v$ paths in G .

Thm 3.26. (Global Version of Menger's Theorem)

1. A gr. is k -conn. iff it contains k internally vx-

disjoint paths between any two vxs.

2. A gr. is k -edge-conn. iff it contains k edge-disjoint paths between any two vxs.

4 Eu. & Ha. cyc.

4.1 Eul. trails & tours

Def 4.2. A trail is a walk with no repeated edges.

Def 4.3. An Eulerian trail in a (multi)gr. $G = (V, E)$ is a walk in G passing through every edge exactly once. If this walk is closed (starts and ends at the same vx) it is called an Eulerian tour.

Thm 4.5. A conn. (multi)gr. has an Eulerian tour iff each vx has even degree.

Lem 4.6. Every maximal trail in an even gr. (i.e., a gr. where all the vxs have even degree) is a closed trail.

Cor 4.7. A conn. multi G has an Eulerian trail iff it has either 0 or 2 vxs of odd degree.

4.2 Hamilton paths and cycles

Def 4.8. A Hamilton path/-cycle in a G is a path/cycle visiting every vx of G exactly

once. A G is called Hamiltonian if it contains a Hamilton cycle.

Prop 4.10. If G is Hamiltonian then for any set $S \subseteq V$ the gr. $G \setminus S$ has at most $|S|$ conn. components.

Cor 4.11. If a conn. bipartite gr. $G = (V, E)$ with bipartition $V = A \cup B$ is Hamiltonian then $|A| = |B|$.

Thm 4.13 (Dirac 1952). If G is a simple gr. with $n \geq 3$ vxs and if $\delta(G) \geq n/2$, then G is Hamiltonian.

Thm 4.15 (Ore 1960). If G is a simple gr. with $n \geq 3$ vxs such that for every pair of non-adjacent vxs u, v of G we have $d(u) + d(v) \geq |G|$, then G is Hamiltonian.

5 Matchings

Def 5.1. A set of edges $M \subseteq E(G)$ in a G is called a matching if $e \cap e' = \emptyset$ for any pair of edges $e, e' \in M$.

A matching is perfect if $|M| = \frac{|V(G)|}{2}$, i.e. it covers all vxs of G . We denote the size of the maximum matching in G , by $\nu(G)$.

$$G = K_n; \nu(G) = \lfloor \frac{n}{2} \rfloor$$

$$G = K_{s,t}; s \leq t, \nu(G) = s$$

$$\nu(\text{PetersenGraph}) = 5$$

Rem 5.3. A matching in a G corresponds to an independent set in the line gr. $L(G)$.

Def 5.4. A set of vxs $T \subseteq V(G)$ of a gr. G is called a cover of G if every edge $e \in E(G)$ intersects T ($e \cap T \neq \emptyset$), i.e., $G \setminus T$ is an empty graph. Then, $\tau(G)$ denotes the size of the minimum cover.

$$G = K_n : \tau(G) = n - 1$$

$$G = K_{s,t}, s \leq t : \tau(G) = s$$

$$\tau(\text{PetersenGraph}) = 6$$

Prop 5.6. $\nu(G) \leq \tau(G) \leq 2\nu(G)$.

5.2 Hall's Theorem

Thm 5.7 (Hall 1935). A bipartite gr. $G = (V, E)$ with bipartition $V = A \cup B$ has a matching covering A iff $|N(S)| \geq |S| \forall S \subseteq A$

Cor 5.8. If in a bipartite gr. $G = (A \cup B, E)$ we have $|N(S)| \geq |S| - d$ for every set $S \subseteq A$ and some fixed $d \in \mathbb{N}$, then G contains a matching of cardinality $|A| - d$.

Cor 5.9. If a bipartite gr. $G = (A \cup B, E)$ is k -regular with $k \geq 1$, then G has a perfect matching.

Cor 5.10. Every regular gr. of positive even degree has a 2-factor (a spanning 2-regular subgraph).

Rem 5.11. A 2-factor is a disjoint union of cycles covering all the vxs of a graph

Def 5.12. Let A_1, \dots, A_n be a collection of sets. A family $\{a_1, \dots, a_n\}$ is called a system of distinct representatives (SDR) if all the a_i are distinct, and $a_i \in A_i$ for all i .

Cor 5.13. A collection A_1, \dots, A_n has an SDR iff for all $I \subseteq [n]$ we have $|\bigcup_{i \in I} A_i| \geq |I|$.

Thm 5.15 (König 1931). If $G = (A \cup B, E)$ is a bipartite graph, then the maximum size of a matching in G equals the minimum size of a vx cover of G .

5.3 Matchings in general graphs: Tutte's Theorem

Given a G , let $q(G)$ denote the # its odd components, i.e. the ones of odd order. If G has a perfect matching then clearly $q(G \setminus S) \leq |S|$ for all $S \subseteq V(G)$ since every odd component of

GnS will send an edge of the matching to S, and each such edge covers a different vx in S.

Thm 5.16 (Tutte 1947). *A G has a perfect matching iff $q(G \setminus S) \leq |S|$ for all $S \subseteq V(G)$.*

Cor 5.17 (Petersen 1891). *Every 3-regular gr. with no cut-edge has a perfect matching.*

Cor 5.19 (Berge 1958). *The largest matching in an n -vx G covers $n + \min_{S \subseteq V(G)} (|S| - q(G \setminus S))$ vxs.*

6 Planar Graphs

Def 6.1. A polygonal path or polygonal curve in the plane is the union of many line segments such that each segment starts at the end of the previous one and no point appears in more than one segment except for common endpoints of consecutive segments. In a polygonal u, v -path, the beginning of the first segment is u and the end of the last segment is v .

A drawing of a G is a function that maps each vx $v \in V(G)$ to a point $f(v)$ in the

plane and each edge uv to a polygonal $f(u), f(v)$ -path in the plane. The images of vxs are distinct. A point in $f(e) \cup f(e')$ other than a common end is a crossing. A gr. is planar if it has a drawing without crossings. Such a drawing is a planar embedding of G . A plane gr. is a particular drawing of a planar gr. in the plane with no crossings.

Def 6.4. An open set in the plane is a set $U \subset \mathbb{R}^2$ such that for every $p \in U$, all points within some small distance from p belong to U . A region is an open set U that contains a polygonal u, v -path for every pair $u, v \in U$ (that is, it is “path-conn.”). The faces of a plane gr. are the maximal regions of the plane that are disjoint from the drawing.

Thm 6.5 (Jordan curve theorem). *A simple closed polygonal curve C consisting of finitely many segments partitions the plane into exactly two faces, each having C as boundary.*

Rem 6.6. This is not true in three dimensions. In \mathbb{R} there

is a surface called the Möbius band which has only one side.

Rem 6.7. The faces of G are pairwise disjoint (they are separated by the edges of G). Two points are in the same face iff there is a polygonal path between them which does not cross an edge of G . Also, note that a finite gr. has a single unbounded face (the area “outside” of the graph).

Prop 6.8. *A plane forest has exactly one face.*

Def 6.9. The length of the face f in a planar embedding of G is the sum of the lengths of the walks in G that bound it.

Prop 6.11. *If $l(f_i)$ denotes the length of a face f_i in a plane G , then $2e(G) = \sum l(f_i)$.*

Thm 6.12 (Euler’s formula 1758). *If a conn. plane G has exactly n vxs, e edges and f faces, then $n - e + f = 2$.*

Thm 6.14. *If G is a planar gr. with at least three vxs, then $e(G) \leq 3|G| - 6$. If G is also triangle-free, then $e(G) \leq 2|G| - 4$.*

Cor 6.15. *If G is a planar bipartite n -vx gr. with $n \geq 3$ vxs then G has at most $2n - 4$ edges.*

Cor 6.16. *K_5 and $K_{3,3}$ are not planar.*

Rem 6.17 (Maximal planar graphs / triangulations). The proof of Theorem 6.14 shows that having $3n - 6$ edges in a simple n -vx planar gr. requires $2e = 3f$, meaning that every face is a triangle. If G has some face that is not a triangle, then we can add an edge between non-adjacent vxs on the boundary of this face to obtain a larger plane graph. Hence the simple plane graphs with $3n - 6$ edges, the triangulations, and the maximal plane graphs are all the same family.

6.1 Platonic Solids

Def 6.18. A polytope is a solid in 3 dimensions with flat faces, straight edges and sharp corners. Faces of a polytope are joined at the edges. A polytope is convex if the line connecting any two points of the polytope lies inside the polytope.

Cor 6.21. *If K is a convex polytope with v vxs, e edges and f faces then $v - e + f = 2$.*

7 Graph colouring

7.1 Vertex colouring

Def 7.1. A k -colouring of G is a labeling $f : V(G) \rightarrow \{1, \dots, k\}$. It is a proper k -colouring if $(x, y) \in E(G)$ implies $f(x) \neq f(y)$. A gr. G is k -colourable if it has a proper k -colouring. The chromatic number $\chi(G)$ is the minimum k such that G is k -colourable. If $\chi(G) = k$, then G is k -chromatic. If $\chi(G) = k$, but $\chi(H) < k$ for every proper subgr. H of G , then G is colour-critical or k -critical.

$\chi(K_n) = n$

Rem 7.3. The vxs having a given colour in a proper colouring must form an independent set, so $\chi(G)$ is the minimum # independent sets needed to cover $V(G)$. Hence G is k -colourable iff G is k -partite. Multiple edges do not affect chromatic number. Although we define k -colouring using numbers from $\{1, \dots, k\}$ as labels, the numerical values are usually unimportant, and we may use any set of size k as

labels.

7.3 Bounds on χ

Claim 7.6. If H is a subgr. of G then $\chi(H) \leq \chi(G)$.

Cor 7.7. $\chi(G) \geq \omega(G)$

Prop 7.9. $\chi(G) \geq \frac{|V(G)|}{\alpha(G)}$

Claim 7.10. For any gr. $G = (V, E)$ and any $U \subseteq V$ we have $\chi(G) \leq \chi(G[U]) + \chi(G[V \setminus U])$.

Claim 7.11. For any graphs G_1 and G_2 on the same vx set, $\chi(G_1 \cup G_2) \leq \chi(G_1)\chi(G_2)$.

Prop 7.12(i) $\chi(G)\chi(\overline{G}) \geq |G|$

(ii) $\chi(G) + \chi(\overline{G}) \leq |G| + 1$

7.4 Greedy colouring

Def 7.13. The greedy colouring with respect to a vx ordering v_1, \dots, v_n of $V(G)$ is obtained by colouring vxs in the order v_1, \dots, v_n , assigning to v_i the smallest-indexed colour not already used on its lower-indexed neighbours.

Def 7.15. Let $G = (V, E)$ be a graph. We say that G is k -degenerate if every subgr. of G has a vx of degree less than or equal to k .

Prop 7.16. G is k -degenerate iff there is an ordering v_1, \dots, v_n of the vxs of G such that each v_i has at most k neighbours among the vxs v_1, \dots, v_{i-1} .

Def 7.17. Define $dg(G)$ to be the minimum k such that G is k -degenerate.

Rem 7.18. $\delta(G) \leq dg(G) \leq \Delta(G)$.

Thm 7.19. $\chi(G) \leq 1 + dg(G)$

Cor 7.20. $\chi(G) \leq \Delta(G) + 1$.

Rem 7.21. This bound is tight if $G = K_n$ or if G is an odd cycle.

Thm 7.22 (Brooks 1941). If G is a conn. gr. other than a clique or an odd cycle, then $\chi(G) \leq \Delta(G)$.

7.5 Colouring planar graphs

Claim 7.23. A (simple) planar G contains a vx v of degree at most 5.

Cor 7.24. A planar G is 5-degenerate and thus 6-colourable.

Thm 7.25 (5 colour theorem, Heawood 1890). Every planar G is 5-colourable.

Thm 7.26 (Appel-Haken 1977, conjectured by Guthrie in 1852). Every planar gr. is 4-colourable. (the countries of every plane map can be 4-coloured so that neighbouring countries get distinct colours).

7.6 Art gallery thm

Thm 7.28. For any museum with n walls, $\lfloor n/3 \rfloor$ guards suffice.

8 Col. results

Thm 8.1 (Gallai, Roy). If D is an orientation of G with longest path length $l(D)$, then $\chi(G) \leq 1 + l(D)$. Furthermore, equality holds for some orientation of G .

8.1 Large girth and large χ

The bound $\chi(G) \geq \omega(G)$ can be tight, but (surprisingly) it can also be arbitrarily bad. There are graphs having arbitrarily large chromatic number, even though they do not contain K_3 . Many constructions of such graphs are known, though none are trivial. We give one here.

Thm 8.3. Mycielski's con-

struction produces a $(k + 1)$ -chromatic triangle-free gr. from a k -chromatic triangle-free graph.

Def 8.4. The girth of a gr. is the length of its shortest cycle.

Thm 8.5 (Erdos 1959). Given $k \geq 3$ and $g \geq 3$, there exists a gr. with girth at least g and chromatic number at least k .

Thm 8.8. There is a tournament on n vxs where any $\frac{\log_2(n)}{2}$ vxs are beaten by some other vx.

8.2 χ and clique minors

Def 8.9. Let $e = (x, y)$ be an edge of a gr. $G = (V, E)$. By G/e we denote the gr. obtained from G by contracting the edge e into a new vx v_e , which becomes adjacent to all the former neighbours of x and of y .

H is a minor of G if it can be obtained from G by deleting vxs and edges, and contracting edges.

Thm 8.12 (Mader). If the average degree of G is at least $2t - 2$ then G has a K_t minor.

Rem 8.13. It is known that $\overline{d}(G) \geq ct\sqrt{\log(t)}$ already implies the existence of a K_t minor in G , for some constant $c > 0$.

8.3 Edge-colourings

Def 8.14. A k -edge-colouring of G is a labelling $f : E(G) \rightarrow [k]$, the labels are "colours". A proper k -edge-colouring is a k -edge-colouring such that edges sharing a vx receive different colours, equivalently, each colour class is a matching. A G is k -edge-colourable if it has a proper k -edge-colouring. The edge-chromatic number or chromatic index $\chi'(G)$ is the minimum k such that G is k -edge colourable.

Rem 8.15. (i) An edge-colouring of a G is the same as a vx-colouring of its line gr. $L(G)$.

(ii) A G with maximum degree Δ has $\chi'(G) \geq \Delta$ since the edges incident to a vx of degree Δ must have different colours.

(iii) If G has maximum degree Δ then $L(G)$ has maximum degree at most $2(\Delta - 1)$. $\Rightarrow \chi'(G) \leq 2\Delta - 1$

Thm 8.16 (König 1916). *If G is a bipartite multigraph, then $\chi'(G) = \Delta(G)$.*

Thm 8.17 (Vizing). *Let G be a simple gr. with maximum degree Δ . Then $\Delta(G) \leq \chi_0(G) \leq \Delta(G) + 1$.*

8.4 List colouring

Def 8.19. For each vx v in a G , let $L(v)$ denote a list of colours available for v . A list colouring or choice function from a given collection of lists is a proper colouring f such that $f(v)$ is chosen from $L(v)$. A G is k -choosable or k -list-colourable if it has a proper list colouring from every assignment of k -element lists to the vxs. The list chromatic number or choosability $\chi_l(G)$ is the minimum k such that G is k -choosable.

Thm 8.20 (Erdos, Rubin, Taylor 1979). *If $m = \binom{2k-1}{k}$, then $K_{m,m}$ is not k -choosable.*

Def 8.21. Let $L(e)$ denote the list of colours available for e . A list edge-colouring is a proper edge-colouring f with $f(e)$ chosen from $L(e)$ for each e . The list chromatic

index or edge-choosability $\chi'l(G)$ is the minimum k such that G has a proper list edge-colouring for each assignment of lists of size k to the edges. Equivalently, $\chi'l(G) = \chi_l(L(G))$, where $L(G)$ is the line gr. of G .

Thm 8.23 (Galvin 1995). $\chi'l(K_{n,n}) = n$.

Def 8.24. A kernel of a digr. is an independent set S having an edge to every vx outside S . A digr. is kernel-perfect if every induced subdigr. has a kernel. Given a function $f : V(G) \rightarrow \mathbb{N}$, the G is f -choosable if a proper list colouring can be chosen whenever the lists satisfy $|L(x)| \geq f(x)$ for each x .

Lem 8.25. *If D is a kernel-perfect orientation of G and $f(x) = d_D^-(x)$ for all $x \in V(G)$, then G is $(1 + f)$ -choosable.*

9 The Matrix Tree Theorem

Thm 9.1 (Cayley's formula). *There are n^{n-2} labeled trees on n vxs.*

Cayley's formula).

Now consider an arbitrary conn. simple G on vx set $[n]$, and denote the # spanning trees by $t(G)$. The following celebrated result is Kirchhoff's matrix tree theorem. To formulate it, consider the incidence matrix B of G (as in Definition 1.13), and replace one of the two 1's by -1 in an arbitrary manner to obtain the matrix C (we say C is the incidence matrix of an orientation of G). $M = CC^T$ is then a symmetric $n \times n$ matrix, which is

$$\begin{pmatrix} d(1) & 0 & 0 \\ 0 & \ddots & 0 \\ 0 & 0 & d(n) \end{pmatrix} - A_G$$

Thm 9.2 (Matrix tree theorem). *We have $t(G) = \det M_{ii}$ for all $i = 1, \dots, n$, where M_{ii} results from M by deleting the i -th row and the i -th column.*

Thm 9.3 (Binet, Cauchy). *If P is an $r \times s$ matrix and Q is an $s \times r$ matrix with $r \leq s$, then*

$$\det(PQ) = \sum_Z (\det P_Z) (\det Q_Z)$$

where P_Z is the $r \times r$ submatrix of P with column set Z ,

and Q_Z is the $r \times r$ submatrix of Q with the corresponding rows Z , and the sum is over all r -sets $Z \subseteq [s]$.

10 More Thms on Hamiltonicity

Def 10.1. The (Hamiltonian) closure of a G , denoted $C(G)$, is the supergr. of G on $V(G)$ obtained by iteratively adding edges between pairs of nonadjacent vxs whose degree sum is at least n , until no such pair remains.

Thm 10.2 (Bondy Chvátal 1976). *A simple n -vx gr. is Hamiltonian iff its closure is Hamiltonian.*

Thm 10.3 (Chvátal 1972). *Suppose G has vx degrees $d_1 \leq \dots \leq d_n$. If $i < n/2$ implies that $d_i > i$ or $d_{n-i} \geq n-i$, then G is Hamiltonian.*

Thm 10.4 (Chvátal-Erdos 1972). *If $\kappa(G) \geq \alpha(G)$, then G has a Hamiltonian cycle (unless $G = K_2$).*

10.1 Pósa's Lemma

Let P be a path in a G , say from u to v . Given a vx $x \in P$, we write x^- for the vx preceding x on P , and x^+ for the vx following x on P

(whenever these exist). Similarly, for $X \subseteq V(P)$ we put $X^\pm := \{x^\pm : x \in X\}$

If $x \in P \setminus u$ is a neighbour of u in G , then $P \cup \{(u, x)\} \setminus \{(x, x^-)\}$ (which is a path in G with vx set $V(P)$) is said to have been obtained from P by a rotation fixing v . A path obtained from P by a (possibly empty) sequence of rotations fixing v is a path derived from P . The set of starting vxs of paths derived from P , including u , will be denoted by $S(P)$. As all paths derived from P have the same vx set as P , we have $S(P) \subseteq V(P)$.

Rem 10.5. If some sequence of rotations can delete the edge (x, x^-) , call this edge a broken edge. Note that every interval of the original path not containing broken edges is traversed by all derived paths as a whole piece (however, the direction can change).

Def 10.6. For a G and a subset $S \subseteq V(G)$, let $\partial S = \{v \in G \setminus S : \exists y \in S, v \sim y\}$.

Lem 10.7. *Let G be a graph, let $P = u \dots v$ be a longest*

path in G , and put $S := S(P)$. Then $\partial S \subseteq S^- \cup S^+$.

Lem 10.8. Let G be a graph, let $P = u \dots v$ be a longest path in G , and put $S := S(P)$. If $\deg(u) \geq 2$ then G has a cycle containing $S \cup \partial S$.

Cor 10.9. Fix $k \geq 2$ and let G be a gr. such that for all $S \subseteq V(G)$ with $|S| \leq k$, we have $|\partial S| \geq |2S|$. Then G has a cycle of length at least $3k$.

10.2 Tournaments

Def 10.10. A tournament is a directed gr. obtained by assigning a direction to every edge of the complete graph. That is, it is an orientation of K_n .

Thm 10.11. Every tournament has a Hamilton path.

Def 10.12. A tournament is strongly conn. if for all u, v there is a directed path from u to v .

Thm 10.13. A tournament T is strongly conn. iff it has a Hamilton cycle.

11 Kuratowski's Theorem

Def 11.1. A subdivision of a gr. H is a gr. obtained from

H by replacing the edges of H by internally vx disjoint paths of non-zero length with the same endpoints.

Rem 11.3. If G contains a subdivision of H , it also contains an H -minor.

Def 11.4. A Kuratowski gr. is a gr. which is a subdivision of K_5 or $K_{3,3}$. If G is a gr. and H is a subgr. of G which is a Kuratowski gr. then we say that H is a Kuratowski subgr. of G .

Thm 11.5 (Kuratowski 1930). A gr. is planar iff it has no Kuratowski subgraph.

Def 11.6. A straightline drawing of a planar G is a drawing in which every edge is a straight line.

Thm 11.7. If G is a gr. with no Kuratowski subgr. then G has a straightline drawing in the plane.

11.1 Convex drawings of 3-conn. graphs

Def 11.8. A convex drawing of G is a straightline drawing in which every non-outer face of G is a convex polygon, and the outer face is the

complement of a convex polygon. (That is, the boundary of each face is the boundary of a convex polygon).

Thm 11.9 (Tutte 1960). If G is a 3-conn. gr. which has no Kuratowski subgraphs then G has a convex drawing in the plane with no three vxs on a line.

Lem 11.10 (Thomassen 1980). Every 3-conn. G with at least five vxs has an edge e such that G/e is 3-conn.

Lem 11.11. If G has no Kuratowski subgraphs, then G/e has no Kuratowski subgraph, for any edge $e \in E(G)$.

11.2 Reducing the general case to the 3-conn. case

Def 11.12. Given a subdivision H' of H , we call the vxs of the original gr. branch vxs.

Fact 2. We make three observations.

1. In a Kuratowski subgraph, there are three internally vx-disjoint paths connecting any two branch vxs. For K_5 -subdivisions, we even have four such paths.

2. In a Kuratowski subgraph, there are four internally vx-disjoint paths between any two pairs of branch vxs.
3. Any cycle in a subdivision contains at least three branch vxs.

Prop 11.14. Let G be a gr. with at least 4 vxs which has no Kuratowski subgraph, and suppose that adding an edge-joining any pair of non-adjacent vxs creates a Kuratowski subgraph. Then G is 3-conn.

12 Ramsey Theory

Prop 12.1. Among six people it is possible to find three mutual acquaintances or three mutual non-acquaintances.

As we shall see, given a natural number s , there is an integer R such that if $n \geq R$ then every colouring of the edges of K_n with red and blue contains either a red K_s or a blue K_s . More generally, we define the Ramsey number $R(s, t)$ as the smallest value of N for which every red-blue colouring of K_N yields a red K_s or a blue K_t . In partic-

ular, $R(s, t) = 1$ if there is no such N such that in every red-blue colouring of K_N there is a red K_s or a blue K_t . It is obvious that $R(s, t) = R(t, s)$ for every $s, t \geq 2$ and $R(s, 2) = R(2, s) = s$.

Thm 12.2 (Erdős, Szekeres). The function $R(s, t)$ is finite for all $s, t \geq 2$. Quantitatively, if $s > 2$ and $t > 2$ then $R(s, t) \leq R(s - 1, t) + R(s, t - 1)$ and $R(s, t) \leq \binom{s+t-2}{s-1}$.

Thm 12.3. Given k and s_1, s_2, \dots, s_k , if N is sufficiently large, then every colouring of K_N with k colours is such that for some $i, 1 \leq i \leq k$, there is a K_{s_i} coloured with the i -th colour. The minimal value of N for which this holds is usually denoted by $R_k(s_1, \dots, s_k)$, and it satisfies $R_k(s_1, \dots, s_k) \leq R_k - 1(R(s_1, s_2), s_3, \dots, s_k)$.

Thm 12.4. Let $\min\{s, t\} > 3$. Then $R^{(3)}(s, t) \leq R(R^{(3)}(s - 1, t), R^{(3)}(s, t - 1)) + 1$

12.1 Applications

Thm 12.5 (Erdős-Szekeres 1935). Given an integer m , there exists a (least) integer

$N(m)$ such that every set of at least $N(m)$ points in the plane, with no three collinear, contains an m -subset forming a convex m -gon.

12.2 Bounds on Ramsey numbers

Thm 12.6 (Erdős 1947). For $p \geq 3$, we have $R(p, p) > 2^{p/2}$.

Thm 12.7. We have $R_k(3) \stackrel{\text{def}}{=} R_k(3, \dots, 3) \leq \lfloor e \cdot k! \rfloor + 1$.

12.3 Ramsey theory for integers

Thm 12.8 (Schur 1916). For every $k \geq 1$ there is an integer m such that every k -colouring of $[m]$ contains integers x, y, z of the same colour such that $x + y = z$.

12.4 Graph Ramsey numbers

Def 12.9. Let G_1, G_2 be graphs. $R(G_1, G_2)$ is the minimal N such that any red/blue colouring of K_N contains either a red copy of G_1 , or a blue copy of G_2 .

Rem 12.10. Note that $R(G_1, G_2) \leq R(|G_1|, |G_2|)$.

Thm 12.11 (Chvatal 1977). If T is any m -vx tree, then $R(T, K_n) = (m-1)(n-1) + 1$

13 Extremal problems

Def 13.2. $ex(n, H)$ is the maximal value of $e(G)$ among graphs G with n vxs containing no H as a subgraph.

13.1 Turán's theorem

Def 13.4. We call the gr. K_{n_1, \dots, n_r} with $|n_i - n_j| \leq 1$ the Turán graph, denoted by $T_{n,r}$.

Thm 13.5 (Turán 1941). Among all the n -vx simple graphs with no $(r+1)$ -clique, $T_{n,r}$ is the unique gr. having the maximum # edges.

Question 13.6. Let $a_1, \dots, a_n \in \mathbb{R}^d$ be vectors such that $|a_i| \geq 1$ for each $i \in [n]$. What is the maximum # pairs satisfying $|a_i + a_j| < 1$?

Claim 13.7. There are at most $\lfloor \frac{n^2}{4} \rfloor$ such pairs.

Def 13.8. For some fixed gr. H , we define $\pi(H) = \lim_{n \rightarrow \infty} ex(n, H) / \binom{n}{2}$

Thm 13.9 (Erdős-Stone). Let H be a gr. of chromatic number $\chi(H) = r + 1$. Then for every $\varepsilon > 0$ and large enough n , $(1 - \frac{1}{r}) \frac{n^2}{2} \leq ex(n, H) \leq (1 - \frac{1}{r}) \frac{n^2}{2} + n^2 \varepsilon$

13.2 Bipartite Turán Theorems

Thm 13.11. If a G on n vxs contains no 4-cycles, then $e(G) \leq \lfloor \frac{n}{4}(1 + \sqrt{4n-3}) \rfloor$.

Thm 13.13 (Kovári-Sós-Turán). For any integers $r \leq s$, there is a constant c such that every $K_{r,s}$ -free gr. on n vxs contains at most $cn^{1-\frac{1}{r}}$ edges. In other words, $ex(n, K_{r,s}) \leq cn^{1-\frac{1}{r}}$

Thm 13.14. There is c depending on k such that if G is a gr. on n vxs that contains no copy of C_{2k} , then G has at most $cn^{1+\frac{1}{k}}$ edges.

Question 13.15. Given n points in the plane, how many pairs can be at distance 1?

Thm 13.16 (Erdős). There are at most $cn^{3/2}$ pairs.

14 Exercises

14.1 Assignment 1

P 3. Prove that if a G is not conn. then its complement \bar{G} is conn. converse is not true.

P 4. Show that every gr. on at least two vxs contains two vxs of equal degree.

P 5. Prove that every gr. with $n \geq 7$ vxs and at

least $5n - 14$ edges contains a subgr. with minimum degree at least 6.

P 6. Show that in a conn. gr. any two paths of maximum length share at least one vx.

P 7. Prove that a gr. is bipartite iff it contains no cycle of odd length.

14.2 Assignment 2

P 1. Show that in a tree containing an even # edges, there is at least one vx with even degree.

P 2. Given a G and a vx $v \in V(G)$, $G - v$ denotes the subgr. of G induced by the vx set $V(G) \setminus \{v\}$. Show that every conn. G of order at least two contains vxs x and y such that both $G - x$ and $G - y$ are conn.

P 3. Let T be an n -vx tree with exactly $2k$ odd-degree vxs. Prove that T decomposes into k paths (i.e. its edge-set is the disjoint union of k paths).

P 4. Prove that a conn. G is a tree iff any family of pairwise (vx-)intersecting paths P_1, \dots, P_k in G have a common vx.

P 5(a) Prüfer codes corresponding to stars (i.e. to trees isomorphic to $K_{1,n-1}$) = 1 value.

(b) Prüfer codes containing exactly 2 different values = 2 connected stars

P 6. Let T be a forest on vx set $[n]$ with components T_1, \dots, T_r . Prove, by induction on r , that the # spanning trees on $[n]$ containing T is $n^{r-2} \prod_{i=1}^r |T_i|$. Deduce Cayley's formula.

14.3 Assignment 3

P 1. Prove that a conn. G is k -edge-conn. iff each block of G is k -edge-conn.

P 2. Let G be a gr. and suppose some two vxs $u, v \in V(G)$ are separated by $X \subseteq V(G) \setminus \{u, v\}$. Show that X is a minimal separating set (i.e. there is no proper subset $Y \subset X$ that separates u and v) iff every vx in X has a neighbor in the component of $G - X$ containing u and another in the component containing v .

P 3. Show that if G is a gr. with $|V(G)| = n \geq k + 1$ and $\delta(G) \geq (n + k - 2)/2$ then G is k -conn.

P 4. Prove that a G with at

least 3 vxs is 2-conn. iff for any three vxs x, y, z there is a path from x to z containing y .

P 5. Let G be a k -conn. graph, where $k \geq 2$. Show that if $|V(G)| \geq 2k$ then G contains a cycle of length at least $2k$.

14.4 Assignment 4

P 1. Show that if $k > 0$ then the edge set of any conn. gr. with $2k$ vxs of odd degree can be split into k trails.

P 2. Let G be a conn. gr. that has an Euler tour. T / F?

- (a) If G is bipartite then it has an even # edges. T
- (b) If G has an even # vxs then it has an even # edges. F
- (c) For edges e and f sharing a vx, G has an Euler tour in which e and f appear consecutively. F

P 3. Let G be a conn. gr. on n vxs with minimum degree δ . Show that

- (a) if $\delta \leq \frac{n-1}{2}$ then G contains a path of length 2δ , and
- (b) if $\delta \geq \frac{n-1}{2}$ then G contains a Hamiltonian path.

P 4. Show that the maximum # edges in a non-Hamiltonian gr. on $n \geq 3$ vxs is $\binom{n-2}{1} + 1$.

14.5 Assignment 5

P 1. Let G be a conn. gr. on more than 2 vxs such that every edge is contained in some perfect matching of G . Show that G is 2-edge-conn.

P 2(a) Let G be a gr. on $2n$ vxs that has exactly one perfect matching. Show that G has at most n^2 edges.

(b) Construct such a G containing exactly n^2 edges for any $n \in \mathbb{N}$.

P 3. Let A be a finite set with subsets A_1, \dots, A_n , and let d_1, \dots, d_n be positive integers. Show that there are disjoint subsets $D_k \subseteq A_k$ with $|D_k| = d_k$ for all $k \in [n]$ if and only if $|\bigcup_{i \in I} A_i| \geq \sum_{i \in I} d_i$.

P 4. Suppose M is a matching in a bipartite gr. $G = (A \cup B, E)$. We say that a path $P = a_1 b_1 \dots a_k b_k$ is an augmenting path in G if (b) $b_i a_{i+1} \in M$ for all $i \in [k-1]$ and a_1 and b_k are not covered by M . The name comes from

the fact that the size of M can be increased by flipping the edges along P (in other words, taking the symmetric difference of M and P): by deleting the edges $b_i a_{i+1}$ from M and adding the edges $a_i b_i$ instead.

- (a) Prove Hall's theorem by showing that if Hall's condition is satisfied and M does not cover A , then there is an augmenting path in G .
- (b) Show that if M is not a maximum matching (i.e. there is a larger matching in G) then the gr. contains an augmenting path. Is this true for non-bipartite graphs as well?

P 5. Show that for $k \geq 1$, every k -regular $(k-1)$ -edge-conn. gr. on an even # vxs contains a perfect matching.

14.6 Assignment 6

P 2(a) Show that every planar gr. has a vx of degree at most 5. Is there a planar gr. with minimum degree 5?

(b) Show that any planar bipartite gr. has vx of degree at most 3. Is there a planar bipartite gr. with

minimum degree 3?

P 3. Show that a conn. plane G is bipartite iff all its faces have even length.

P 4. Let G be a gr. on $n \geq 3$ vxs and $3n - 6 + k$ edges for some $k > 0$. Show that any drawing of G in the plane contains at least k crossing pairs of edges.

P 5. Let G be a plane gr. with triangular faces and suppose the vxs are colored arbitrarily with three colors. Prove that there is an even # faces that get all three colors.

P 6. Let S be a set of $n \geq 3$ points in the plane such that any two of them have distance at least 1. Show that there are at most $3n - 6$ pairs of distance exactly 1.

14.7 Assignment 7

P 1. T / F?

(a) If G and H are graphs on the same vx set, then $dg(G \cup H) \leq dg(G) + dg(H)$. F

(b) If G and H are graphs on the same vx set, then $\chi(G \cup H) \leq \chi(G) + \chi(H)$. F

(c) Every G has a $\chi(G)$ -coloring where $\alpha(G)$ vxs

get the same color. F

P 2. G has the property that any two odd cycles in it intersect (they share at least one vx in common). Prove that $\chi(G) \leq 5$.

P 3. For a vx v in a conn. G , let G_r be the subgr. of G induced by the vxs at distance r from v . Show that $\chi(G) \leq \max_{0 \leq r \leq n} \chi(G_r) + \chi(G_{r+1})$.

P 4. Let l be the length of the longest path in a G . Prove $\chi(G) \leq l + 1$ using the fact that if a gr. is not d -degenerate then it contains a subgr. of minimum degree at least $d + 1$.

P 5. Suppose the complement of G is bipartite. Show that $\chi(G) = \omega(G)$.

14.8 Assignment 8

P 1. For a given natural number n , let G_n be the following gr. with $\binom{n}{2}$ vxs and $\binom{n}{3}$ edges: the vxs are the pairs (x, y) of integers with $1 \leq x < y \leq n$, and for each triple (x, y, z) with $1 \leq x < y < z \leq n$, there is an edge joining vx (x, y) to vx (y, z) . Show that for any natural number k , the gr. G_n is triangle-free and has chro-

matic number $\chi(G_n) > k$ provided $n > 2k$.

P 2. Show that the theorem of Mader implies the following weakening of Hadwiger's conjecture: Any G with $\chi(G) \geq 2^{t-2} + 1$ has a K_t -minor.

P 3. Find the edge-chromatic $\# K_n$ (don't use Vizing's theorem).

P 4. Let G be a conn. k -regular bipartite gr. with $k \geq 2$. Show, using König's theorem, that G is 2-conn.

14.9 Assignment 9

P 1. Prove that every G of maximum degree Δ has an equitable $(\Delta + 1)$ -edge-coloring, i.e. one where each color class contains $\lfloor e = (\Delta + 1) \rfloor$ or $\lceil e = (\Delta + 1) \rceil$ edges, where e is the $\#$ edges in G .

P 2. The cartesian product $H \times G$ of graphs H and G is the gr. with vx set $V(H) \times V(G)$, with an edge between (v, u) and (v', u') if $v = v'$ and u is adjacent to u' in G , or if $u = u'$ and v is

adjacent to v' in H . Prove that if $\chi'(H) = \Delta(H)$ then $\chi'(H \times G) = \Delta(H \times G)$

P 3. Show that $\chi(C_n) = \chi_l(C_n)$ for any $n \geq 3$.

P 4. Let G be a bipartite gr. on n vxs. Prove that $\chi_l(G) \leq 1 + \log_2(n)$ using the probabilistic method.

P 5. Let G be a complete r -partite gr. with all parts of size 2. (In other words, G is K_{2r} minus a perfect matching.) Show, using a combination of induction and Hall's theorem, that $\chi_l(G) = r$.

14.10 Assignment 10

P 1. How many spanning trees does $K_{r,s}$ have? $r^{s-1}s^{r-1}$

P 2. Find the $\#$ spanning trees of $K_n - e$ (the complete gr. on n vxs with one edge removed): $(n-2)n^{n-3}$

P 3. In this exercise we prove the following alternative form of the matrix-tree theorem. For an n -vx conn. G , the $\#$ spanning trees in G is equal to the product of the nonzero eigenvalues of

the Laplacian matrix M of G , divided by n . (This matrix M is as in the lecture notes).

P 4(a) Prove that any n -by- n bipartite gr. with minimum degree $\delta > n/2$ contains a Hamilton cycle.

(b) Show that this is not necessarily the case if $\delta \leq n/2$.

14.11 Assignment 11

P 1. T / F?

(a) If every vx of a tournament has positive in- and out-degree, then the tournament contains a directed Hamilton cycle. F

(b) If a tournament has a directed cycle, then it has a directed triangle. T

P 2. Let G be a gr. on $n \geq 3$ vxs with at least $\alpha(G)$ vxs of degree $n-1$. Show that G is Hamiltonian.

P 3. Suppose G is a gr. on n vxs where all the degrees are at least $\frac{n+q}{2}$. Show that any set F of q independent edges is contained in a Hamiltonian cycle.

14.12 Assignment 12

P 1. The lower bound for $R(p, p)$ that you learn in

the lectures is not a constructive proof: it merely shows the existence of a red-blue coloring not containing any monochromatic copy of K_p by bounding the $\#$ bad (b) graphs. Give an explicit coloring on $K_{(p-1)^2}$ that proves $R(p, p) > (p-1)^2$.

P 2. Prove that for every fixed positive integer r , there is an n such that any coloring of all the subsets of $[n]$ using r colors contains two non-empty disjoint sets X and Y such that X, Y and $X \cup Y$ have the same color.

P 3. Prove that for every $k \geq 2$ there exists an integer N such that every coloring of $[N]$ with k colors contains three distinct numbers a, b, c satisfying $ab = c$ that have the same color.

P 4. For every $k \geq 2$ there is an N such that any k -coloring of $[N]$ contains three distinct integers a, b, c of the same color satisfying $a + b = c$.

P 5(a) Let $n \geq 1$ be an integer. Show that any se-

quence of $N \geq R(n, n)$ distinct numbers, a_1, \dots, a_N contains a monotone (increasing or decreasing) subsequence of length n .

Let $k, l \geq 1$ be integers and show that any sequence of $kl + 1$ distinct numbers a_1, \dots, a_{kl+1} contains a monotone increasing subsequence of length $k + 1$ or a monotone decreasing subsequence of length $l + 1$.

14.13 Assignment 13

P 1. Let H be an arbitrary fixed gr. and prove that the sequence $ex(n, H) / \binom{n}{2}$ is (not necessarily strictly) monotone decreasing in n .

P 2. Among all the n -vx K_{r+1} -free graphs, the Turan gr. $T_{n,r}$ contains the maximum $\#$ triangles (for any $r, n \geq 1$).

P 3. Let X be a set of n points in the plane with no two points of distance greater than 1. Show that there are at most $\frac{n^2}{3}$ pairs of points in X that have distance greater than $\frac{1}{\sqrt{2}}$.