

THE RAGE WAS NEVER ABOUT THE SOUND. IT NEVER IS.

MISOPHONIA

◇ PROTOCOL · REGULATION IN THE ADULT NERVOUS SYSTEM

A sound enters the room -- a chew, a sniff, a pen click -- and something in you goes to war. Not annoyance. Not preference. A full nervous system response: heat, rage, the urge to leave your own skin.

This is not a character flaw and it is not something you're imagining. It's a documented pattern in the brain's salience network, and adults who live with it have spent years mistaking a wiring difference for a personal failure.

This protocol does not promise the sounds will stop bothering you. It gives you a way to make the reaction smaller and your choices bigger.

-
- EVIDENCE VS. STORY
 - BODY SCAN
 - MIRROR CHECK + FUTURE ECHO
 - COST LEDGER
 - LIE DETECTOR
 - IN-THE-MOMENT GROUNDING

THE REASON BEHIND THIS PROTOCOL

THIS IS WIRING, NOT WEAKNESS.

Misophonia was given its first expert consensus definition in 2022: a disorder of decreased tolerance to specific everyday sounds, most often oral or nasal -- chewing, breathing, sniffing, throat-clearing -- producing a reaction far past irritation. Brain research on hundreds of adults has since confirmed a disorder-specific signature in the anterior insula and salience network, the same circuitry that flags danger, not simple sound.

It has no standalone entry in the DSM-5 or ICD-11 yet. That reflects how young the research is, not whether the experience is real. National data on adults with misophonia shows the scale of what it actually costs: nearly half report trouble with social activities, roughly a third report work impairment, and a similar share report strain in close relationships.

As of early 2026, the first randomized controlled trials in adults confirmed it: both cognitive behavioral therapy and acceptance and commitment therapy measurably reduce symptoms. The goal was never total silence. It's a smaller reaction, and a bigger life around it.

This protocol is not a substitute for that treatment -- it's a structured way to work between sessions, or before you have one. If reactions are frequent, intense, or shrinking your life through avoidance, a clinician trained in evidence-based, transdiagnostic therapy (CBT/ACT) is the fastest path forward. An audiologist evaluating sound tolerance can also help rule out related conditions like hyperacusis.

01

EVIDENCE VS. STORY · THE TRIGGER LOG

WHAT SOUND, AND WHAT STORY ARE YOU ADDING TO IT?

Separate the acoustic fact from the meaning your mind attaches to it.

What is the exact sound, and who or what makes it?

The sound is...

What do you tell yourself the sound means about the person making it, or about you?

The story I add is...

02

BODY SCAN · THE FIRST SECOND

WHERE DOES THE REACTION HIT FIRST?

Misophonia is a body-first hijack before it's a thought. Track exactly where and how fast.

Where in your body does the reaction land first, and how quickly after the sound starts?

It hits my...

What is the urge your body produces -- flee, confront, freeze?

My body wants to...

03

MIRROR CHECK + FUTURE ECHO · THE HISTORY OF THE TRIGGER

**WHEN DID THIS SOUND FIRST
BECOME DANGEROUS?**

Many trigger sounds trace back to a specific relationship or period. Find the thread.

When do you first remember this sound, or one like it, causing this reaction?

The first time I remember this was...

In the moment it happens now, who does your body react as if it's actually with?

Some part of me still reacts like it's...

04 COST LEDGER · WHAT AVOIDANCE HAS BOUGHT

WHAT HAS THIS COST YOU -- AND WHAT HAS AVOIDING IT COST YOU?

Misophonia often shrinks a life through avoidance. Take the full account, not just the sound's price.

What has this trigger cost you in relationships, meals, or shared spaces?

It has cost me...

What has avoiding the trigger cost you instead -- the plans skipped, the rooms left?

Avoiding it has cost me...

05 LIE DETECTOR · THE COVER STORY

WHAT DO YOU SAY INSTEAD OF THE TRUTH?

Most adults with misophonia develop a cover story. Name it, then draft the honest version.

What do you usually say when you leave a room or snap at someone, instead of naming the trigger?

What I usually say is...

What would the honest, plain version of that sentence be?

The honest version is...

REGULATE DURING THE TRIGGER, NOT JUST AFTER IT.

GROUND YOURSELF WHILE IT'S HAPPENING, AND BEFORE IT HAPPENS AGAIN.

WHEN IT'S HAPPENING RIGHT NOW

IN THE MOMENT

Use this the second you feel the reaction start, before you speak or leave.

- 01 Name the sound plainly, in your head: "That is a chewing sound. It is not a threat."
- 02 Exhale longer than you inhale -- four counts in, seven counts out, three rounds.
- 03 Give your attention a job: count objects of one color in the room, or press your feet into the floor and name the sensation.

PLAN THE EXIT BEFORE YOU NEED ONE

BEFORE THE TRIGGER

Adults who manage this well plan in advance. They don't invent the plan mid-reaction.

- 01 Choose one low-drama exit line for shared spaces: "I need to step out for a minute."
- 02 Keep one physical tool on hand -- earbuds, a white-noise app, a seat near an exit.
- 03 Tell one trusted person your plan in advance, so leaving doesn't require an explanation in the moment.

THE WORK ISN'T DONE UNTIL IT LIVES IN YOUR ACTUAL LIFE.

06

INTEGRATION · CARRY IT FORWARD

WHAT DOES ONE SMALLER REACTION LOOK LIKE THIS WEEK?

The goal was never silence. It's a smaller reaction, and a bigger life around it.

Name one specific situation this week where you'll use the in-the-moment tools instead of the old response.

This week, I will use the tools when...

What does it look like to tell one person the truth about this, instead of the cover story?

I will tell the truth to...