

Please write clearly in block capitals.

Centre number

|  |  |  |  |  |
|--|--|--|--|--|
|  |  |  |  |  |
|--|--|--|--|--|

Candidate number

|  |  |  |  |
|--|--|--|--|
|  |  |  |  |
|--|--|--|--|

Surname

---

Forename(s)

---

Candidate signature

---

## Level 2 Certificate FURTHER MATHEMATICS

Paper 2

Calculator

Thursday 21 June 2018


Afternoon

Time allowed: 2 hours

### Materials

For this paper you must have:

- a calculator
- mathematical instruments.



### Instructions

- Use black ink or black ball-point pen. Draw diagrams in pencil.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- Do all rough work in this book. Cross through any work you do not want to be marked.
- In all calculations, show clearly how you work out your answer.

### Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 105.
- You may ask for more answer paper, graph paper and tracing paper. These must be tagged securely to this answer book.
- The use of a calculator is expected but calculators with a facility for symbolic algebra must **not** be used.

### For Examiner's Use

| Pages        | Mark |
|--------------|------|
| 3            |      |
| 4–5          |      |
| 6–7          |      |
| 8–9          |      |
| 10–11        |      |
| 12–13        |      |
| 14–15        |      |
| 16–17        |      |
| 18–19        |      |
| 20–21        |      |
| 22–23        |      |
| 24–25        |      |
| 26–27        |      |
| 28–29        |      |
| <b>TOTAL</b> |      |



J U N 1 8 8 3 6 0 2 0 1


IB/M/Jun18/E5

**8360/2**

**Formulae Sheet**


**Volume of sphere**  $= \frac{4}{3} \pi r^3$

**Surface area of sphere**  $= 4\pi r^2$




**Volume of cone**  $= \frac{1}{3} \pi r^2 h$

**Curved surface area of cone**  $= \pi r l$



**In any triangle ABC**

**Area of triangle**  $= \frac{1}{2} ab \sin C$



**Sine rule**  $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$

**Cosine rule**  $a^2 = b^2 + c^2 - 2bc \cos A$

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

**The Quadratic Equation**

The solutions of  $ax^2 + bx + c = 0$ , where  $a \neq 0$ , are given by  $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

**Trigonometric Identities**

$\tan \theta \equiv \frac{\sin \theta}{\cos \theta} \quad \sin^2 \theta + \cos^2 \theta \equiv 1$



Answer **all** questions in the spaces provided.

1 The  $n$ th term of a sequence is  $\frac{1420 - 5n}{1420 + 5n}$

1 (a) Work out the **position** of the term that has the value zero.

[2 marks]

$$\frac{1420 - 5n}{1420 + 5n} = 0$$

$$1420 - 5n = 0$$

$$1420 = 5n$$

$$n = 284$$

Answer \_\_\_\_\_

1 (b) Write down the limiting value of the sequence as  $n \rightarrow \infty$

[1 mark]

$$\lim_{n \rightarrow \infty} = \frac{-5n}{+5n} = \frac{-5}{+5} = -1$$

Answer  $x \rightarrow -1$  \_\_\_\_\_

Turn over for the next question



2  $P(-3, -10)$  and  $Q(a, b)$  are points on a straight line with gradient 12

Work out one possible pair of integer values for  $a$  and  $b$ .

[2 marks]

$$-3 + 1 = -2$$

$$-10 + 12 = 2$$

$$a = \underline{-2}$$

$$b = \underline{2}$$



3  $p = \frac{m+2}{m^2+1}$

3 (a) Work out the value of  $p$  when  $m = -5.5$

[1 mark]

$$= \frac{-5.5+2}{(-5.5)^2+1}$$

$$= \frac{-3.5}{31.25}$$

Answer

$$-\frac{14}{125}$$

3 (b) Work out the values of  $m$  when  $p = 2$

[3 marks]

$$2 = \frac{m+2}{m^2+1}$$

$$2m^2 + 2 - m - 2 = 0$$

$$2m^2 - m = 0$$

$$m(2m-1) = 0$$

$$m=0, 2m-1=0$$

$$2m=1$$

$$m=\frac{1}{2}$$


Answer

$$0, \frac{1}{2}$$

Turn over for the next question



- 4  $A, B, C$  and  $D$  are points on a circle, centre  $O$ .



$$\begin{aligned} x + (180 - y) &= 180 \\ x - y &= 0 \\ x &= y \end{aligned}$$

Which statement is correct?

Tick **one** box.

[1 mark]

$x + y = 180^\circ$  and  $w = 2x$

☐

$x + y = 180^\circ$  and  $x = 2w$

☐

$x = y$  and  $w = 2x$

☒

$x = y$  and  $x = 2w$

☐



5 On the grid, draw the graph of  $y = f(x)$

$$f(x) = x + 4 \quad -4 \leq x < 0$$

$$= 4 - 3x \quad 0 \leq x < 2$$

$$= -2 \quad 2 \leq x \leq 5$$

[4 marks]



Turn over for the next question



6  $f(x) = x^2 - 7$  for all values of  $x$   
 $g(x) = 1 - 3x$  for  $-4 \leq x \leq 4$

- 6 (a) Work out the range of  $f(x)$ .  
 Give your answer as an inequality.

[1 mark]

Answer  $f(x) \geq -7$

- 6 (b) Work out the range of  $g(x)$ .  
 Give your answer as an inequality.

[2 marks]

$x =$        $1 - 3x$   
 $-4$        $13$   
 $0$        $1$   
 $4$        $-11$

Answer  $-11 \leq g(x) \leq 13$





6 (c) Solve  $2f(x) = g(x)$

You **must** show your working.

Give your answers to 3 decimal places.

[4 marks]

$$2x^2 - 14 = 1 - 3x$$

$$2x^2 + 3x - 15 = 0$$

$$2(x^2 + 1.5x) = 15$$

$$2\left(x + 0.75\right)^2 - \frac{9}{16} = 15$$

$$2(x + 0.75)^2 - \frac{9}{8} = 15$$

$$2(x + 0.75)^2 = 16\frac{1}{8}$$

$$(x + 0.75)^2 = 8\frac{1}{16} = \frac{129}{16}$$

$$x + 0.75 = \pm \frac{\sqrt{129}}{4}$$


$$x = \frac{-1.5 \pm \sqrt{129}}{4}$$

Answer 2.089, -3.589

Turn over for the next question



7

 $PQRS$  is a trapezium.

The area of the trapezium is 63 square units.

Work out the value of  $a$ .

[2 marks]

$$A = \frac{a+b}{2} h$$

$$63 = \frac{(9-5) + (10-2)}{2} a$$

$$63 = \frac{4+8}{2} a$$


$$63 = 6a$$

$$a = 10\frac{1}{2}$$

Answer \_\_\_\_\_



8

Here is a sketch of triangle  $ABC$ . $P$  is a point on  $AB$ . $AP : PB$  is  $3 : 1$ Work out the length  $PC$ .

Give your answer to 4 significant figures.

[4 marks]

$$\vec{AB} = (6, -5) \quad \frac{\vec{AB}}{4} = \left(\frac{3}{2}, -\frac{5}{4}\right)$$

$$P = B - \frac{\vec{AB}}{4} = \left(7 - \frac{3}{2}, 1 + \frac{5}{4}\right) = \left(5\frac{1}{2}, 2\frac{1}{4}\right)$$

$$PC = \sqrt{\left(12 - 2\frac{1}{4}\right)^2 + \left(9 - 5\frac{1}{2}\right)^2}$$

$$= \sqrt{\left(9\frac{3}{4}\right)^2 + \left(3\frac{1}{2}\right)^2}$$

$$= 10.36$$

Answer \_\_\_\_\_ units



9

$$y = \frac{2x^7 + 15x^2}{3x} = \frac{2x^6 + 15x}{3} = \frac{2}{3}x^6 + 5x$$

Work out the value of  $x$  when  $\frac{dy}{dx} = 133$

[4 marks]

$$\frac{dy}{dx} = 4x^5 + 5$$

$$4x^5 + 5 = 133$$

$$4x^5 = 128$$

$$x^5 = 32$$

$$x = 2$$

Answer 2



10

The transformation matrix  $\begin{pmatrix} a & b \\ 2a & 3b \end{pmatrix}$  maps the point  $(1, -3)$  onto the point  $(1, 4)$

Work out the values of  $a$  and  $b$ .

You **must** show your working.

[5 marks]

$$\begin{pmatrix} a & b \\ 2a & 3b \end{pmatrix} \begin{pmatrix} 1 \\ -3 \end{pmatrix} = \begin{pmatrix} 1 \\ 4 \end{pmatrix}$$

$$\begin{pmatrix} a-3b \\ 2a-9b \end{pmatrix} = \begin{pmatrix} 1 \\ 4 \end{pmatrix}$$

$$\begin{array}{r} \textcircled{A} \quad a-3b=1 \\ \textcircled{B} \quad 2a-9b=4 \\ - \quad \textcircled{2A} \quad 2a-6b=2 \\ \hline -3b=2 \\ b=-\frac{2}{3} \end{array}$$

$$\begin{array}{l} a-3b=1 \\ a+\frac{6}{3}=1 \\ a+2=1 \\ a=-1 \end{array}$$

$$a = -1$$

$$b = -\frac{2}{3}$$

Turn over for the next question

Turn over ►



11

Expand and simplify fully  $(x+2)(x+3)(x+4)$  [3 marks]

$$= (x^2 + 5x + 6)(x + 4)$$

$$= x^3 + 4x^2 + 5x^2 + 20x + 6x + 24$$

$$= x^3 + 9x^2 + 26x + 24$$

Answer  $x^3 + 9x^2 + 26x + 24$



- 12 (a) Write  $\frac{7}{9x} + \frac{2}{3x^2}$  as a single fraction in its simplest form.

[3 marks]

$$= \frac{7(3x^2) + 2(9x)}{(9x)(3x^2)} = \frac{2(x^2 + 18x)}{27x^3} = \frac{3(7x+6)}{3(9x^2)} = \frac{7x+6}{9x^2}$$

Answer \_\_\_\_\_

- 12 (b) Show that  $\frac{x^4}{x+4} \times \frac{x+2}{x} \div \frac{x^2}{3x+12}$

simplifies to the form  $ax^2 + bx$  where  $a$  and  $b$  are integers.

[4 marks]


$$\begin{aligned}
 &= \frac{x^3(x+2)}{x+4} \times \frac{3x+12}{x} \div \frac{x^2}{3x+12} \\
 &= \frac{x^3(x+2)(3x+12)}{x+4} = \frac{(x^2+2x)(3x+12)}{x+4} \\
 &= \frac{3x^3 + 18x^2 + 24x}{x+4} = \frac{3x(x^2 + 6x + 8)}{x+4} = \frac{3x(x+4)(x+2)}{x+4} \\
 &= 3x(x+2) = 3x^2 + 6x
 \end{aligned}$$

Turn over for the next question

Turn over ►



13 (a) Here is a sketch of  $y = \tan x$  for  $0^\circ \leq x \leq 360^\circ$



How many solutions of  $\tan x = k$  where  $k > 0$  are between  $90^\circ$  and  $360^\circ$ ?

*tan x > 0*

[1 mark]

Answer

*180 ?*

*1*






13 (b)  $0 < p < 1$

How many solutions of  $\sin x = p - 1$  are between  $0^\circ$  and  $180^\circ$ ?

You may use a sketch graph to help you.

[1 mark]

$$-1 < \sin x < 0$$



Answer


0

13 (c) State the coordinates of each point where the graph

$$y = \cos x \quad \text{for } 0^\circ \leq x \leq 360^\circ$$

meets or intersects an axis.

[2 marks]



Answer

$(0, 1), (90, 0), (270, 0)$



- 14 (a) Factorise fully  $12pq^3r - 18pq^2r^2 + 24pq^2r$

[2 marks]

$$6pq^2r(2q - 3r + 4)$$

Answer \_\_\_\_\_

- 14 (b) Factorise fully  $6(y+3)^5 + 4(y+3)^4$

Give your answer in its simplest form.

Do **not** attempt to expand  $(y+3)^5$  or  $(y+3)^4$ 

[3 marks]

$$\begin{aligned} &(y+3)^4(6(y+3) + 4) \\ &(y+3)^4(6y+18+4) \\ &(y+3)^4(6y+22) \\ &2(y+3)^4(3y+11) \end{aligned}$$

Answer \_\_\_\_\_

- 14 (c) Factorise fully  $48 - 75x^2$

[2 marks]

$$\begin{aligned} &= -3(25x^2 - 16) \\ &= -3(5x+4)(5x-4) \end{aligned}$$

Answer \_\_\_\_\_



15

Work out the rate of change of  $y$  with respect to  $x$  at the point on the curve

$$y = x^2(x^2 - 9) \quad \text{where} \quad x = -2$$

You **must** show your working.

[4 marks]

$$y = x^4 - 9x^2$$

$$\frac{dy}{dx} = 4x^3 - 18x, \quad x = -2, \quad = 4 \times -8 - 18 \times -2$$

$$= -32 + 36$$

$$= 4$$

Answer 4

Turn over for the next question



16

$$A = 2 - 5x \quad B = 3x - 1 \quad C = x^2$$

Show that  $(2A + 3B)^2 \equiv A + B + C$

[4 marks]

$$(2A + 3B)^2 \equiv (2(2 - 5x) + 3(3x - 1))^2$$

$$\equiv (4 - 10x + 9x - 3)^2$$

$$\equiv (1 - x)^2$$

$$\equiv x^2 - 2x + 1$$

$$A + B + C \equiv x^2 + 3x - 1 + 2 - 5x$$

$$\equiv x^2 - 2x + 1$$

SAME

17

A circle has equation  $x^2 + y^2 = 29$

$P$  is the point  $(-5, 2)$

17 (a)

Show that  $P$  is on the circle.

[1 mark]

$$= 2^2 + (-5)^2$$


$$= 4 + 25 = 29$$



- 17 (b) The tangent to the circle at  $P$  intersects the  $x$ -axis at point  $Q$ .

Work out the  $x$ -coordinate of  $Q$ .

You **must** show your working.



[4 marks]

$$\frac{dy}{dx} = \frac{-2}{5}, \text{ gradient} = \frac{5}{2}$$

$$y = mx + c$$

$$y = \frac{5x}{2} + c$$

$$2 = \frac{5 \times -5}{2} + c$$

$$2 = -\frac{25}{2} + c$$

$$c = \frac{29}{2} = 14\frac{1}{2}$$

$$y = \frac{5}{2}x + \frac{29}{2}$$

$$2y = 5x + 29$$

$$0 = 5x + 29$$

$$-29 = 5x$$

$$x = -\frac{29}{5}$$

Answer \_\_\_\_\_



- 18 (a) Work out all the **integer** values of  $x$  for which

$$-5 < 4x + 3 \leq 13$$

$$-8 < 4x \leq 10$$

$$-2 < x \leq 2.5$$

[3 marks]

Answer -1, 0, 1, 2

- 18 (b) Work out the range of values of  $x$  for which

$$x^2 - 11x + 28 > 0$$

You **must** show your working.

$$(x-7)(x-4) > 0$$

$$x < 4 \text{ and } x > 7$$

Answer \_\_\_\_\_

[3 marks]



19

Use **matrix multiplication** to show that, in the  $x$ - $y$  plane,

- a reflection in the line  $y = -x$ , followed by
- a rotation,  $90^\circ$  anticlockwise about the origin, followed by
- a reflection in the  $x$ -axis

is equivalent to a transformation by the identity matrix.

**[5 marks]**

$$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$


??  
, ,

Turn over for the next question

Turn over ►



20

 $PQRSTU$  is a triangular prism. $PQRS$  is a rectangle and angle  $QRU = 90^\circ$  $PQ = 10 \text{ cm}$        $QR = 12 \text{ cm}$        $UR = 7 \text{ cm}$  $M$  is the midpoint of  $PQ$ .

20 (a)

Calculate the size of the angle between the line  $UM$  and the plane  $PQRS$ .

[4 marks]

$$a^2 + b^2 = c^2$$

$$UQ = \sqrt{7^2 + 12^2} = \sqrt{193}$$

$$UM = \sqrt{193 + 5^2} = \sqrt{218}$$

$$RM = \sqrt{12^2 + 5^2} = 13$$

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

$$UMR = \cos^{-1} \left( \frac{218 + 13^2 - 7^2}{2 \times \sqrt{218} \times 13} \right)$$

$$= \cos^{-1} (0.88 \dots)$$

$$\approx 28.3^\circ$$

Answer 28.3° degrees



20 (b)

Calculate the size of the angle between the planes  $UMR$  and  $UQR$ . AKA

[2 marks]

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

$$MQR = \cos^{-1} \left( \frac{13^2 + 12^2 - 5^2}{2 \times 13 \times 12} \right)$$

$$= \cos^{-1} \left( \frac{12}{13} \right)$$

$$= 22.6$$

Answer \_\_\_\_\_ degrees

Turn over for the next question



21 The continuous curve  $y = f(x)$  has exactly two stationary points.


Here is some information about the curve.

|                 |                 |                 |                 |                 |
|-----------------|-----------------|-----------------|-----------------|-----------------|
| $x < -1$        | $x = -1$        | $-1 < x < 2$    | $x = 2$         | $x > 2$         |
| $\frac{dy}{dx}$ | $\frac{dy}{dx}$ | $\frac{dy}{dx}$ | $\frac{dy}{dx}$ | $\frac{dy}{dx}$ |
| is positive     | is zero         | is negative     | is zero         | is negative     |

$$f(-1) = 3 \quad \text{and} \quad f(2) = 1$$

State the coordinates **and** the nature of each of the stationary points.

[3 marks]



stationary point ( -1 , 3 ) nature Max

stationary point ( 2 , 1 ) nature ? inflection



- 22 (a)  $8 \cos x + 5 \sin x = 0$  where  $90^\circ < x < 180^\circ$

Work out the size of angle  $x$ .

[3 marks]

$$\begin{aligned} & \div \cos x \quad ? \\ & 8 + 5 \frac{\sin x}{\cos x} = 0 \\ & 5 \tan x = -8 \\ & \tan x = -1.6 \\ & \theta = -57.99, +180 = 122 \end{aligned}$$

Answer 122 degrees

- 22 (b)  $6 \sin^2 x + 4 \cos^2 x \equiv A + B \cos^2 x$  where A and B are integers.

Work out the values of A and B.

You **must** show your working.

[2 marks]

$$\begin{aligned} & 6 \sin^2 x + 4 \cos^2 x \equiv ( \\ & \equiv 6 \sin^2 x + 4 \cos^2 x \\ & \equiv 6(1 - \cos^2 x) + 4 \cos^2 x \\ & \equiv 6 - 6 \cos^2 x + 4 \cos^2 x \end{aligned}$$



A = 6 B = -2



23

For each of these two function machines, when the input is  $a$  the output is  $b$ .

$k > 0$  and  $k \neq 1$  and  $a > 0$



Work out an expression for  $a$  in terms of  $k$ .

Give your answer in its simplest form.

5  
[6 marks]

$$k(a^2 + 3) = b, \quad ka^2 + 3k = b$$

$$(ak)^2 + 3 = b, \quad a^2k^2 + 3 = b$$

$$ka^2 + 3k = a^2k^2 + 3$$

$$a^2k - a^2k^2 = 3 - 3k$$

$$a^2(k - k^2) = 3 - 3k$$

$$a^2 = \frac{3 - 3k}{k - k^2}$$

$$a = \sqrt{\frac{3 - 3k}{k - k^2}}$$

*cancel out (1-k)*

$$a = \sqrt{\frac{3}{k}}$$

Answer \_\_\_\_\_



24

Work out the value of  $p$  when

$$9^{0.5p} \times 81 = 27^{2p-1}$$

$$3^{2 \cdot 0.5p} \times 3^4 = 3^{2p-1}$$

$$3^{p+4} = 3^{2p-1}$$

$$p+4 = 2p-1$$

$$4+1 = 2p-p$$

$$p = 5$$

[4 marks]

Answer

END OF QUESTIONS



**There are no questions printed on this page**

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**



**There are no questions printed on this page**

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**



**There are no questions printed on this page**

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**

**Copyright information**

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third party copyright material will be published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from [www.aqa.org.uk](http://www.aqa.org.uk) after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2018 AQA and its licensors. All rights reserved.

