

# Pure Maths Consolidation Notes

Jack Maguire

# Contents

<b>I Book 1</b>	<b>3</b>
<b>3 Inequalities</b>	<b>4</b>
3.1 Basics . . . . .	4
3.2 Set Notation . . . . .	4
3.3 Quadratics . . . . .	4
<b>5 Coordinate Geometry</b>	<b>6</b>
5.1 Characteristics of Lines . . . . .	6
5.1.1 Gradient . . . . .	6
5.1.2 Y-Intercept . . . . .	7
5.1.3 Lines between points . . . . .	7
5.2 Solving Line Equations . . . . .	8
<b>6 Circles</b>	<b>9</b>
6.1 Intersections between lines and circles . . . . .	9
<b>8 Binomial Expansion</b>	<b>10</b>
<b>9 Trig Ratios</b>	<b>11</b>
<b>10 Trig Functions</b>	<b>12</b>
<b>12 Differentiation</b>	<b>13</b>
<b>13 Integration</b>	<b>14</b>
<b>14 Logs &amp; Exponents</b>	<b>15</b>
14.1 Logarithms . . . . .	15
14.1.1 Log Laws . . . . .	15
14.2 Exponentials . . . . .	17
14.2.1 Euler's Number - $e$ . . . . .	17
14.3 Modelling With Exponentials . . . . .	17
<b>II Other Bits</b>	<b>19</b>
<b>1 Transforming &amp; Sketching Graphs</b>	<b>20</b>
1.1 Graph Appearances . . . . .	20
1.1.1 Transformations . . . . .	20
1.1.2 Combining Transformations . . . . .	21
1.1.3 Graph Shapes . . . . .	22
1.2 Solving Using Graphs . . . . .	23
<b>2 Polynomial Division, Factor Theorem &amp; Cubics</b>	<b>25</b>
2.1 Polynomial Division . . . . .	25
2.2 Factor Theorem . . . . .	26

<b>III Book 2</b>	<b>28</b>
<b>1 Algebraic Methods</b>	<b>29</b>
<b>2 Functions &amp; Graphs</b>	<b>30</b>
<b>6 Trig Functions</b>	<b>31</b>
<b>7 Trig Modelling</b>	<b>32</b>
<b>8 Parametric Equations</b>	<b>33</b>
<b>9 Differentiation</b>	<b>34</b>
<b>11 Integration</b>	<b>35</b>

Part I

Book 1

## Chapter 3

# Inequalities

### 3.1 Basics

Very similar to solving a normal equation with one caveat - every time you multiply by -1, you need to flip the sign.

**Example 3.1.1**  $5 - 3x \geq 21$

$$\begin{aligned}5 - 3x &\geq 21 \\ -3x &\geq 16 \\ 3x &\leq -16 \\ x &\leq -\frac{16}{3}\end{aligned}$$

**Example 3.1.2**  $17 + x \leq 32 + 3x \leq 21 + x$

$$\begin{aligned}17 + x &\leq 32 + 3x \leq 21 + x \\ 17 &\leq 32 + 2x \leq 21 \\ -15 &\leq 2x \leq -11 \\ -\frac{15}{2} &\leq x \leq -\frac{11}{2}\end{aligned}$$

### 3.2 Set Notation


Whilst we can use  $\leq$  and the other signs, we can also use set notation, which makes some easier, and is especially useful for quadratics.

$$\begin{array}{llll}x > 4 & \rightarrow & \{x : x < 4\} & \equiv & x \in (4, \text{inf}) \\ -15 \leq x \leq -11 & \rightarrow & \{x : -15 \leq x \leq -11\} & \equiv & x \in [-15, -11]\end{array}$$

### 3.3 Quadratics


The difficulty comes with quadratics which have multiple x-solutions. We need to draw a graph, and then check which way around which we should give our answer - a single set, or a union of 2.

**Example 3.3.1**  $2x^2 + 5x - 3 < 0$



$$\begin{aligned}
 2x^2 + 5x - 3 &< 0 \\
 (2x - 1)(x + 3) &< 0 \\
 -3 < x < \frac{1}{2} \\
 \left\{ x : -3 < x < \frac{1}{2} \right\}
 \end{aligned}$$

**Example 3.3.2**  $x^2 > 10 - 3x$



$$\begin{aligned}
 x^2 + 3x - 10 &> 0 \\
 (x + 5)(x - 2) &> 0 \\
 \{x : x < -5\} \cup \{x : x > 2\}
 \end{aligned}$$

## Chapter 5

# Coordinate Geometry


A line is a function which is straight, and they have operations that can be performed on them. If the function is not of first order, then they can have multiple gradients. All lines follow  $y = mx + c$ , where  $m$  is the gradient, and  $c$  is the y-intercept.

### 5.1 Characteristics of Lines


#### 5.1.1 Gradient

**Definition 5.1.1: Gradient**

The gradient of a line represents how shallow or deep it is.

**Definition 5.1.2: Perpendicular Lines**

One line is perpendicular to another if  $m_a = \frac{-1}{m_b}$ , ie they meet at  $90^\circ$




## 5.1.2 Y-Intercept

### Definition 5.1.3: Y Intercept

The Y-Intercept is where the line intersects the y-axis.

### Definition 5.1.4: Parallel Lines

One line is parallel to another if the gradients are identical, but the y-intercepts are different



## 5.1.3 Lines between points

### Definition 5.1.5: Midpoint

The midpoint of a line is the average between 2 points.



### Question 1: Midpoint between 2 points

**Example 5.1.1** Between (1, 4) and (5, 1)

$$\begin{aligned} &= \left( \frac{1+5}{2}, \frac{4+1}{2} \right) \\ &= (3, 2.5) \end{aligned}$$

### Definition 5.1.6: Distance

To find the distance between 2 points, use Pythagoras.

### Question 2: Distance between 2 points

**Example 5.1.2** Between (1, 4) and (5, 1)

$$\begin{aligned} c^2 &= a^2 + b^2 \\ c &= \sqrt{a^2 + b^2} \\ &= \sqrt{(5-1)^2 + (4-1)^2} \\ &= \sqrt{4^2 + 3^2} &= 5 \end{aligned}$$

## 5.2 Solving Line Equations

Mainly just a question of plugging in formulas and knowing the above. One other useful thing is this:

$$y - y_1 = m(x - x_1)$$

This equation can be used to construct a line at  $(x_1, y_1)$  given gradient  $m$ . This can also be used to find the y intercept.

## Chapter 6

# Circles

Circles all follow a formula:  $(x - a)^2 + (y - b)^2 = r^2$ .  $(a, b)$  is the centre of the circle, and  $r$  is the radius. Questions might ask you to describe a circle given a formula - just rearrange till you can get to the formula. When rearranging from the formula, be careful to preserve all solutions, as  $y$  is squared so you need to preserve negative values of  $y$ .

### 6.1 Intersections between lines and circles

Substitute into the circle equation. Make sure to check the discriminant or graph to check how many solutions exist.

**Question 3: Intersection between  $y = 2x + 1$  and  $(x - 3)^2 + (y + 1)^2 = 64$**

$$\begin{aligned}(x - 3)^2 + (2x + 1 + 1)^2 &= 64 \\ x^2 - 6x + 9 + 4x^2 + 8x + 4 &= 64 \\ 5x^2 + 2x - 51 &= 0\end{aligned}$$

$$2^2 - 4 * 5 * -51 > 0 \therefore 2 \text{ Intersections}$$

$$\begin{aligned}(5x + 17)(x - 3) &= 0 \\ x &= 3, -3.4 \\ y &= 2x + 1 \\ y &= 7, -5.7\end{aligned}$$

$$= (3, 7)(-3.4, -5.8)$$

## Chapter 8

# Binomial Expansion

## Chapter 9

# Trig Ratios

## Chapter 10

# Trig Functions

## Chapter 12

# Differentiation

## Chapter 13

# Integration

# Chapter 14

## Logs & Exponents

### 14.1 Logarithms

#### Definition 14.1.1: Logarithms

$$a^x = y \therefore \log_a y = x$$

Exponential Fact	Logarithm Fact
------------------	----------------

$10^3 = 1000$	$\log_{10} 1000 = 3$
---------------	----------------------

$5^4 = 625$	$\log_5 625 = 4$
-------------	------------------

$36^{\frac{1}{2}} = 6$	$\log_{36} 6 = \frac{1}{2}$
------------------------	-----------------------------

$2^{-3} = \frac{1}{8}$	$\log_2 \frac{1}{8} = -3$
------------------------	---------------------------

#### 14.1.1 Log Laws

#### Definition 14.1.2: Log & Index Laws

Laws of Indices	Laws of Logs
-----------------	--------------

$a^x * a^y = a^{x+y}$	$\log_a x + \log_a y = \log_a mn$
-----------------------	-----------------------------------

$a^x \div a^y = a^{x-y}$	$\log_a x - \log_a y = \log_a \frac{x}{y}$
--------------------------	--

$(a^x)^y = a^{xy}$	$\log_a x * \log_a y = \log_a m^n$
--------------------	------------------------------------



**Question 4:**  $\log_4 \frac{x}{x-1} = \log_4 3 + \log_4 2$

$$\log_4 \frac{x}{x-1} = \log_4 6$$

$$\frac{x}{x-1} = 6$$

$$x = 6x - 6$$

...

**Question 5:**  $\log_7 4x = \log_7 \frac{1}{x-6} + 1$

$$\log_7 \frac{4}{\frac{1}{x-6}} = 1$$

$$\log_7 4x(x-6) = 1$$

$$\log_7 4x^2 - 24x = 1$$

$$4x^2 - 24x = 7$$

...

**Question 6:**  $2^x = 75$

$$x = \log_2 75$$

$$x = 6.23$$

**Question 7:**  $5^{2x} - 6(5^x) = 0$

$$\text{let } y = 5^x$$

$$y^2 - 6y - 7 = 0$$

$$(y-7)(y+1) = 0$$

Logarithms always positive  $\therefore y = 7$

$$5^x = 7$$


$$x = \log_5 7$$

$$x = 1.21$$

**Question 8:**  $3^{2x+1} = 2^{5x}$ 

$$\begin{aligned}
 \log_e 3^{2x+1} &= \log_e 2^{5x} \\
 (2x+1) \log_e 3 &= (5x) \log_e 2 \\
 2(\log_e 3)x + \log_e 3 &= 5(\log_e 2)x \\
 \log_e 3 &= 5(\log_e 2)x - (2\log_e 3)x \\
 \log_e 3 &= x(5\log_e 2 - 2\log_e 3) \\
 x &= \frac{\log_e 3}{5\log_e 2 - 2\log_e 3} \\
 x &= 0.866
 \end{aligned}$$

## 14.2 Exponentials



Graphs with anything to the power of  $x$  always have a y-intercept of 1, because anything to the power of 0 is equal to 1.

### 14.2.1 Euler's Number - $e$

**Definition 14.2.1:**  $e$ 

$e$  is defined as having the following characteristics:

- $y = e^x, \frac{\Delta y}{\Delta x} = e^x$
- $y = e^{kx}, \frac{\Delta y}{\Delta x} = k e^{kx}$

**Question 9:**  $y = Ae^{kx}$ , and passes through  $(0, 6)$  and  $1, 9$ . Find  $A$  and  $k$

$$6 = Ae^0$$

$$A = 6$$

$$9 = 6e^k$$

$$e^k = \frac{9}{6}$$

$$k = \ln \frac{9}{6}$$

$$k = 0.41$$

## 14.3 Modelling With Exponentials

All Exponentials can be written with  $e$ , which makes calculus much easier.

Given  $y = Ae^{kx}$ , if  $k > 0$  we get exponential growth, and if  $k < 0$  then we get exponential decay.

## Part II

# Other Bits

# Chapter 1

## Transforming & Sketching Graphs

### 1.1 Graph Appearances

#### 1.1.1 Transformations

##### Definition 1.1.1: Translation

- The graph of  $f(x - a)$  is the graph of  $f(x)$  translated right by  $a$  units.
- The graph of  $f(x) + b$  is the graph of  $f(x)$  translated upwards by  $b$  units.


##### Definition 1.1.2: Scaling


###### Note

Never say shrink: always say stretch by a factor  $e$  where  $|e| < 1$

- The graph of  $cf(x)$  is the graph of  $f(x)$  stretched vertically by a factor of  $c$ .
- The graph of  $f(dx)$  is the graph of  $f(x)$  stretched horizontally by a factor of  $d^{-1}$ .

##### Example 1.1.1 Basic Graph Transformations





### 1.1.2 Combining Transformations

#### Question 10: Combining Transformations

$$y = f(-2x)$$

This is obtained from  $f(x)$  by doing the following:

1. Flip horizontally.
2. Stretch horizontally by a factor of 0.5.

#### Question 11: Combining Transformations


$$y = cf\left(\frac{1}{a} * (x - b)\right) + d$$

This is obtained from  $f(x)$  by doing the following:

1. Shift to the right  $b$  units.
2. Stretch horizontally by a factor of  $a$ .
3. Stretch vertically by a factor of  $c$ .
4. Shift upwards by  $d$  units.

### 1.1.3 Graph Shapes

#### Example 1.1.2 Basic Power Graphs



These are all of the basic graph shapes, and can be transformed just like  $y = x^2$  above.

## 1.2 Solving Using Graphs

Find one or more  $y = \dots$  equation, plot it, find the  $x$  position of any intercepts. If only one equation, find intersections with  $y = 0$ .

### Question 12: Solving $5 = 6x + 8$

$$5 = 6x + 8 \quad (1.1)$$

$$-3 = 6x \quad (1.2)$$

$$(1.3)$$


#### Example 1.2.1 Using Algebra

$$6x = -3 \quad (1.4)$$

$$x = \frac{6}{-3} \quad (1.5)$$

$$x = -\frac{1}{2} \quad (1.6)$$

#### Example 1.2.2 Using a Graph



Whilst this might seem less useful for basic equations, this can become much more useful for more complicated questions like below.



**Question 13: Finding the intersection of  $y = 3x^2 - 2x - 21$  and  $y = (x - 3)(x + 3)$**

**Example 1.2.3** Using Algebra

1. Set equal to each other
2. Simplify
3. Check how many roots exist using the discriminant
4. Work out all roots (possibly using factor theorem which can take a while)

$$3x^2 - 2x - 21 = (x - 3)(x + 3) \quad (1.7)$$

$$3x^2 - 2x - 21 = x^2 - 9 \quad (1.8)$$

$$2x^2 - 2x - 12 = 0 \quad (1.9)$$


$$x^2 - x - 6 = 0 \quad (1.10)$$

$$(x - 3)(x + 2) = 0 \quad (1.11)$$

$$x = 3, -2 \quad (1.12)$$

**Example 1.2.4** Using a Graph

1. Plot
2. Check intersections.



Intersects at  $x = -2, 3$ .

## Chapter 2

# Polynomial Division, Factor Theorem & Cubics

## 2.1 Polynomial Division

Any expression  $\frac{f(x)}{g(x)}$  can be expressed as  $g(x) \text{ rem } r(x)$ . For example,  $\frac{11}{4} \equiv 2 \text{ rem } 3$ .

There are 2 main methods of dividing polynomials - Long Division and Synthetic. Synthetic is usually considered easier, but a question might ask for Long Division, so learn both.

**Question 14:**  $x^3 - 17x + 6 \div x - 3$

### Example 2.1.1 Synthetic Division

Synthetic Division isn't very hard, but has steps you need to carefully follow. Firstly, copy all of the coefficients into the top row, and the negative of the divisor constant into the left column in the row below. For the first coefficient, copy it directly to the bottom. Then, for all of the others follow these steps:

1. Multiply the result from the order above by the negative of the constant, and copy it to the middle row.
2. Add that to the original and place the result in the bottom row.

$$\begin{array}{r|rrrr} & 1 & 0 & -17 & 6 \\ 3 & & 3 & & 9 & -24 \\ \hline & 1 & 3 & -8 & -18 \end{array}$$

$$\therefore \frac{x^3 - 17x + 6}{x - 3} = x^2 + 3x - 8 \text{ rem } -18$$

### Example 2.1.2 Long Division

Long Division involves a few main steps, which you repeat for every term of the dividend. It starts with laying out the equation in the box as you would for normal long division, and then you do the following:

1. Divide the dividend term by the divisor term an order below, and add that to the result at the top
2. Copy the dividend term a row below
3. Multiply the next dividend term by the negative of the constant in the divisor.
4. Treat what you've written down as long subtraction.
5. Copy the rest of the row down into the results bit.

Continue until you get to a lone constant, and that is the remainder

$$\begin{array}{r} x^2 + 3x - 8 \\ x - 3 \overline{) x^3 \phantom{+ 3x^2} - 17x + 6} \\ \underline{-x^3 + 3x^2} \phantom{+ 6} \\ 3x^2 - 17x \phantom{+ 6} \\ \underline{-3x^2 + 9x} \phantom{+ 6} \\ -8x + 6 \\ \underline{8x - 24} \\ -18 \end{array}$$

$$\therefore \frac{x^3 - 17x + 6}{x - 3} = x^2 + 3x - 8 \text{ rem } -18$$

## 2.2 Factor Theorem

The factor theorem is a method of solving cubic equations.

### Definition 2.2.1: Factor Theorem

Given  $f(x)$ , if  $f(y)$  is a solution, then  $(x - y)$  is a factor of  $f(x)$

To solve an equation using factor theorem, usually we need to follow a few steps:

1. Work through factors of the constant until you find one that is a factor of the whole equation.
2. Create a trial factorising function.
3. Expand the faux-factorised function, and equate coefficients.
4. Write a proper expanded function.
5. Properly factorise.

**Question 15:** Given one solution is an integer, solve  $2x^3 + x^2 - 18x + 9$

**Example 2.2.1** Solving with Factor Theorem

$$f(x) = 2x^3 + x^2 - 18x - 9$$

$$f(1) = 2 + 1 - 18 - 9 \neq 0$$

$$f(-1) = 2 + 1 + 18 - 9 \neq 0$$

$$f(3) = 2 * 27 + 9 - 54 - 9 = 0$$

$$f(3) = 0 \quad \therefore (x - 3) \text{ is a factor}$$

$$f(x) = (x - 3)(2x^2 + bx + 3)$$

$$= \dots - 6x^2 + bx^2 + \dots$$

$$b - 6 = 1$$

$$b = 7$$

$$f(x) = (x - 3)(2x^2 + 7x + 3)$$

$$= (x - 3)(2x + 1)(x + 3)$$

$$\therefore x = -3, \frac{1}{2}, 3$$

Part III

Book 2

## Chapter 1

# Algebraic Methods

## Chapter 2

# Functions & Graphs

## Chapter 6

# Trig Functions



## Chapter 7

# Trig Modelling

## Chapter 8

# Parametric Equations

## Chapter 9

# Differentiation

## Chapter 11

# Integration