

$$1) x = \frac{3}{2} \ln(y^2 + 9)$$

$$\frac{dx}{dy} = \frac{3}{2} \times \frac{1}{y^2 + 9} \times 2y$$

$$= \frac{3y}{y^2 + 9}$$

$$\frac{dx}{dy} = \frac{y^2 + 9}{3y}$$

$$= \frac{y^2}{3y} + \frac{9}{3y}$$

$$= \frac{y}{3} + \frac{3}{y}$$

C3 Z

Jack Maguire

Question 1

$$x = \ln(y^2 + 9)^{\frac{3}{2}}$$

$$= (\ln(y^2 + 9))^{\frac{3}{2}}$$

$$u = \ln(y^2 + 9)$$

$$\frac{\Delta u}{\Delta x} = \frac{2y}{y^2 + 9} \quad \text{Log Laws}$$

$$\frac{\Delta y}{\Delta x} = \frac{3}{2} (\ln(y^2 + 9))^{\frac{1}{2}} \frac{2y}{y^2 + 9}$$

$$= \frac{6y}{2y^2 + 18} \sqrt{\ln(y^2 + 9)}$$

Question wrong? Cannot get to simplify and desmos disagrees

Question 2

$$\frac{\sin 2\phi}{\sin \phi} - \frac{\cos 2\phi}{\cos \phi} \equiv \sec \phi$$

$$\text{LHS} \equiv \frac{\sin 2\phi}{\sin \phi} - \frac{\cos 2\phi}{\cos \phi}$$

$$\equiv \frac{2 \sin \phi \cos \phi}{\sin \phi} - \frac{2 \cos^2 \phi - 1}{\cos \phi}$$

$$\equiv 2 \cos \phi - 2 \cos \phi + \sec \phi \equiv \sec \phi \equiv \text{RHS} \quad \text{QED}$$



Question 3


α

$$v = 1 + 2 \ln x$$

$$v' = \frac{2}{x}$$

$$u = x$$

$$u' = 1$$

$$\begin{aligned} \frac{\Delta y}{\Delta x} &= \frac{vu' - v'u}{v^2} \\ &= \frac{1 + 2 \ln x - \frac{2x}{x}}{(1 + 2 \ln x)^2} \\ &= \frac{2 \ln x - 1}{1 + 4 \ln x + 4(\ln x)^2} \\ 0 &= \frac{2 \ln x - 1}{1 + 4 \ln x + 4(\ln x)^2} \\ 0 &= 2 \ln x - 1 \\ \frac{1}{2} &= \ln x \\ x &= e^{\frac{1}{2}} = \sqrt{e} \\ y &= \frac{x}{1 + 2 \ln x} \\ &= \frac{\sqrt{e}}{1 + 1} \\ &= \frac{1}{2} \sqrt{e} \\ \text{TP} &= (\sqrt{e}, \frac{1}{2} \sqrt{e}) \end{aligned}$$


b

$$v = (1 + 2 \ln x)^2$$

$$v' = \frac{4 + 8 \ln x}{x}$$

$$u = 2 \ln x - 1$$

$$u' = \frac{2}{x}$$

$$\frac{\Delta y}{\Delta x} = \frac{vu' - v'u}{v^2}$$

$$\frac{\Delta^2 y}{\Delta x^2} = \frac{\frac{2}{x}(1 + 2 \ln x)^2 - \frac{4+8 \ln x}{x}(2 \ln x - 1)}{(1 + 2 \ln x)^4}$$

$$\frac{\Delta^2 y}{\Delta x^2 \sqrt{e}} = \frac{\frac{2}{\sqrt{e}}(1 + 1)^2 - \frac{4+4}{\sqrt{e}}(1 - 1)}{(1 + 1)^4}$$

$$= \frac{\frac{2}{\sqrt{e}}}{(1 + 1)^2}$$

$$= \frac{2}{4\sqrt{e}}$$

$$= \frac{1}{2\sqrt{e}}$$

$\frac{1}{2\sqrt{e}} > 0 \therefore$ is a minimum point. ✓

Question 4

a

$$x = \tan y$$

$$\frac{\Delta x}{\Delta y} = \sec^2 y = 1 + \tan^2 y = 1 + x^2$$

$$\therefore \frac{dy}{dx} = \frac{1}{1+x^2}$$

$$\frac{\Delta y}{\Delta x} = \cos^2 y$$

$$= \cos^2 (\tan^{-1} x)$$

???

b

$$\frac{\Delta y}{\Delta x} = 1 + x^2 - \frac{8}{1+x^2} - 6x$$

$$= x^2 - 6x - \frac{8}{1+x^2} + 1$$

$$0 = x^2 - 6x - \frac{8}{1+x^2} + 1$$

$$= (1+x^2)x^2 - (1+x^2)6x - 8 + (1+x^2)$$

$$= x^2 + x^4 - 6x - 6x^3 - 8 + 1 + x^2$$

$$= x^4 - 6x^3 + 2x^2 - 6x - 7$$

???

$$\frac{dy}{dx} = \frac{1}{x^2+1} - 9x \cdot \frac{1}{1+x^2} \times 2x - 6x$$

$$= \frac{1}{1+x^2} - \frac{18x^2}{1+x^2} - 6x$$

$$0 = \frac{1-18x^2}{1+x^2} - 6x$$

$$6x + 6x^3 = 1 - 18x^2$$

$$6x^3 + 18x^2 - 6x - 1 = 0$$

c

$$f(x) = 6x^3 + 14x - 1 \quad \text{value}$$

$$f(0) = -1$$

$$f(1) = 19$$

Between positive and negative \therefore root in between. ✓

d

???

NL

e

???

..

Question 5

a

$$H = k \left(1 - e^{-\frac{1}{12}t} \right)$$

$$41 = k \left(1 - e^{-\frac{5}{12}} \right)$$

$$\frac{41}{k} = 1 - e^{-\frac{5}{12}}$$

$$k = \frac{41}{1 - e^{-\frac{5}{12}}}$$

$$k = 120.319 \dots$$

$$= 120$$



b

$$H = 120 \left(1 - e^{-\frac{1}{12}t} \right)$$

$$\frac{90}{120} = 1 - e^{-\frac{1}{12}t}$$

$$\frac{1}{4} = e^{-\frac{1}{12}t}$$

$$\ln \frac{1}{4} = -\frac{1}{12}t$$

$$-12t = -\ln 4$$

$$t = 12 \ln 4$$

$$= 24 \ln 2$$



c

$$H = 120 \left(1 - e^{-\frac{1}{12}t} \right)$$

$$\frac{H}{120} = 1 - e^{-\frac{1}{12}t}$$

$$1 - \frac{H}{120} = e^{-\frac{1}{12}t}$$

$$-\frac{1}{12}t = \ln \left(1 - \frac{H}{120} \right)$$

$$t = -12 \ln \left(1 - \frac{H}{120} \right)$$

$$H = 120 \left(1 - e^{-\frac{1}{12}t} \right)$$

$$a = 1 - e^{-\frac{1}{12}t}$$

$$\frac{\Delta a}{\Delta t} = \frac{1}{12} e^{-\frac{1}{12}t}$$

$$\frac{\Delta H}{\Delta t} = 120 \frac{1}{12} e^{-\frac{1}{12}t}$$

$$= 10 e^{-\frac{1}{12}t}$$

$$= 10 e^{-12 \ln \left(1 - \frac{H}{120} \right)}$$

$$= \frac{-10}{12} H$$

$$\begin{aligned} \frac{H}{120} &= 1 - e^{-\frac{1}{12}t} \\ e^{-\frac{1}{12}t} &= 1 - \frac{H}{120} \\ 10 e^{-\frac{1}{12}t} &= 10 - \frac{H}{12} \end{aligned}$$

d

$$\begin{aligned} \frac{\Delta H}{\Delta t} &= \frac{-10}{12} H \frac{\Delta H}{\Delta t} \cdot 0.5 &= -\frac{10}{12} 7.5 \\ &= \frac{-25}{4} &\text{ECF} \end{aligned}$$

e

$$\frac{\Delta H}{\Delta t} = 120 \frac{1}{12} e^{-\frac{1}{12}t}$$

$$0 = 10 e^{-\frac{1}{12}t}$$

$$e^{-\frac{1}{12}t} = 0$$

$$\begin{aligned} t \rightarrow \infty, e^{-\frac{1}{12}t} &\rightarrow 0 \\ \therefore H &\rightarrow 120 \\ t &= 120 \end{aligned}$$

Since $\mathbb{R} \not\subseteq \ln 0$, we can see that this model exponentially grows, that is, there is no maximum height.

Question 6

a

$$2 \leq f(x) \leq 4$$

$$a = 3$$

$$\pi \rightarrow 2\pi \therefore b = 0.5$$

b

$$y = 3 + \cos \frac{x}{2}$$

$$x = 3 + \cos \frac{y}{2}$$

$$\cos \frac{y}{2} = x - 3$$

$$f^{-1}(x) = 2 \cos^{-1}(x - 3)$$

c

$$2 \leq x \leq 4$$

$$0 \leq f(x) \leq 2\pi$$

d

$$\frac{\Delta y}{\Delta x} = -\frac{1}{2} \sin \frac{x}{2} \frac{\Delta y}{\Delta x} \frac{4}{3}\pi$$

$$= -\frac{1}{2} \sin \frac{2\pi}{3}$$

$$= -\frac{1}{2} \frac{\sqrt{3}}{2}$$

$$= -\frac{\sqrt{3}}{4}$$

e

$$-\frac{\sqrt{3}}{4} \quad \times \text{reciprocal}$$

$$\begin{aligned} 71) \quad & \sin(A+B) + \sin(A-B) \\ &= 2 \sin A \cos B \\ & \underline{A+B = P} \\ & \underline{A-B = Q} \\ & A = \frac{P+Q}{2}, \quad B = \frac{P-Q}{2} \\ \therefore \sin P + \sin Q &= 2 \sin \frac{P+Q}{2} \cos \frac{P-Q}{2} \end{aligned}$$

$$\begin{aligned} 76) \quad & 2 \sin \left(\frac{50+\theta}{2} \right) \cos \left(\frac{50-\theta}{2} \right) = \sin 30 \\ & 2 \sin 30 \cos 20 = \sin 30 \\ & \sin 30 (2 \cos 20 - 1) = 0 \\ & \text{solve.} \\ & = 0, 30, 60, 120, 150, 180 \end{aligned}$$

Question 7

a

$$\sin P + \sin Q \equiv 2 \sin \left(\frac{P+Q}{2} \right) \cos \left(\frac{P-Q}{2} \right)$$

$$\text{RHS} \equiv 2 \sin \left(\frac{P+Q}{2} \right) \cos \left(\frac{P-Q}{2} \right)$$

$$\equiv 2 \sin \left(\frac{P}{2} + \frac{Q}{2} \right) \cos \left(\frac{P}{2} - \frac{Q}{2} \right)$$

$$\equiv 2 \left(\sin \frac{P}{2} \cos \frac{Q}{2} + \sin \frac{Q}{2} \cos \frac{P}{2} \right) \left(\cos \frac{P}{2} \cos \frac{Q}{2} + \sin \frac{P}{2} \sin \frac{Q}{2} \right)$$

$$\equiv 2 \left(\sin \frac{P}{2} \cos \frac{P}{2} \cos^2 \frac{Q}{2} + \sin^2 \frac{P}{2} \sin \frac{Q}{2} \cos \frac{Q}{2} + \sin \frac{Q}{2} \cos^2 \frac{P}{2} \cos \frac{Q}{2} + \sin \frac{P}{2} \sin^2 \frac{Q}{2} \cos \frac{P}{2} \right)$$

???

$$\equiv \sin P + \sin Q \equiv \text{RHS} \quad \text{QED}$$

b

$$\sin \theta - \sin 3\theta + \sin 5\theta = 0???$$

Question 8

278

