

Fowler's theorem for involutions.

Sylvain Cappell, S. Weinberger, and M. Yan

Fowler, in his Ph.D. thesis, proved that if Γ is a uniform lattice in a real semisimple group with odd torsion in Γ then there is no compact closed manifold M whose universal cover is rationally acyclic. A proof can be found in [W2]. We show that the same is true for Γ with 2-torsion.

Without loss of generality (by considering a normal subgroup of finite index), it suffices to prove this for the special case where $\Gamma = \pi \rtimes \mathbf{Z}_2$ for a torsion free group π , a lattice in G , for which there is an involution on $M = K \backslash G / \pi$ (by isometries with the locally symmetric metric) whose fixed set F is not empty. (F might be disconnected; for simplicity we will write what follows just for the connected case – there are no differences in the general case.)

Now suppose that X^m is a manifold with fundamental group Γ , Y its 2-fold cover, and suppose that the universal cover of X (and therefore Y) are rationally acyclic. We will consider the symmetric signatures of Y in the (symmetric = quadratic L-group) $L(\mathbf{R}\pi)$, where \mathbf{R} is the real numbers. There is an equivalence $f: Y \rightarrow M$ which (while not degree one) gives an equivalence of symmetric signatures (because over \mathbf{R} , all degrees have square roots, so the symmetric signature is only sensitive to the sign of the degree of the map). Since the Novikov conjecture is true for π , the assembly map from $H_m(B\pi; L(\mathbf{R})) \rightarrow L_m(\mathbf{R}\pi)$ is injective, and this detects in the degree m piece $H_m(B\pi; \mathbf{Z})$ the class that these manifolds represent in group homology. It follows that this map is degree one. $f_*[Y] = [M]$.

Now we use a cobordism argument from [W1]. We now consider the image of the fundamental class of any manifold Z with fundamental group π involution inducing this automorphism of π and the image of $[Z]$ in $H_m(B\Gamma; \mathbf{Z}_2)$. It follows from standard equivariant homotopy theory that Z has an equivariant map, g , to M , and thus there is a map from its fixed set $Z^{\mathbf{Z}_2} \rightarrow F$. We claim that $g_*[Z] = g_*[Z^{\mathbf{Z}_2}]$ where we make use of the map from $\mathbf{Z}_2 \times \pi_1 F \rightarrow \Gamma$ (and the periodicity on the group homology of \mathbf{Z}_2 to raise the dimension from that of F to $\dim M$).

This cobordism is between Z and a projective space bundle over \mathbf{Z}^2 - namely the projectivized normal bundle to \mathbf{Z}^2 . (The fundamental class of the latter is the desired element by the Leray-Hirsch theorem.) It is explicitly $Z \times [0, 1]$ and on $Z \times \{1\}$ mod out in the complement of the equivariant regular neighborhood of \mathbf{Z}^2 the $\mathbf{Z}/2$ action.

Thus for Y , this image is 0, since the action is free. For M however, this is always nonzero. The action by \mathbf{Z}_2 by isometries has fixed set which is aspherical and indeed the Borel

construction for the action on M shows that $Z_2 \times F \rightarrow \Gamma$ induces an injection on homology in dimension $\dim(M/Z_2)$ (and an isomorphism in higher dimensions, see [B]). Since the fundamental class of an aspherical manifold is always nontrivial in its group homology, we have a contradiction.

References

- [B] A. Borel, A seminar on transformation groups, Princeton University Press 1960
- [W1] S. Weinberger, Group actions and higher signatures II, CPAM 1987
- [W2] S. Weinberger, Variations on a theorem of Borel, Cambridge University Press 2022