

Multi-domain Bivariate Spectral Local Linearisation method for solving non-similar boundary layer partial differential equations

MAGAGULA VUSI MPENDULO

Supervisors: Prof. S.S. Motsa & Prof. P. Sibanda



The 40th South African Symposium of Numerical and Applied Mathematics

22 - 24 March 2016



Overview

- 1 AIM
- 2 SOLUTION PROCEDURE
- 3 RESULTS AND DISCUSSION
- 4 CONCLUSION & FUTURE RESEARCH DIRECTION
- 5 REFERENCES

Aim & Objectives

Aim

To present an extension of the **Bivariate Spectral Local Linearisation Method (BSLLM)**[1] for solving non-similar nonlinear PDEs over **large time intervals**.

- BSLLM uses Chebyshev-Gauss-Lobatto points (see [3, 4])
- Drawback of BSLLM - accuracy deteriorates over **large time intervals**.
- New approach termed - **Multi-domain Bivariate Spectral Local Linearisation Method (MD-BSLLM)**

Objectives

- Solve non-linear non-similar boundary layer equations over a large time domain using the MD-BSLLM.
- Validate the results using a series solution approach.

Solution Method

The solution approach involves

- Domain decomposition
- linearisation and decoupling
- bivariate interpolation.
- pseudo-spectral approximation

Numerical Experiment

$$\begin{aligned} \frac{\partial^3 f}{\partial \eta^3} + \frac{1}{4}(n+3)f \frac{\partial^2 f}{\partial \eta^2} - \frac{1}{2}(n+1) \left(\frac{\partial f}{\partial \eta} \right)^2 + \zeta \frac{\partial^2 f}{\partial \eta^2} + (1-w)g + wh \\ = \frac{1}{4}(1-n)\zeta \left[\frac{\partial f}{\partial \eta} \frac{\partial^2 f}{\partial \zeta \partial \eta} - \frac{\partial^2 f}{\partial \eta^2} \frac{\partial f}{\partial \zeta} \right], \end{aligned} \quad (1)$$

$$\frac{1}{Pr} \frac{\partial^2 g}{\partial \eta^2} + \frac{1}{4}(n+3)f \frac{\partial g}{\partial \eta} + \zeta \frac{\partial g}{\partial \eta} = \frac{1}{4}(1-n)\zeta \left[\frac{\partial f}{\partial \eta} \frac{\partial g}{\partial \zeta} - \frac{\partial g}{\partial \eta} \frac{\partial f}{\partial \zeta} \right], \quad (2)$$

$$\frac{1}{Sc} \frac{\partial^2 h}{\partial \eta^2} + \frac{1}{4}(n+3)f \frac{\partial h}{\partial \eta} + \zeta \frac{\partial h}{\partial \eta} = \frac{1}{4}(1-n)\zeta \left[\frac{\partial f}{\partial \eta} \frac{\partial h}{\partial \zeta} - \frac{\partial h}{\partial \eta} \frac{\partial f}{\partial \zeta} \right], \quad (3)$$

subject to

$$f(\zeta, 0) = 0, \quad \frac{\partial f}{\partial \eta}(\zeta, 0) = 0, \quad g(\zeta, 0) = h(\zeta, 0) = 1, \quad \frac{\partial f}{\partial \eta}(\zeta, \infty) = g(\zeta, \infty) = h(\zeta, \infty) = 0.$$

- **Formulated and Solved** by **Hussain et.al.** [2] using finite-difference based **Keller-box technique**
- Results were validated using **Series Solutions**, for small and large ζ

Domain Decomposition




Figure: Multi-domain Grid

- The patching condition requires that

$$f^{(l)}(\eta, \zeta_{l-1}) = f^{(l-1)}(\eta, \zeta_{l-1}), \quad \eta \in [a, b], \quad (4)$$

where $f^{(l)}(\eta, \zeta)$ denotes the solution of equation (2) at each sub-interval Γ_l

Linearisation and Decoupling

- The PDEs are linearised using the **quasi-linearisation method**.
 - an iterative procedure based on the **Taylor-series expansion** about the previous estimates of the solution.
- Applying the **quasi-linearisation method** on one function at a time, results in three **decoupled linear PDEs**.
- In each sub-interval $[\zeta_{l-1}, \zeta_l]$, these decoupled linear PDEs can be solved iteratively in a sequential manner until the desired solution is obtained.

$$\beta_{0,r} \frac{\partial^3 f_{r+1}^{(l)}}{\partial \eta^3} + \beta_{1,r} \frac{\partial^2 f_{r+1}^{(l)}}{\partial \eta^2} + \beta_{2,r} \frac{\partial f_{r+1}^{(l)}}{\partial \eta} + \beta_{3,r} f_{r+1}^{(l)} + \beta_{4,r} \frac{\partial f_{r+1}^{(l)}}{\partial \xi} + \beta_{5,r} \frac{\partial f_{r+1}^{\prime(l)}}{\partial \xi} = R_{f,r}^{(l)}, \quad (5)$$

$$\sigma_{1,r} \frac{\partial^2 g_{r+1}^{(l)}}{\partial \eta^2} + \sigma_{2,r} \frac{\partial g_{r+1}^{(l)}}{\partial \eta} + \sigma_{3,r} g_{r+1}^{(l)} + \sigma_{4,r} \frac{\partial g_{r+1}^{(l)}}{\partial \xi} = R_{g,r}^{(l)}, \quad (6)$$

$$\omega_{1,r} \frac{\partial^2 h_{r+1}^{(l)}}{\partial \eta^2} + \omega_{2,r} \frac{\partial h_{r+1}^{(l)}}{\partial \eta} + \omega_{3,r} h_{r+1}^{(l)} + \omega_{4,r} \frac{\partial h_{r+1}^{(l)}}{\partial \xi} = R_{h,r}^{(l)}, \quad (7)$$

Pseudo-spectral Approximation

- Assume that the solution at each sub-interval Γ_l , denoted by $f^{(l)}(\eta, \zeta)$, can be approximated by a **bivariate Lagrange interpolation** polynomial of the form

$$f^{(l)}(\eta, \zeta) \approx \sum_{i=0}^{N_x} \sum_{j=0}^{N_t} f^{(l)}(\eta_i, \zeta_j) \mathcal{L}_i(\eta) \mathcal{L}_j(\zeta). \quad (8)$$

- Transpiration parameter derivative values are computed at the grid points (η_i, ζ_j) :

$$\left. \frac{\partial f^{(l)}}{\partial \zeta} \right|_{(\eta_i, \zeta_j)} = \left(\frac{2}{\zeta_l - \zeta_{l-1}} \right) \sum_{v=0}^{N_t} d_{jv} f^{(l)}(\eta_j, \zeta_v) = \left(\frac{2}{\zeta_l - \zeta_{l-1}} \right) \sum_{v=0}^{N_t} d_{jv} \mathbf{F}_v^{(l)} \quad (9)$$

- The n th order space derivative is defined as

$$\left. \frac{\partial^n f^{(l)}}{\partial \eta^n} \right|_{(\eta_i, \zeta_j)} = \left(\frac{2}{\eta_\infty} \right)^n \sum_{\rho=0}^{N_x} D_{i\rho}^n f^{(l)}(\eta_\rho, \zeta_j) = \left(\frac{2}{\eta_\infty} \right)^n \mathbf{D}^n \mathbf{F}_j^{(l)}, \quad j = 0, 1, 2, \dots, N_t, \quad (10)$$

where the vector $\mathbf{F}_j^{(l)}$ is defined as

$$\mathbf{F}_j^{(l)} = [f^{(l)}(\eta_0, \zeta_j), f^{(l)}(\eta_1, \zeta_j), \dots, f^{(l)}(\eta_{N_x}, \zeta_j)]^T. \quad (11)$$

Pseudo-spectral Approximation

- Substituting equations (9) and (10) into equation (5), we get

$$\left[\beta_{0,r} \mathbf{D}^3 + \beta_{1,r} \mathbf{D}^2 + \beta_{2,r} \mathbf{D} + \beta_{3,r} \right] \mathbf{F}_{r+1,j}^{(l)} + \beta_{4,r} \sum_{v=0}^{N_t} d_{jv} \mathbf{F}_{r+1,v}^{(l)} + \beta_{5,r} \sum_{v=0}^{N_t} d_{jv} \mathbf{D} \mathbf{F}_{r+1,v}^{(l)} = \mathbf{R}_{f,r}^{(l)}, \quad (12)$$

for $j = 0, 1, 2, \dots, N_t$.

- The patching condition requires that

$$f_{r+1}^{(l)}(\eta_i, \zeta_{(l-1,j)}) = f_{r+1}^{(l-1)}(\eta_i, \zeta_{(l-1,j)}), \quad \eta \in [a, b], \quad (13)$$

- The initial unsteady solution of equation (5) when $\zeta = 0$ corresponds to $t = t_{N_t} = -1$.
- Equation (12) is evaluated for $j = 0, 1, \dots, N_t - 1$

$$\left[\beta_{0,r} \mathbf{D}^3 + \beta_{1,r} \mathbf{D}^2 + \beta_{2,r} \mathbf{D} + \beta_{3,r} \right] \mathbf{F}_{r+1,j}^{(l)} + \beta_{4,r} \sum_{v=0}^{N_t-1} d_{jv} \mathbf{F}_{r+1,v}^{(l)} + \beta_{5,r} \sum_{v=0}^{N_t-1} d_{jv} \mathbf{D} \mathbf{F}_{r+1,v}^{(l)} = \mathbf{R}_{1,j}^{(l)}, \quad (14)$$

and

$$\mathbf{R}_{1,j}^{(l)} = \mathbf{R}_{f,r}^{(l)} - \beta_{4,r} d_{jN_t} \mathbf{F}_{N_t}^{(l)} - \beta_{5,r} d_{jN_t} \mathbf{D} \mathbf{F}_{N_t}^{(l)}.$$

- Imposing boundary conditions for $j = 0, 1, \dots, N_t - 1$, equation (14) can be expressed as the following

$$N_t(N_x + 1) \times N_t(N_x + 1)$$

matrix system

$$\begin{bmatrix} A_{0,0} & A_{0,1} & \cdots & A_{0,N_t-1} \\ A_{1,0} & A_{1,1} & \cdots & A_{1,N_t-1} \\ \vdots & \vdots & \ddots & \vdots \\ A_{N_t-1,0} & A_{N_t-1,1} & \cdots & A_{N_t-1,N_t-1} \end{bmatrix} \begin{bmatrix} \mathbf{F}_0^{(l)} \\ \mathbf{F}_1^{(l)} \\ \vdots \\ \mathbf{F}_{N_t-1}^{(l)} \end{bmatrix} = \begin{bmatrix} \mathcal{R}_{1,0}^{(l)} \\ \mathcal{R}_{1,1}^{(l)} \\ \vdots \\ \mathcal{R}_{1,N_t-1}^{(l)} \end{bmatrix}, \quad (15)$$

where

$$A_{i,i} = \beta_{0,r} \mathbf{D}^3 + \beta_{1,r} \mathbf{D}^2 + \beta_{2,r} \mathbf{D} + \beta_{3,r} \mathbf{I} + \beta_{4,r} d_{ii} \mathbf{I} + \beta_{5,r} d_{ii} \mathbf{D} \quad (16)$$

$$A_{i,j} = \beta_{4,r} d_{ij} \mathbf{I} + \beta_{5,r} d_{ij} \mathbf{D}, \quad \text{when } i \neq j, \quad (17)$$

Results

- Comparison of Multi-domain bivariate spectral local linearisation solution for the **skin friction** against the series solution for large ζ .

MD-BSLLM		Series Solution for large ζ
ζ	$f''(0, \zeta)$	$f''(0, \zeta)$
5	0.3088214	0.3088214
10	0.1547399	0.1547399
15	0.1031717	0.1031717
20	0.0773803	0.0773803
25	0.0619045	0.0619045
30	0.0515872	0.0515872
35	0.0442176	0.0442176
40	0.0386905	0.0386905

Table: $N_x = 60, N_t = 5, p = 20$

Results

- Comparison of Multi-domain bivariate spectral local linearisation solution for the **Sherwood number** against the series solution for large ζ

MD-BSLLM		Series Solution for large ζ
ζ	$-h'(0, \zeta)$	$-h'(0, \zeta)$
5	3.0018658	3.0018658
10	6.0002332	6.0002332
15	9.0000691	9.0000691
20	12.0000292	12.0000292
25	15.0000149	15.0000149
30	18.0000086	18.0000086
35	21.0000054	21.0000054
40	24.0000036	24.0000036

Table: $N_x = 60, N_t = 5, p = 20$

Results

- Comparison of Multi-domain bivariate spectral local linearisation solution for the **Nusselt number** against the series solution for large ζ

	MD-BSLLM	Series Solution for large ζ
ζ	$-g'(0, \zeta)$	$-g'(0, \zeta)$
5	3.5018961	3.5018961
10	7.0002370	7.0002370
15	10.5000702	10.5000702
20	14.0000296	14.0000296
25	17.5000152	17.5000152
30	21.0000088	21.0000088
35	24.5000055	24.5000055
40	28.0000037	28.0000037

Table: $N_x = 60, N_t = 5, P = 20$

Conclusion

- The MD-BSLLM method can be used to solve non-linear non-similar boundary layer equations over a large time domain.
- We were able to validate the results using a series solution approach.

Future Research Direction

- Solve different **types** of NPDEs arising from Physics, Mathematical Biology, etc.

References



Motsa, S.S. and Animasaun, I.L., 2015. A new numerical investigation of some thermo-physical properties on unsteady MHD non-Darcian flow past an impulsively started vertical surface. Thermal Science, 19(suppl. 1), 249-258.



Hussain S., Hossain M.A., Natural convection flow from a vertical permeable flat plate with variable surface temperature and species concentration, Engineering Computations, Vol. 17 No. 7, 2000, 789-812



Trefethen, Lloyd N. Spectral methods in MATLAB. Vol. 10. Siam, 2000.



Canuto C., Hussaini M. Y., Quarteroni A., and T. A. Zang, Spectral Methods, Evolution to Complex Geometries and Applications to Fluid Dynamics, Springer-Verlag, Berlin, 2007.

Thank you ...