Historical and Conditional Forecasts of Sampled Net-Position Peaks

Common targets, source-specific results and exploratory diagnostics

23 October 2026

Abstract

How well can a dealer forecast the largest absolute recorded net position before a reporting horizon ends? Such forecasts motivate evaluating a monitoring input for the measured position series. We compare historical peak constructions with ridge quantile regression and local quantiles on a market maker's FX client-flow series and a weekly Treasury-dealer aggregate. The common target includes the observed issue position and uses completed training labels. First, six-record FX regression lowers matched-history pinball loss by 14.77%, while breaches rise from 360 to 376. Second, the Treasury primary deteriorates by 50.79%, with breaches rising from 28 to 54. Third, lower pinball loss often accompanies more breach occasions. The evidence is specific to the inspected records, stipulated score and source-measurement conventions.

1. Introduction

Dealers absorb customer order imbalances while managing inventory risk and scarce balance-sheet capacity. Inventory affects optimal dealer pricing (Ho and Stoll, 1981), FX dealing (Lyons, 1995), and observed inventory adjustment in dealership markets (Hansch, Naik and Viswanathan, 1998). Treasury intermediation likewise depends on dealer capacity (Duffie et al., 2023). A forecast of the maximum recorded position magnitude over a future horizon may support monitoring of the measured position: an endpoint can return toward its starting value after an intervening large position. Translating that monitoring signal into a particular margin or financing decision requires the corresponding contractual and balance-sheet inputs.

We ask whether mature conditional estimators improve forecasts of an absolute sampled-position peak relative to simple historical constructions. Every report is evaluated against the same peak response and upper-quantile score on common issue origins. The endpoint-derived bound introduced below is a benchmark rather than a direct peak-quantile estimate. The historical alternatives use past headrooms, paths translated to the current signed position, or native peaks floored at the observed current magnitude. The conditional alternatives use ridge quantile regression, nearest-neighbor quantiles and scaled nearest-neighbor quantiles.

The matched-history FX regression comparison lowers mean loss by 14.77%, but increases breaches from 360 to 376. The original FX primary uses a different historical window and lowers loss by 12.66%. The separately declared Treasury primary deteriorates by 50.79%, with 54 rather than 28 breaches. Short-history FX regression also deteriorates at the longer horizon; a Treasury neighbor gain reverses on the sparse grid. These adverse outcomes are central findings.

The paper makes three contributions. First, it places established forecasting engines and historical constructions on a common completed-peak response with an explicit issue-known floor. Second, it controls completed-label pools and update schedules and retains the complete favorable and adverse model menu. Third, it documents how score, breach frequency, report coherence and fixed-report preferences can disagree across two financial measurement processes. The contribution is an application and model-validation benchmark using mature methods.

Section 2 locates the study in the literature. Sections 3–5 describe data, forecasts and evaluation. Sections 6–7 report FX and Treasury results. Section 8 collects exploratory robustness evidence, Section 9 discusses mechanisms and limitations, and Section 10 concludes. Detailed historical menus, geometry and supplementary diagnostics appear in the appendices.

2. Related literature

Wisniewski, Lindsay and Lindsay (2020) and the net-position forecasting chapter of Wisniewski (2024) are the closest empirical lineage. They already study conditional, neighbor and normalized endpoint forecasts on the FX source family, using lagged positions and calendar information. Here the response is a single upper quantile of a future absolute sampled maximum, including the current magnitude. Marginal endpoint intervals and their averaged losses address a different functional; the present comparison fits the established engines directly to completed peak labels.

Financial path extrema have a substantial prior literature. MaxVaR evaluates worst intermediate mark-to-market loss (Boudoukh et al., 2004). Liao and Theodosopoulos (2005) study the timing of mark-to-market for collateralized contingent credit exposure. Van den Hoogen (2017) models a conditional intraday liquidity buffer through the maximum cumulative payment deficit; Petrunia, Sabetti and Voia (2019) study a cross-participant end-of-day debit maximum.

Arjani, Li and Sabetti (2020) forecast remaining-day liquidity-demand ratios. These targets clarify the specific contribution of a sampled net-position envelope.

Regression quantiles (Koenker and Bassett, 1978), sequential neighbor quantiles (Biau and Patra, 2011), conditional historical filtering (Hull and White, 1998; Barone-Adesi et al., 1999), and quantile scoring (Gneiting, 2011) supply the mature methods. Quantile forests (Meinshausen, 2006), CAViaR (Engle and Manganelli, 2004), sequential conformal prediction (Xu and Xie, 2023), trajectory calibration (Zhou, Lindemann and Sesia, 2024) and expected-shortfall curves (Sözen, 2026) are further relevant alternatives. The executed estimator menu is deliberately bounded.

The resulting distinction is the measured peak response, known issue floor, completed-label timing, controlled history/update comparisons and qualified aggregate application. The benchmark measures how these mature procedures behave on the specified task; its supporting finite geometry explains construction differences.

3. Data

The FX input is the one-hour member of the Net Position Time Series Dataset, version 2: 18,576 records from 5 February 2014 01:00 through 24 January 2017 00:00. Wisniewski et al. (2020) describe aggregate client-flow positions across five liquid currency pairs for one market maker. We use datetime and the signed NetPosUsd scalar, rather than reconstructing raw currency positions, hedge transactions or cash movements. The source has 18,420 one-hour and 155 forty-nine-hour adjacent gaps. We preserve chronology without sorting, removing gap-crossing windows or interpolation. Recorded-clock and vintage limits are collected in Section 9.

The Treasury application uses the New York Fed Primary Dealer Statistics series PDPOSGS-B: Treasury-bill long positions less short positions, measured at weekly Wednesday close in market/fair-value USD millions. The accepted snapshot contains 706 records. It combines dealer inventories, valuation and reporting composition. This supplies an aggregate financial measurement task alongside the individual market-maker client-flow series.

Treasury validation issue dates are 24 July 2019 to 15 July 2020. The final H=24 validation label ends at index 404 on 30 December 2020, before evaluation begins at 405 on 6 January 2021. The last common issue is 681 on 22 April 2026; its H=6 label ends at 687 on 3 June, and H=24 at 705 on 7 October 2026. FX validation ends at origin 11902; its latest H=24 label ends at 11926, before evaluation origin 11927.

Table 1. Frozen source-specific chronological budgets. All indices are zero based; W and d count source records. H is 6 or 24 in every cell.

Source	Validation origins	Evaluation origins	W	Refit d
FX	10463–11902 (1440)	11927–18551 (6625)	720 / 2880	24
Treasury	329–380 (52)	405–681 (277)	52 / 260	13

All horizons share the source-specific evaluation issue origins. Horizon H counts subsequent source records, so the response uses H+1 observed positions including issue time. FX gaps are retained in chronological order. Treasury transitions follow the weekly economic-date clock. Exact data identities, access routes and reproduction commands are retained in the repository guide.

4. Target and forecasting methods

Notation. Distinct symbols for response, refits, history, distances and preferences.

Symbol	Definition
N_t; B_t	Signed position; observed issue magnitude $\lvert N_t\rvert$
$P_{t,H}$; $G_{t,H}$	Sampled absolute peak; headroom $P_{t,h}-B_t$
$\kappa(t)$; W; d	Most recent refit index; label-pool size; refit step
C_t^m; Floor	Issued report for method m; the known-floor report
d_{abs}; d_{signed}	Common-set anchor-distance radii
$\tilde{N}_{i,h}^{tr}$; $M_{i,h}$	Translated historical path; AR residual-replay path
a_κ; $z_{i,\kappa}$	Training response scale; standardized predictors
a; $\omega=a/(1-a)$	Shortfall weight; shortfall/excess preference ratio
J_λ; D_λ	Penalized regression objective; feasible dual objective

Let N denote the signed source-reported net position. The issue magnitude B is observed when the report is issued. Define the sampled peak P and its nonnegative headroom G by

$$B_t=|N_t|;\quad P_{t,H}=\max_{0\le h\le H}|N_{t+h}|;\quad G_{t,H}=P_{t,H}-B_t\ge 0.$$

At issue t, the most recent scheduled refit is κ(t). Every training label is complete by that refit, and the next refit updates the transform, scale, coefficients and historical pool together:

$\kappa(t)=t_0+d\,\lfloor (t-t_0)/d \rfloor; \quad T_{\kappa,H,W}=\{\kappa-H-W+1,\ldots,\kappa-H\}.$

For a completed episode i, let u and l be its largest and smallest increments from its starting position, including the zero increment. Translating its ordered path to $x=N_t$ produces $V_{i,t}=\max\{x+u_i,-x-l_i\}$. On a common episode set A, Q selects the duplicate-preserving corrected empirical order statistic. The historical forecasts are

$C_t^F=B_t+Q\{G_{i,H}:i\in A\}; \quad C_t^S=B_t+Q\{V_{i,t}-B_t:i\in A\}; \quad C_t^K=\max\{B_t,Q\{P_{i,H}:i\in A\}\}.$

Rolling controls update their completed-label set at each issue. Frozen controls use the same completed-label pool and refit schedule as the fitted procedures, while their current issue anchor continues to update. Earlier one-state controls retrieve nearby signed starting positions; their complete earlier menus remain in Appendix B.

The 33 raw coordinates are N(t) through N(t−24), current magnitude, short/long past increment RMS, sine/cosine of recorded hour, sine/cosine of weekday (Monday zero), and whether the just-observed adjacent gap exceeds its nominal interval. RMS spans are 24/168 FX records and 6/26 Treasury records; nominal gaps are 3600 and 604800 seconds. Calendar coordinates follow the stored source clock; constant training columns are centered exactly at their repeated value.

$v_L(t)=[L^{-1}\sum_{j=t-L+1}^{t}(N_j-N_{j-1})^2]^{1/2}; \quad z_{i,\kappa,j}=(x_{ij}-\mu_{\kappa j})/s_{\kappa j}.$

Training-only centering and population standard deviations define z. An exactly constant coordinate receives unit scale after centering. Matching completed response origins retains the different past-only lookbacks required by the feature construction. The response scale a is computed from completed training headrooms and magnitudes:

$a_\kappa=\max\{\text{mean}_{i\in T}(G_{i,H}),10^{-6}\,\text{mean}_{i\in T}|N_i|,10^{-12}\}.$

QR minimizes the normalized headroom regression-quantile objective with an unpenalized intercept and a ridge penalty on slopes. The fixed menu is λ in $\{0.001,0.01,0.1\}$; $\tau=19/20$. Training optimizes the unprojected response, while validation scores the final projected report in native units.

$\min_{\beta_0,\beta} W^{-1}\sum_{i\in T}\rho_\tau(G_{i,H}/a_\kappa-\beta_0-z_{i,\kappa}^T\beta)+(\lambda/2)\|\beta\|_2^2.$

$C_t^{QR}=B_t+\max\{0,a_\kappa(\beta_{0,\kappa}+z_{t,\kappa}^T\beta_\kappa)\}.$

The intercept is unpenalized and slopes receive a ridge penalty. Adding B to a headroom quantile translates it into a peak quantile; projecting a report up to the known response floor weakly improves realized pinball loss. The equivalent peak parameterization and numerical conventions are retained in Appendix C.

KNN uses training-standardized features with the frozen group weights and recency tie-breaking rule specified in Appendix C. It reports B plus a local headroom quantile. Scaled KNN divides each selected headroom by its past short RMS and multiplies the resulting quantile by the query RMS, using the same fixed stabilizer within each refit block.

$s_{i|\kappa}=\max\{v_{\text{short}}(i),10^{-6}a_\kappa\}; \quad C_t^{\text{scaled}}=B_t+s_{t|\kappa}Q_k(\{G_{i,H}/s_{i|\kappa}:i\in N_{t,\kappa,k}\}).$

The neighbor menus are {180,360,720} at both FX windows, {20,40,52} at Treasury W52, and {40,80,160} at W260. The implementation uses exact order statistics and fails on an unavailable rank. For k=52, $\lceil (k+1).95\rceil=51$, while the ordinary .95 empirical pinball quantile has rank 50. At W720, W2880 and W260, however, W times .95 is integer: the lower ordinary rank and the adjacent corrected rank are both endpoints of the empirical pinball minimizer interval. The W52 distinction is specific to its noninteger probability-times-pool-size case. A later endpoint/rank ablation is retained in Appendix D. Plain KNN at k=W equals frozen F exactly; membership is then the complete training pool. Scaled KNN still changes responses at k=W. Long-window menus restrict neighborhood sizes below the full pool. Appendix C records numerical acceptance and failures.

Post-result / exploratory classical maximal-inequality benchmark

Proposition 1 (classical Lévy bound, direct scalar application). Let the future real increments $X_1,\ldots,X_H$ be independent and individually symmetric about zero, with $S_0=0$ and $S_h=\sum_{j=1}^{h}X_j$. They need not have the same distribution or finite moments. For a fixed starting position x, write $B=|x|$ and $q=\inf\{u:\Pr(|S_H|\le u)\ge 0.975\}$. Lévy's maximal inequality implies the following strict-breach bound; Appendix J gives the proof. This is a classical result applied to this target, not a new theorem (Sussman, 2023, Proposition 3.4).

$\Pr(\max_{0\le h\le H}|x+S_h|>B+q)\le\Pr(\max_{1\le h\le H}|S_h|>q)\le 2\Pr(|S_H|>q)\le 0.05.$

The operational report L uses the same frozen completed-origin pool and refit times as F. Its endpoint sample is $|N_{i+H}-N_i|$ and its one-based order rank is $\lceil 39(W+1)/40 \rceil$. Add the resulting estimate to B_t. This empirical plug-in is distinct from the known population quantile in Proposition 1. Overlapping endpoint samples and dependent, changing series do not inherit the proposition's population coverage. Conditional coverage additionally requires the independence and symmetry premises and the correct endpoint quantile conditional on issue information. The bound supplies no pinball-loss ordering against F.

5. Evaluation design

The FX primary is H6 QR W2880 minus original rolling F W720; the Treasury primary is H6 QR W260 minus frozen F W260. These are the two declared source-specific comparisons. The FX H24 QR W2880 minus original rolling S W720 comparison is a legacy-reference sensitivity: its reference was chosen in the preceding, already inspected study.

Every selected conditional family is compared with frozen F, S and K at matching source, horizon and history. The complete menu also retains rolling-versus-frozen update controls. The declared source primaries remain fixed throughout every post-result analysis.

Forecasts are scored in native source units using the stipulated upper-quantile pinball preference. The paired loss difference is left minus right, so a negative difference favors the left procedure:

$e=P-C;\quad L_{.95}=.95(e)_+ +.05(-e)_+;\quad D=M^{-1}\sum_t[L(P_t,C_t^{left})-L(P_t,C_t^{right})].$

Mean positive overrun averages $(P-C)_+$ over all issue origins; mean unused allowance averages $(C-P)_+$. Breaches use strict P>C. Conditional event severity uses each forecast's own breach occasions. Annual groups and the globally anchored H+1 grid supply the original sensitivity views.

Table 2. Complete validation-selected parameters. QR entries are ridge penalties; KNN entries are neighborhood counts. W52 plain KNN selects the full pool at both horizons.

Source	H	W	QR λ	KNN k	Scaled KNN k
FX	6	720	0.01	180	360
FX	6	2880	0.01	180	180
FX	24	720	0.01	180	180
FX	24	2880	0.001	180	180
Treasury	6	52	0.1	52	52
Treasury	6	260	0.001	80	80
Treasury	24	52	0.001	52	40
Treasury	24	260	0.1	160	160

The original conditional-study design and selected parameters were publicly frozen before their respective executions (see the linked reproduction record). The FX evaluation interval had previously been inspected. All new inference and robustness analyses are labeled post-result / exploratory; the round2 additions have local execution bindings, rather than a new public preregistration.

Post-result inference

After exact trace reproduction, we compare paired loss differences using Diebold–Mariano (1995) statistics with Bartlett Newey–West (1987) covariance. The fixed bandwidth is $L=\max\{H,\lfloor 4(n/100)^{2/9}\rfloor\}$; doubled bandwidth supplies a sensitivity check. Harvey, Leybourne and Newbold (1997) supply the reported small-sample multiplier, here used as an approximation for pinball loss. Fixed finite estimation windows motivate the unconditional predictive-ability interpretation of Giacomini and White (2006), under mixing, adequate high-order moments and positive long-run variance. Shared training can extend dependence beyond the target horizon. These are working assumptions for this post-result analysis.

$DM=\bar{d}/\sqrt{(\Omega/n)};\quad \Omega=\hat{\gamma}_0+2\sum_{j=1}^{L}(1-j/(L+1))\hat{\gamma}_j;\quad DM_{HLN}=DM\sqrt{[(n+1-2H+H(H-1)/n)/n]}.$

Kupiec (1995) proportion-of-failures and Christoffersen (1998) independence/conditional-coverage diagnostics use the original first-origin-plus-j(H+1) grid. Conditional coverage here adds full-sample-grid POF to the transition-independence LR. A DQ-type full-sample HAC Wald adaptation of Engle and Manganelli (2004) uses the intercept, issue-known capacity and matured hit lags H+1 through H+4. Singular covariance or an unidentified transition row yields NA. The fixed 72-member common-history family receives Holm (1979) adjustment; exact identical losses use conservative p=1 solely for multiplicity bookkeeping. Appendix H and the output CSVs retain all statistics and statuses.

6. Results: FX

Table 3. Complete selected conditional menu with breach rates and same-history frozen-F contrasts. Loss is in source-specific USD millions; B1 statistics are post-result / exploratory.

H	W	Method	Loss	Breach rate	Δ frozen F	DM / HLN
6	720	QR	0.433	8.015%	-2.117%	t=-0.616; p=0.538; HLN p=0.538
6	720	KNN	0.436	4.815%	-1.394%	t=-0.719; p=0.472; HLN p=0.473
6	720	Scaled KNN	0.466	4.060%	+5.349%	t=1.245; p=0.213; HLN p=0.214
6	2880	QR	0.386	5.675%	-14.767%	t=-5.866; p=4.47e-09; HLN p=4.83e-09
6	2880	KNN	0.411	4.815%	-9.302%	t=-3.615; p=0.0003; HLN p=0.000306
6	2880	Scaled KNN	0.426	4.121%	-5.795%	t=-1.255; p=0.209; HLN p=0.21
24	720	QR	1.035	13.947%	+31.504%	t=3.223; p=0.00127; HLN p=0.00133
24	720	KNN	0.836	7.592%	+6.197%	t=1.527; p=0.127; HLN p=0.128
24	720	Scaled KNN	0.897	6.642%	+13.886%	t=1.541; p=0.123; HLN p=0.125
24	2880	QR	0.735	8.015%	-3.758%	t=-0.504; p=0.614; HLN p=0.615
24	2880	KNN	0.770	6.883%	+0.837%	t=0.185; p=0.854; HLN p=0.854
24	2880	Scaled KNN	0.848	5.962%	+11.085%	t=1.325; p=0.185; HLN p=0.187

The H6 primary QR W2880 loss is USD 385,844.89, against USD 441,774.04 for original rolling F W720. The direct paired mean is −USD 55,929.15, or −12.660%. Against W2880 frozen F it is −14.767%; against frozen S at that same window it is also favorable. Thus the added conditional benchmark changes the older FX ranking. The cross-window primary changes history length and update frequency as well as the estimator; the matched-pool controls address these procedural differences while retaining estimator and feature distinctions.

The H6 score gain comes with more breach occasions. QR has 376 strict breaches (5.675%) versus 332 (5.011%) for old rolling F. Mean positive overrun falls from USD 166,531.06 to 156,123.03, while mean unused allowance falls by USD 920,830.41. More occasions breach, with a lower unconditional overrun contribution. Conditional severity averages different method-specific event sets; the statistic therefore compares different incident compositions. At W720, H6 QR improves on frozen F by only 2.117%, while scaled KNN is worse by 5.349%.

At H24, QR W2880 loss is USD 734,695.07 against USD 748,383.54 for old rolling S W720: −USD 13,688.47 (−1.829%). Breaches increase from 287 (4.332%) to 531 (8.015%); mean positive overrun rises from USD 149,813.88 to 317,678.12, while unused allowance falls. With W720, QR instead loses to frozen F by 31.504% and frozen S by 37.095%. In the inspected FX record, the H24 regression ranking changes with the history window.

The H6 primary loss difference is negative in both issue-year groups: −12.012% over 6,262 origins in 2016 and −29.513% over the 363-origin partial 2017. H24 improvement against its legacy reference is only −0.555% in 2016 and −33.007% in partial 2017; about 70.83% of the small total net gain comes from that short tail segment. The fixed H+1 grids retain favorable H6/H24 differences of −16.388%/−8.027%, with 947/265 origins. These subdivisions and grids describe sensitivity within the inspected FX record; their interpretation follows Section 9.

Figure 1 compares every selected conditional procedure with its matched frozen-F control; Table 3 adds breach rates and the corresponding same-history changes. Figure 3 displays every original annual primary comparison.




Figure 1. All 24 selected conditional forecasts versus frozen F with the same source, horizon, history window and refit schedule. Entries show within-reference mean-loss percentages; negative favors the conditional procedure. Source panels have explicit scales and are not pooled. Full-pool KNN identities are marked.

7. Results: Treasury

Table 4. Complete selected conditional menu with breach rates and same-history frozen-F contrasts. Loss is in source-specific USD millions; B1 statistics are post-result / exploratory.

H	W	Method	Loss	Breach rate	Δ frozen F	DM / HLN
6	52	QR	5095.519	30.325%	+83.354%	t=3.057; p=0.00224; HLN p=0.00298
6	52	KNN	2779.054	3.971%	+0.000%	IDENTICAL_LOSSES
6	52	Scaled KNN	4208.293	3.249%	+51.429%	t=3.682; p=0.000231; HLN p=0.000365
6	260	QR	4159.673	19.495%	+50.786%	t=2.669; p=0.00761; HLN p=0.00939
6	260	KNN	2663.561	10.108%	-3.447%	t=-0.442; p=0.658; HLN p=0.665
6	260	Scaled KNN	3127.173	8.303%	+13.358%	t=0.945; p=0.345; HLN p=0.355
24	52	QR	12165.574	47.653%	+227.411%	t=2.238; p=0.0252; HLN p=0.0415
24	52	KNN	3715.693	11.913%	+0.000%	IDENTICAL_LOSSES
24	52	Scaled KNN	5625.667	11.552%	+51.403%	t=2.797; p=0.00516; HLN p=0.011
24	260	QR	5204.760	30.325%	+109.843%	t=1.667; p=0.0955; HLN p=0.128
24	260	KNN	2257.963	4.693%	-8.964%	t=-1.941; p=0.0523; HLN p=0.0768
24	260	Scaled KNN	4137.264	2.888%	+66.805%	t=4.923; p=8.53e-07; HLN p=9.8e-06

The declared H6 QR W260 primary has mean loss 4,159.673 versus 2,758.665 for matched frozen F, in USD millions. Its paired increase is 1,401.008 (+50.786%), with 54 rather than 28 strict breaches (19.495% versus 10.108%). Mean positive overrun rises from 1,395.639 to 3,131.862 and unused allowance falls from 28,656.166 to 23,688.088. Under the stipulated score, the more heavily weighted overrun increase dominates the reduced unused quantity. Numerically accepted fits produce this adverse accuracy result.

Every Treasury QR cell loses to its own frozen F control: +83.354% and +50.786% at H6 for W52/W260, and +227.411% and +109.843% at H24. Short-history H24 QR breaches 132 of 277 responses. All four selected scaled-KNN cells also have higher loss than their matched F controls. The primary QR difference is adverse in each complete year

2021–2025. Its favorable −14.404% in 2026 covers only 16 issue origins through April, with zero breaches for either method. The full adverse primary and this partial-period result are both retained.

Plain KNN W52 selects k=52 at both horizons and equals frozen F at every stored issue. These rows use the complete pool, so they demonstrate the full-pool fallback. At W260, selected KNN has secondary all-origin loss reductions of 3.447% (H6) and 8.964% (H24) against F. The adverse QR primary remains the declared test case. Each selected conditional procedure loses to at least one retained historical reference at its horizon. The menu is reported as a complete descriptive comparison.

The secondary H24 KNN W260 comparison has 13 versus 12 breaches and mean overrun 327.816 versus 306.848, alongside reduced unused allowance. On the fixed 12-origin H+1 grid its mean loss is 3,111.788 against 2,807.400 for F, reversing to +10.842%. The H6 KNN sensitivity remains favorable: its 40-origin difference remains −22.071%. These exact grids remove shared recorded label observations but retain common training histories and possible time dependence. The sparse reversal documents instability in the secondary Treasury gain.

Table 4 retains every conditional family and history. Figure 2 juxtaposes the declared primaries and the FX legacy sensitivity, keeping score changes, breach-rate changes and severity separate.

Declared comparisons: score and tail tradeoffs

	FX H6 primary	FX H24 legacy	Treasury H6 primary
Mean pinball loss change (%)	-12.660	-1.829	+50.786
Breach-rate change (pp)	+0.664	+3.683	+9.386
Mean overrun difference	-0.010	+0.168	+1736.223
Mean unused difference	-0.921	-3.463	-4968.078

Overrun/unused differences: source-specific USD millions; source amounts stay separate.

Figure 2. The two declared source primaries and FX H24 legacy-reference sensitivity: relative mean-loss, overrun and unused-allowance changes, plus breach-rate differences in percentage points. These are separate descriptive quantities, separate measures of forecast behavior. A reduced unused quantity is not automatically overall improvement.

Every annual primary and legacy comparison

FX H6 primary		FX H24 legacy		Treasury H6 primary	
2016 (n=6262)	-12.01%	2016 (n=6262)	-0.56%	2021 (n=52)	+19.11%
2017 (n=363)	-29.51%	2017 (n=363)	-33.01%	2022 (n=52)	+44.28%
				2023 (n=52)	+118.29%
				2024 (n=52)	+48.18%
				2025 (n=53)	+12.00%
				2026 (n=16)	-14.40%

All years retained; the final issue-year groups are partial. These are within-record sensitivities.

Figure 3. Every issue-year group for the two declared primaries and FX H24 legacy sensitivity. Values are paired within-reference loss percentages, with origin counts and partial years labeled. They are subdivisions of the same source records, not independent replications, confidence intervals or retrospectively selected switching regimes.

8. Robustness

Post-result / exploratory robustness

AR has lower loss in five of eight cells against S, five against F and five against QR; these sets differ. Relative to QR it loses in three of four FX cells and improves in all four Treasury cells. The exploratory AR comparisons show a different observed loss pattern from the adverse Treasury QR primary. Against S, the FX H6 changes are −1.536%/−0.546% at W720/W2880, while H24 changes are +2.386%/+5.147%. Treasury changes are −13.444%/−18.887% at H6 and +35.125%/−24.118% at H24 for W52/W260. The complete table retains the original model identities. Against S, AR mean report and mean unused allowance decrease in every cell, while strict breach occasions increase in all eight and mean positive overrun increases in seven. Treasury H24 W260 loss falls from 2478.871 to 1881.016 USD millions (−24.118%), yet breaches rise from 12 to 59 of 277 and mean positive overrun rises from 306.848 to 892.437. Mean unused allowance falls from 43747.307 to 20664.031. Under the stipulated score, .05 times the unused reduction exceeds .95 times the added overrun. This explains the observed loss change without asserting safer limits, calibrated tails or observed financing savings.

A later post-inspection rank/endpoint ablation adds a same-pool, same-update intercept-only empirical headroom quantile, choosing its lower minimizer endpoint. The matched FX H6 W2880 QR comparison still lowers loss by 14.773%, with 376 rather than 365 breaches; the matched Treasury H6 W260 QR comparison still raises loss by 46.058%, with 54 rather than 32 breaches. All four Treasury QR cells remain adverse against this added report. The lower-rank rule improves on original frozen F in three cells and loses in five, so the empirical minimizer endpoint can perform worse in evaluation. The two headline signs persist under this specific control. Table D3 retains every cell and the later design history.

The two FX H6 comparisons reduce both mean positive overrun and mean unused allowance. Their score improvement holds across the linear preference range, while strict breach occasions still increase. By contrast, the FX H24 legacy comparison increases overrun while reducing unused allowance: its 1.829% pinball improvement reverses at $\omega=20.630898$ ($a=0.953770$), close to 19. The Treasury H6 primary becomes preferable only below $\omega=2.861429$ ($a=0.741028$); its adverse result at 19 remains intact. These crossings describe the fixed sample means. For each of the eight source/H/W cells, the complete lower envelope retains QR, KNN, scaled KNN, matched frozen F/S/K, AR and I. The eight reports share the completed-label history length and update schedule within a cell. Rolling and one-state reports are excluded from these frontiers; the original cross-window primary references remain separately labeled in Table F4. Appendix E retains every full and sparse-grid frontier, endpoint and tie convention. A supplementary figure displays the whole preference range and a high-shortfall-weight detail. The full-grid Treasury H6 W260 QR is never optimal within these eight fixed reports, including both endpoints. At $a=.95$ the preferred report changes on four of eight sparse grids: FX H24 W720 and Treasury H6 W52/W260 and H24 W52. Treasury H24 grids have only 12 origins. These changes delimit sample stability under the original sparse-grid construction.

Post-result diagnostics clarify the issued reports rather than changing their rankings. Every common realized target satisfies $P(H6)\leq P(H24)$, yet the separately fitted reports can cross: FX W720 QR crosses on 640/6,625 issues, and Treasury W52 QR on 91/277. Some short Treasury historical F/S controls also cross because the horizons use different mature episode sets. Appendix D retains all 45 original pairs and a separately frozen addition covering all four AR pairs. AR has zero observed FX crossings, but Treasury W52/W260 cross on 59/26 of 277 issues. The horizon-specific fits and mature scenario sets differ, producing the observed incoherence. Figure 3 retains the complete annual comparison; all detailed coherence and fixed-terminal diagnostics remain in Appendix D.

Table 5. Summary of post-result / exploratory robustness; complete numerical tables remain in the appendices.

Analysis	Main finding	Retained adverse evidence
AR replay	Improves on QR in every Treasury cell	Breach occasions rise against S in every cell
Empirical rank	Both matched H6 QR directions persist	Lower empirical rank loses to F in five cells
Preference frontier	FX H6 reduces both loss components	FX H24 reverses near the original weight
Horizon coherence	Targets are ordered by construction	Separately fitted reports cross

Post-result predictive accuracy and calibration

Table 6. Post-result / exploratory paired predictive-accuracy tests. Negative t favors QR; normal DM and Student-reference HLN p values are two sided. Holm p uses the fixed 72 common-history family; original cross-window FX comparisons are outside it. Original primaries and legacy remain fixed.

Comparison	L	DM t	DM p	HLN t	HLN p	Holm p
FX H6 primary	10	-5.18	2.2e-07	-5.18	2.31e-07	Outside family
FX H6 matched	10	-5.87	4.47e-09	-5.86	4.83e-09	2.86e-07
FX H24 legacy	24	-0.204	0.839	-0.203	0.839	Outside family
Treasury H6 primary	6	2.67	0.00761	2.62	0.00939	0.32

Table 7. Post-result / exploratory coverage diagnostics. k/n counts strict breaches on the fixed H+1 grid; POF/independence/CC are chi-square-reference p values. DQ is the overlap-aware full-sample HAC adaptation. Boundary and small-sample qualifications apply.

Report	k/n	POF p	Ind p	CC p	DQ p
FX QR H6	50/947	0.695	0.388	0.638	0.736
FX rolling F H6	49/947	0.807	0.712	0.907	0.00798
FX matched F H6	49/947	0.807	0.368	0.647	0.0407
FX QR H24	21/265	0.0433	0.0866	0.0299	0.101
FX legacy S H24	10/265	0.339	0.0454	0.0855	0.0299
Treasury QR H6	7/40	0.00413	0.775	0.0157	0.00891
Treasury F H6	4/40	0.199	0.338	0.277	8.52e-05

The original FX H6 primary and matched-history comparison have negative DM statistics, while the Treasury primary has a positive statistic (Table 6). The FX H24 legacy has DM p=0.839 and HLN p=0.839, so its 1.83% descriptive improvement is not statistically significant at the 5% working-reference level. At the declared bandwidth, 19 of 72 Holm-adjusted working-reference p values fall below 5%, compared with 18 at doubled bandwidth. The declared-bandwidth count includes 15 favorable and 4 adverse mean differences for the conditional method. The matched FX H6 comparison has Holm p=2.86e-07; the Treasury primary has Holm p=0.32 and does not meet that family-adjusted threshold. Its descriptive loss deterioration remains an adverse finding. On the Treasury primary grid, 7/40 strict breaches yield POF p=0.00413 and CC p=0.0157. These coverage diagnostics assess different nulls from paired equal predictive loss.

Figure 4 fixes τ=0.95 and varies the elementary-score threshold θ (Ehm et al., 2016). Both primary diagrams cross: QR has favorable and unfavorable threshold regions. Consequently neither report empirically dominates the other over the entire class of consistent 0.95-quantile scores. The check evaluates every unique forecast/outcome breakpoint privately; the displayed 201-point grid is an aggregate visualization. This differs from the archived preference frontier, which holds reports fixed and changes shortfall/excess weights. The calibration-and-sharpness perspective of Gneiting, Balabdaoui and Raftery (2007) motivates separate assessment of forecast concentration and calibration; the scalar capacity report addresses its stated functional.




Figure 4a. FX original primary elementary quantile scores at fixed τ=0.95. Lower is better. The full-breakpoint check finds crossing; the display grid alone is not used to establish that finding.




Figure 4b. Treasury original primary elementary quantile scores at fixed τ=0.95. Lower is better. The full-breakpoint check finds crossing; the display grid alone is not used to establish that finding.

Table 8. Post-result / exploratory parsimonious QR. Four predictors plus intercept, expanded validation penalty menu, original split/refit schedule. Percent changes compare loss against the original QR and matched F. Every original primary remains unchanged.

Source	H	W	λ	Δ QR %	Δ F %	Breaches	Status
Treasury	6	52	10	-46.9	-2.59	15	COMPLETE
Treasury	6	260	10	-32.2	2.17	32	COMPLETE
Treasury	24	52	10	-67.6	5.95	36	COMPLETE
Treasury	24	260	10	-52.3	0.11	15	COMPLETE
FX	6	720	0.01	3.67	1.48	414	COMPLETE
FX	6	2880	0.01	12.3	-4.29	399	COMPLETE
FX	24	720	0.01	-11.4	16.5	701	COMPLETE
FX	24	2880	0.01	0.898	-2.89	508	COMPLETE

Post-result / exploratory Lévy plug-in benchmark

Table 9. All eight classical-bound plug-in L cells. Loss uses source-native units (FX USD; Treasury USD millions). Δ F/QR compares the same W and refit schedule; positive values are adverse for L. DM p is the unadjusted working-reference paired L–F test; Appendix J retains both comparators and a separate 16-member Holm family. Original primaries are unchanged.

Source	H	W	Mean loss	Hits / n	Δ F %	Δ QR %	DM p vs F
FX	6	720	5.795e+05	101/6625	30.93	33.76	2.988e-15
FX	6	2880	5.903e+05	75/6625	30.39	52.98	6.695e-15
FX	24	720	8.728e+05	136/6625	10.85	-15.71	0.06462
FX	24	2880	8.24e+05	174/6625	7.941	12.16	0.03958
Treasury	6	52	2873	10/277	3.37	-43.62	0.3317
Treasury	6	260	2434	15/277	-11.77	-41.49	0.1785
Treasury	24	52	4694	42/277	26.32	-61.42	0.1167
Treasury	24	260	2725	22/277	9.868	-47.64	0.2746

The plug-in reduces FX breach counts relative to F in every cell while increasing pinball loss. In Treasury H24 it has both higher loss and more breach occasions than F; its Treasury H6/W260 loss is lower than F. Thus the empirical report does not have a uniform conservative or accuracy ordering. Corrected level-AR sensitivities give observed peak-breach rates of 0–0.913%. Negative-increment AR reaches 5.032%, an excess of 0.032 percentage points with Monte Carlo SE 0.06913 percentage points. These oracle-threshold mechanisms supply no empirical-coverage guarantee (Appendix J).

Post-result / exploratory published-aggregate panel

A metadata/calendar rule selected 12 additional NYFed net-position series before model evaluation: five nominal-coupon maturity buckets, four TIPS buckets, agency/GSE debt, MBS and corporate totals. The two longer-maturity coupon keys lack the required complete history and are retained as skips. Every included series uses the original 706-date economic grid and the same validation/evaluation/refit indices. The QR33 and C2 structures retain their penalty menus; a common strict complete-menu policy and larger-penalty exact tie rule apply to this new panel. All 192 procedure-cell outputs complete. These series share dealers, reporting regimes and related balance-sheet components; they are additional aggregate tasks rather than independent institutions.

Table 10. Complete panel summary: number of four H/W cells with lower pinball loss than matched F. These are descriptive counts, not independent trials or evidence selected after screening losses. Appendix K retains every loss change and breach count.

Series key	QR33 lower / cells	C2 lower / cells	L lower / cells
PDPOSMBS-TOT	0/4	0/4	2/4
PDPOSCS-TOT	0/4	3/4	1/4
PDPOSFGS-TOT	1/4	1/4	3/4
PDPOSTIPS-L2	0/4	0/4	4/4
PDPOSTIPS-G11	0/4	0/4	1/4
PDPOSTIPS-G2	0/4	0/4	0/4
PDPOSTIPS-G6L11	2/4	4/4	0/4
PDPOSGSC-L2	0/4	2/4	2/4
PDPOSGSC-G2L3	0/4	0/4	2/4
PDPOSGSC-G3L6	0/4	0/4	2/4
PDPOSGSC-G6L7	0/4	0/4	1/4
PDPOSGSC-G7L11	0/4	1/4	1/4

QR33 lowers loss against F in 3 of 48 cells; C2 does so in 11 and L in 19. The remaining adverse cells are retained, including QR33 loss changes as high as 544.3% and C2 changes as high as 299.1%. This extension does not support a general conditional-model advantage, and simplification alone does not resolve the weekly-task failures.

Post-result / exploratory mature conditional baselines

Table 11. Fixed FHS, GARCH-scaled peak-headroom POT and true leaf-distribution QRF on all original cells. Each entry gives Δ loss% versus matched frozen F / strict breaches; negative Δ favors the new method. NA retains a failed or structurally ineligible complete-cell comparison; issued counts are not scored as partial successes. Full native losses, reasons and comparisons are in Appendix L.

Source	H	W	FHS: Δ% / hits	Peak POT: Δ% / hits	QRF: Δ% / hits
FX	6	720	NA (240 issued)	NA (240 issued)	5.793 / 402
FX	6	2880	-1.258 / 278	3.269 / 363	-4.48 / 354
FX	24	720	NA (240 issued)	NA (240 issued)	16.02 / 779
FX	24	2880	6.255 / 136	10.61 / 369	7.335 / 600
Treasury	6	52	-1.883 / 7	NA (0 issued)	-6.83 / 16
Treasury	6	260	-8.276 / 4	-10.35 / 17	6.198 / 33
Treasury	24	52	60.26 / 0	NA (0 issued)	4.279 / 44
Treasury	24	260	125.1 / 0	32.09 / 20	33.94 / 62

These baselines use the original issue origins, mature-label windows and scheduled refits. FHS recursively simulates signed position paths from empirical GARCH-standardized innovations (Barone-Adesi et al., 1999). Peak POT fits GPD exceedances of GARCH-scaled mature peak headrooms; it is an adaptation of conditional EVT, not a literal McNeil–Frey (2000) return-innovation implementation. QRF uses the training-response distribution in matching tree leaves (Meinshausen, 2006), rather than quantiles of tree means. The configurations are fixed resource-bounded choices. Complete failures and adverse loss or breach changes remain visible.

Against matched F, the complete-cell counts are FHS: 3 lower-loss and 3 higher-loss cells out of 6 complete comparisons; PEAK_POT: 1 lower-loss and 3 higher-loss cells out of 4 complete comparisons; QRF: 2 lower-loss and 6 higher-loss cells out of 8 complete comparisons. These are descriptive counts across related histories and targets, not independent replications. The failed cells limit method coverage; they do not establish predictive inferiority.

9. Discussion and limitations

Feature transport and the Treasury failure

The executed QR specifications perform poorly on the weekly Treasury task. Their 25 level coordinates span the current observation through 24 weeks of history. The two hour coordinates, two weekday coordinates and gap indicator are exactly constant on this weekly source. Training-only centering makes those five columns exactly zero, so they provide no predictive variation and are not a demonstrated mechanism of overfitting. The original nominal design has 33 predictors plus an intercept, or 34 coefficients, with at most 28 varying columns and 29 potentially predictive coefficients before other collinearity, against W=52 overlapping training labels in the shorter history. All four Treasury penalty selections are menu endpoints. Its 52 validation origins contain at most 3 disjoint H24 label paths, a path count rather than an effective-sample-size estimate. Correlated lag levels, overlapping labels, restricted tuning, target dynamics and reporting composition remain competing explanations. Neither the parsimonious sensitivity nor the aggregate panel separates those mechanisms.

The four-predictor sensitivity uses B, signed N and short/long increment RMS, with validation penalties {0.001,0.01,0.1,1,10}. It lowers loss against the original QR in 5 of 8 cells and against matched F in 3 cells. In the Treasury primary cell its loss change is -32.2% relative to the original QR and 2.17% relative to F. All adverse cells remain in Table 8. The exercise changes features and the penalty menu jointly, so it measures a parsimonious specification sensitivity rather than identifying a feature-only cause or rehabilitating the original primary.

The contrast between the sources concerns a separately fitted procedure family: each source receives its own transforms, coefficient estimates, temporal scales and validation selection. The Treasury deterioration is an accuracy failure of the executed QR specifications. The exploratory AR comparison shows that the observed Treasury loss ranking differs across the executed families.

Limitations

The FX evaluation interval was previously inspected, and the subsequent robustness analyses are post-result / exploratory. These records do not restore a compatible unseen dealer sample. Treasury is a downloaded-vintage, economic-date aggregate replay rather than an original-release public-information backtest; cross-dealer netting, valuation and composition distinguish it from an individual dealer's book.

Shared histories, overlapping labels, finite validation and investigator selection limit statistical generalization. The known floor can create an outcome atom: a valid upper quantile need not have exactly the nominal strict-breach

probability. Numerical acceptance supplies no conditional-coverage guarantee. The horizon-specific mature scenario sets differ, so a zero observed crossing count cannot establish general coherence.

Stipulated shortfall/excess weights are not observed financing prices. Quotes, fills, hedges, book decomposition, collateral and cash flows would be required to map this passive envelope into an operational inventory, margin or funding policy. This study evaluates forecasts, not such interventions; Avellaneda and Stoikov (2008) provide a distinct active inventory-sensitive quoting framework.

Further alternatives include CAViaR, sequential conformal procedures, richer mean or volatility dynamics, and canonical innovation-tail conditional EVT. The fixed FHS, QRF and peak-POT additions in Appendix L extend the executed menu, with their failures and resource bounds retained. Same mature-label pools and update schedules improve procedural comparability while leaving different feature lookbacks, model assumptions and estimation rules. Compatible new dealer observations and decision-linked inputs remain necessary for broader portability.

A post-result / exploratory selected-fit readback examines the Treasury C2 menu boundary directly (Appendix I). At $\lambda=10$, standardized response-normalized slope norms range from 0.001497 to 0.01937. Removing slopes while keeping each fitted intercept changes capacity by a mean absolute 136.8–832.3 source USD millions across the four cells. Relative to the lower empirical headroom-quantile report I, observed loss is 0.2432–2.192% lower; this descriptive difference has no dedicated selection-adjusted test. The fits have small normalized slopes and high headroom correlation with I, but are not identical to I or F. A boundary penalty and high capacity correlation do not by themselves demonstrate zero feature value. Three of the four C2 cells still have higher loss than F.

The expanded aggregate panel strengthens the negative transport finding but does not isolate its mechanism. Features and tuning menus, finite sample size, target behavior and reporting-regime changes can all matter. Downloaded cross-era histories are not original-release vintages; metadata/provenance for each key are retained. A prospective public-release protocol freezes the four procedures and selected historical-validation penalties before future issue artifacts. The first planned new release is after the protocol freeze, rather than retroactively labeling the already published October 14 observation as unseen. Because the release clock lags economic dates, the target includes prediction of unpublished measurements, with a nowcast/forecast mixture. A dated local amendment fixes future path values at their first actual capture and logs later revisions separately; operational capture and missing-week handling still require a bound adapter (Appendix K). Results await prior timestamped report issuance and target publication; no live prospective validation is claimed.

10. Conclusion

Mature conditional estimators change the historical ranking on the sampled FX peak task, while the Treasury regression primary deteriorates. Matched-history FX H6 QR lowers mean loss by 14.77% and raises breaches from 360 to 376; the Treasury primary raises loss by 50.79% and breaches from 28 to 54. Short-history QR deterioration and the sparse-grid Treasury neighbor reversal remain central adverse findings. The exploratory AR, rank, preference and coherence comparisons show why score improvements, breach frequency and report shape should be evaluated together. The study contributes a controlled application and model-validation benchmark for measured position peaks. Its next evidential step is compatible new financial observations and an explicit decision mapping.

The exploratory 12-series extension also finds higher loss than F in 45/48 QR33 cells and 37/48 parsimonious cells, further limiting general performance claims.

Fixed FHS, peak-POT and leaf-distribution QRF add adverse outcomes and explicit numerical exclusions rather than a universal accuracy ranking. Compatible unseen-dealer evidence remains an open requirement.

References

Arjani, N., F. Li and L. Sabetti (2020). Monitoring Intraday Liquidity Risks in a Real Time Gross Settlement System. Payments Canada discussion paper, 23 April. official report.

Avellaneda, M. and S. Stoikov (2008). High-frequency trading in a limit order book. *Quantitative Finance* 8(3), 217–224. DOI:10.1080/14697680701381228.

Barone-Adesi, G., K. Giannopoulos and L. Vosper (1999). VaR without correlations for portfolios of derivative securities. *Journal of Futures Markets* 19(5), 583–602. DOI:10.1002/(SICI)1096-9934(199908)19:5<583::AID-FUT5>3.0.CO;2-S.

Biau, G. and B. Patra (2011). Sequential Quantile Prediction of Time Series. *IEEE Transactions on Information Theory* 57(3), 1664–1674. DOI:10.1109/TIT.2011.2104610.

Board of Governors of the Federal Reserve System. FR 2004: Government Securities Dealer Reports. Official reporting framework, 22 October 2026.

Boudoukh, J., M. Richardson, R. Stanton and R.F. Whitelaw (2004). MaxVaR: Long-Horizon Value at Risk in a Mark-to-Market Environment. *Journal of Investment Management* 2(3), 14–19. author manuscript.

Boyd, S., N. Parikh, E. Chu, B. Peleato and J. Eckstein (2011). Distributed Optimization and Statistical Learning via the Alternating Direction Method of Multipliers. *Foundations and Trends in Machine Learning* 3(1), 1–122. DOI:10.1561/2200000016. author PDF.

Casgrain, P., M. Larsson and J. Ziegel (2023). Anytime-valid sequential testing for elicitable functionals via supermartingales. arXiv:2204.05680v2, 4 June.

Christoffersen, P.F. (1998). Evaluating Interval Forecasts. *International Economic Review* 39(4), 841–862. DOI.

davidlindsay1979 (n.d.). *Net Position Time Series Dataset* (Version 2) [Data set]. Kaggle. Dataset record.

Diebold, F.X. and R.S. Mariano (1995). Comparing Predictive Accuracy. *Journal of Business & Economic Statistics* 13(3), 253–263. DOI.

Duffie, D., M. Fleming, F. Keane, C. Nelson, O. Shachar and P. Van Tassel (2023). Dealer Capacity and U.S. Treasury Market Functionality. Federal Reserve Bank of New York Staff Report No. 1070, August, revised October. DOI:10.59576/sr.1070. official report.

Ehm, W., T. Gneiting, A. Jordan and F. Krüger (2016). Of quantiles and expectiles: consistent scoring functions, Choquet representations and forecast rankings. *Journal of the Royal Statistical Society, Series B* 78(3), 505–562. DOI.

Engle, R.F. and S. Manganelli (2004). CAViaR: Conditional Autoregressive Value at Risk by Regression Quantiles. *Journal of Business & Economic Statistics* 22(4), 367–381. DOI:10.1198/073500104000000370.

Federal Reserve Bank of New York (2026). Primary Dealer Statistics, series PDPOSGS-B; downloaded snapshot checked 22 October 2026. Definitions and publication information; source route. © 2026 Federal Reserve Bank of New York. Content from the New York Fed subject to the Terms of Use at newyorkfed.org.

Giacomini, R. and H. White (2006). Tests of Conditional Predictive Ability. *Econometrica* 74(6), 1545–1578. DOI.

Gneiting, T. (2011). Making and Evaluating Point Forecasts. *Journal of the American Statistical Association* 106(494), 746–762. DOI:10.1198/jasa.2011.r10138.

Gneiting, T., F. Balabdaoui and A.E. Raftery (2007). Probabilistic forecasts, calibration and sharpness. *Journal of the Royal Statistical Society, Series B* 69(2), 243–268. DOI.

Hansch, O., N.Y. Naik and S. Viswanathan (1998). Do Inventories Matter in Dealership Markets? Evidence from the London Stock Exchange. *Journal of Finance* 53(5), 1623–1656. DOI: 10.1111/0022-1082.00067.

Harvey, D., S. Leybourne and P. Newbold (1997). Testing the equality of prediction mean squared errors. *International Journal of Forecasting* 13(2), 281–291. DOI.

Henzi, A. and J.F. Ziegel (2022). Valid sequential inference on probability forecast performance. arXiv:2103.08402v3, 1 July.

Ho, T. and H.R. Stoll (1981). Optimal dealer pricing under transactions and return uncertainty. *Journal of Financial Economics* 9(1), 47–73. DOI: 10.1016/0304-405X(81)90020-9.

Holm, S. (1979). A Simple Sequentially Rejective Multiple Test Procedure. *Scandinavian Journal of Statistics* 6(2), 65–70. Publisher record.

Hull, J. and A. White (1998). Incorporating volatility updating into the historical simulation method for value-at-risk. *Journal of Risk* 1(1), 5–19. DOI:10.21314/jor.1998.001.

Koenker, R. and G. Bassett, Jr. (1978). Regression Quantiles. *Econometrica* 46(1), 33–50. DOI:10.2307/1913643.

Kupiec, P.H. (1995). Techniques for Verifying the Accuracy of Risk Measurement Models. *Journal of Derivatives* 3(2), 73–84. DOI.

Liao, J. and T.V. Theodosopoulos (2005). Study on optimal timing of mark-to-market for contingent credit risk control. arXiv:math/0506077v1, 4 June 2005.

Lyons, R.K. (1995). Tests of microstructural hypotheses in the foreign exchange market. *Journal of Financial Economics* 39(2–3), 321–351. DOI: 10.1016/0304-405X(95)00832-Y.

McNeil, A.J., and Frey, R. (2000). Estimation of tail-related risk measures for heteroscedastic financial time series: an extreme value approach. *Journal of Empirical Finance*7(3–4),271–300. DOI:10.1016/S0927-5398(00)00012-8.

Meinshausen, N. (2006). Quantile Regression Forests. *Journal of Machine Learning Research* 7, 983–999. Official article.

Newey, W.K. and K.D. West (1987). A Simple, Positive Semi-Definite, Heteroskedasticity and Autocorrelation Consistent Covariance Matrix. *Econometrica* 55(3), 703–708. Publisher record.

Petrunia, R., L. Sabetti and M. Voia (2019). Tail Risk in the Automated Clearing Settlement System (ACSS). Payments Canada Discussion Paper No. 7, January 2019 update of September 2018 original. official discussion paper.

Politis, D.N. and J.P. Romano (1994). The Stationary Bootstrap. *Journal of the American Statistical Association* 89(428), 1303–1313. DOI:10.1080/01621459.1994.10476870.

Sussman, E. (2023). A strengthened Orlicz–Pettis theorem via Itô–Nisio. *Annals of Functional Analysis* 14, article22, Proposition3.4. DOI:10.1007/s43034-022-00225-1.

Sözen, C. (2026). Monotone forward expected shortfall paths with joint conformal calibration. *AIMS Mathematics* 11(9), 27665–27688. Published 1 September 2026. DOI:10.3934/math.20261106.

van den Hoogen, S. (2017). Intraday liquidity risk estimation using transaction data: an extreme value theory approach. MSc Applied Mathematics thesis, Delft University of Technology, August. Institutional record.

Wisniewski, W., D. Lindsay and S. Lindsay (2020). Application of Conformal Prediction Interval Estimations to Market Makers' Net Positions. Proceedings of the Ninth Symposium on Conformal and Probabilistic Prediction and Applications, *Proceedings of Machine Learning Research* 128, 285–301. Official proceedings.

Wisniewski, W.J. (2024). Net Position and Retail Trader Behaviour: Implications for Market Makers' Risk Management Strategies. Doctoral thesis, Royal Holloway, University of London. Institutional original.

Xu, C. and Y. Xie (2023). Sequential Predictive Conformal Inference for Time Series. Proceedings of the 40th International Conference on Machine Learning, *Proceedings of Machine Learning Research* 202, 38707–38727. Official proceedings.

Zhou, Y., L. Lindemann and M. Sesia (2024). Conformalized Adaptive Forecasting of Heterogeneous Trajectories. arXiv:2402.09623v2, 15 May 2024.

Appendix A. Finite geometry and its predictive limits

On a common set and fixed rank, let $d_{abs}=\max_{i\in J}|B-|N_i||$ and $d_{signed}=\max_{i\in J}|x-N_i|$. Elementary order-statistic and projection stability gives $|F-K|\le d_{abs}$, $|S-K|\le d_{signed}$, hence $|F-S|\le d_{abs}+d_{signed}$. Appendix A proves these finite relationships and their predictive limits. Nearest signed-state membership minimizes its maximum signed distance at a given count; it need not minimize forecast loss. These statements do not automatically bound QR, scaled responses or the richer 33-coordinate distance.

A.1. Common-set projection bounds

These construction arguments concern F/S/K on an identical historical set and rank. They are retained from the earlier comparison. A one-dimensional signed-distance minimax property does not extend automatically to the multivariate neighbor distance, and the bounds do not automatically constrain QR or scale-adjusted responses.

These are elementary finite-array arguments, rather than new asymptotic or coverage results. Fix a nonempty common selected set J, an available rank $1\le k\le|J|$, query $x=N_t$, floor $B=|x|$ and completed historical windows. All identities in this appendix concern real arithmetic. Write $B_i=|N_i|$, $d_{i,h}=N_{i+h}-N_i$, with $d_{i,0}=0$, and define

Let $a_i=\min_h d_{i,h}$ and $b_i=\max_h d_{i,h}$, so $a_i\le 0\le b_i$. These are the lower and upper increments denoted l and u in Section 4. The auxiliary nonnegative envelope coordinates below use $u_i=b_i$ and $v_i=-a_i$.

$$A_i=B+P_i-B_i;\quad T_i=\max_{0\le h\le H}|x+d_{i,h}|;\quad C_B(z)=\max(B,z).$$

Both A_i and T_i are at least B. The projection C_B is nondecreasing and 1-Lipschitz. It commutes with a finite order statistic, as does a common translation. Hence $F_J=Q_k(A)$, $S_J=Q_k(T)$ and $K_J=Q_k(C_B(P))$. Since $C_B(A_i)=A_i$,

$$|A_i-C_B(P_i)|\le|A_i-P_i|=|B-B_i|.$$

The reverse triangle inequality bounds each path value's change under a starting-position translation by $|x-N_i|$. Taking a maximum gives $|T_i-P_i|\le|x-N_i|$. Applying the same floor projection preserves that bound because $T_i\ge B$. Finally, paired order statistics are nonexpansive in the maximum coordinate difference: if $|u_i-v_i|\le r$ for every i, then $u_i\le v_i+r$ and conversely; monotonicity and common translation give $|Q_k(u)-Q_k(v)|\le r$. These observations prove the two common-set K bounds stated above. The triangle inequality and $||N_i|-|x||\le|N_i-x|$ prove the stated F/S bound. Keeping the rank fixed while changing one side's historical rows does not by itself justify these common-set comparison bounds.

Among all size-m subsets, selecting the m smallest signed distances minimizes the largest selected distance: any alternative must include an element at least as far away as the mth ordered distance. This finite minimax fact, with the declared tie convention, concerns the signed radius. It does not minimize the absolute-anchor radius, the sum of the two radii, the actual report gap or a forecast loss. For every fixed realized y, pinball loss is $\max(\tau,1-\tau)$-Lipschitz in its report, so a report-gap bound also bounds the absolute loss difference. Such stability supplies no sign for the realized or expected loss difference.

A.2. Envelope plateaus and the role of signed anchors

Put $u_i=b_i\ge 0$ and $v_i=-a_i\ge 0$. The headroom of an increment envelope at any anchor z is

$g_i(z)=\max(z+u_i,-z+v_i)-|z|$; for $z\ge 0$ it equals $\max(u_i,v_i-2z)$, and for $z\le 0$ it equals $\max(u_i+2z,v_i)$.

F pools each envelope's headroom at its historical anchor, $g_i(N_i)$; S pools its headroom at the query, $g_i(x)$. The branches agree at zero and together make g_i 2-Lipschitz, with range lying in $[\min(u_i,v_i),\max(u_i,v_i)]$. Apply the distance and range bounds to each paired row, then use the same-rank order-statistic stability from A.1 to obtain

$$|F_J-S_J|\le\max_{i\in J}\min\{2|x-N_i|,|u_i-v_i|\}.$$

For example, $u_i=v_i$ in every row makes $F_J=S_J$ at arbitrary anchors. Nonnegative query/source anchors together with $u_i\ge v_i$ in every row also suffice; the nonpositive analogue uses $v_i\ge u_i$. These are sufficient conditions, not a characterization: pooled order statistics can coincide even when paired rows differ. The range bound also gives $F_J,S_J\le R_J$, where $R_J=B+Q_k\{\max(u_i,v_i):i\text{ in }J\}$. This is a theoretical same-set ordering; no R retrieval configuration is evaluated in the extension, and the ordering does not make R_J preferable under a given loss.

If every $B_i\le B$, then $A_i\ge C_B(P_i)$ and $F_J\ge K_J$. If every $B_i\ge B$, then $C_B(P_i)=P_i\ge A_i$ and $F_J\le K_J$. Equal absolute anchors force F=K but leave signs relevant to S. A coherent full-pool example uses $c>0$, H=1, n=720, $\tau=19/20$, t=721 and $N_s=(-1)^{s+1}c$ for $0\le s\le 721$. The pool uses completed labels at origins $J=\{1,\ldots,720\}$, with 360 anchors of each sign and current $N_t=c$. All native peaks equal c and headrooms equal zero, so F=K=c. Raw reanchoring gives 360 peaks c and 360 peaks 3c; rank 685 gives S=3c. Thus $d_{abs}=0$ while $d_{signed}=2c$ and $|S-K|=2c$. This artificial path respects maturity and an available corrected rank; it is not an empirical claim or a claim about which rows nearest retrieval selects.

A.3. The same observed history allows opposite loss rankings

Choose $d>0$, $H\geq 1$, $0<\tau<1$ and n large enough that $k=\lceil (n+1)\tau \rceil \leq n$. Set $t=H+n-1$ and $N_s=d(t-s)$ for $0\leq s\leq t$, using $J=\{0,...,n-1\}$. Every selected response ends by t. Each historical peak is its own positive starting value, so $G_i=0$ and $F=0$. Each increment path equals $-dh$; reanchoring at $N_t=0$ gives $S=Hd$. Two continuations of this identical observed history yield:

Table A1. Artificial coherent continuations. Symbols are constructed quantities, not source-record estimates.

Future $N(t+h)$, $1\leq h\leq H$	Actual peak	F loss	S loss
0	0	0	$(1-\tau)Hd$
$-dh$	Hd	τHd	0

Either report can therefore have strictly smaller realized loss under a compatible future. A report ordering also need not hold universally. Consider the artificial history $(0,-4,-1,2)$, with $H=1$, $n=3$, $\tau=3/4$, issue $t=3$ and $J=\{0,1,2\}$. The current anchor and floor are $x=B=2$; all three historical windows are completed. Table A2 shows the native and raw-reanchored paths.

Table A2. Completed $H=1$ windows from one artificial history, in arbitrary position units. P is the native peak, G the native headroom and T the peak after reanchoring at $x=2$. These values are not source-record estimates.

i	Historical pair	P	G	Reanchored pair	T
0	0, −4	4	4	2, −2	2
1	−4, −1	4	0	2, 5	5
2	−1, 2	2	1	2, 5	5

The available corrected rank is $\lceil (3+1)\cdot 3/4 \rceil=3$. The third order statistics of G, $T-B$ and P are 4, 3 and 4, respectively, so $F=6$, $S=5$ and $K=4$. The small pool and $\tau=3/4$ belong only to this illustration. It contains no observation after issue and gives no realized query loss. These constructions establish logical limits, not the conditional law of the recorded series or a population ranking. A deterministic construction can refute an unconditional mathematical assertion while supplying no empirical validation for either rule.

A.4. Raw translation versus fixed-coefficient residual replay

For interpretation, fix common coefficients c and φ in $N_{s+1}=c+\varphi N_s+\varepsilon_s$. Replay a historical window's ordered innovations from query x. The coefficients may be treated as fixed after a refit; this algebra does not assert that they are the true dynamics. Denote replay by M and raw translation by $\tilde{N}$. Subtracting the two recursions and using induction gives

$$M_{i,h}(x)=N_{i+h}+\varphi^h(x-N_i); \quad \tilde{N}_{i,h}^{tr}(x)=N_{i+h}+(x-N_i).$$

Taking the maximum absolute path value yields

$$|\max_h|\tilde{N}_{i,h}^{tr}(x)|-\max_h|M_{i,h}(x)||\leq \max_{0\leq h\leq H}|1-\varphi^h|\ |x-N_i|.$$

For $0\leq\varphi\leq 1$ the coefficient is $1-\varphi^H$; at $\varphi=1$ the paths coincide. The identity needs no innovation law, while treating past innovations as representative of future conditional innovations is a separate assumption. Section 8 fits a specified mean recursion and compares realized loss, but does not validate that conditional innovation assumption. The extrema suffice to evaluate the peak after raw translation, but generally not after the dynamic correction, which depends on path order. The same-rank reports inherit a deterministic bound with max over historical anchor distances, not a guaranteed sign of loss improvement. Conditional validity remains a separate question.

Appendix B. Earlier construction and retrieval evidence

This appendix preserves the complete loss menus of the 21 October historical studies. Monetary displays are source-reported FX USD millions. The original rules include headroom F, raw reanchoring S, a symmetric increment envelope R, native peak K (native-peak K), and the current-floor report Floor. One factor changes from the original H6/W720/τ=.95 specification in each sensitivity cell. All use the same 6,625 issue origins; different preferences are not pooled into an overall winner.

Table B1. Complete original mean loss grid. Δ is the stored paired mean F minus S; negative favors F within that setting.

Setting	F	S	R	K	Floor	Δ F−S
main	0.441774	0.451149	0.527544	0.925885	1.813620	-0.009375
h12	0.592518	0.604781	0.689604	1.058614	2.950594	-0.012263
h24	0.777069	0.748384	0.803106	1.197833	4.630827	+0.028685
n480	0.449843	0.460816	0.532515	0.885243	1.813620	-0.010973
n1440	0.443793	0.448275	0.526979	0.931562	1.813620	-0.004482
tau90	0.659643	0.659499	0.763481	1.477201	1.718166	+0.000144
tau99	0.154991	0.158998	0.175486	0.288834	1.889983	-0.004007

The earlier retrieval extension takes all 720 completed origins, the most recent 360/180, or the nearest signed-state 360/180 from that eligible pool, with recency ties. Corrected ranks for 720/360/180 are 685/343/172. Its H6 nearest 360 K-minus-recent 360 K primary lowers loss by 31.83% against a weak native-level baseline, while both original full 720 F and S remain lower. This prior-menu leadership statement does not extend to the new conditional models in Sections 6–7.

Table B2. Complete earlier retrieval-extension mean loss menu and same-set F-minus-S paired mean, in source-reported USD millions. Full 720 rows repeat original references. Differences are computed before display rounding.

H	Historical set	F loss	S loss	K loss	F−S mean
6	full720	0.441774	0.451149	0.925885	-0.009375
6	recent360	0.454089	0.467678	0.872516	-0.013589
6	nearest360	0.448664	0.448731	0.594811	-0.000067
6	recent180	0.485866	0.478940	0.785878	+0.006926
6	nearest180	0.468133	0.459592	0.525471	+0.008541
24	full720	0.777069	0.748384	1.197833	+0.028685
24	recent360	0.807496	0.824047	1.246630	-0.016551
24	nearest360	0.791192	0.770354	0.960696	+0.020838
24	recent180	0.858214	0.895718	1.284092	-0.037504
24	nearest180	0.847734	0.827305	0.963931	+0.020429

All 12 fixed nearest-minus-recent score contrasts were negative. Each matched S gain accompanied more breaches and greater mean positive overrun; conditional severities describe method-specific event sets. January 2016 was adverse for the primary K retrieval comparison at both horizons. The original primary F-minus-S mean was −USD 9,374.86 (−2.08%), with 332 versus 254 breaches and greater mean overrun; its H24 and τ=.90 rankings reversed. The latter τ=.90 difference was small and did not establish equivalence.

The original diagnostic groups, monthly outcomes and four stationary-bootstrap-style profiles remain in results/diagnostics.csv and stability.csv. The fixed resampling used 999 circular geometric-block replicates per mean length 120,240,720,1440, PCG64 SeedSequence([20261020,length]), and noninterpolated order statistics 25/975. The stationarity, moment and dependence premises behind Politis and Romano (1994) were not established here; those profiles are mechanism sensitivities, not population confidence intervals, and are not applied to the new conditional contrasts. The earlier extension's complete 30-row summary, 22 paired comparisons, 10 geometry rows and 26 monthly display rows remain under results/revision-20261021/. No prior adverse evidence is removed.

Appendix C. Numerical, rank and reproduction conventions

After training standardization, KNN multiplies each signed-level coordinate (lags 0–24) by 1/5. Each coordinate in the two-RMS pair, hour sine/cosine pair and weekday sine/cosine pair is multiplied by $1/\sqrt{2}$; the magnitude and observed-gap coordinates retain unit weight. Distance is squared Euclidean distance in these weighted coordinates. Exact distance ties prefer the more recent training origin. These are the frozen coordinate multipliers, not weights fitted to evaluation outcomes.

All issued reports use stored float64 values. Historical ranks retain repeated responses without interpolation. An unavailable rank stops. The corrected ranks for k=20,40,52,80,160,180,360,720 are 20,39,51,77,153,172,343,685. A corrected rank is not always an empirical pinball minimizer: k=52 selects rank 51 instead of the ordinary .95 rank 50. At integral $k\tau$, the upper adjacent order statistic is an endpoint of the real-arithmetic empirical minimizer interval. Neither fact establishes future coverage.

For S, the translated peak is computed first, current B is subtracted before ranking, then B is added. Final losses use direct stored error P−C, not an algebraically equivalent differently rounded headroom residual. The fitter evaluates its complementary weight as $1-\tau$; final scoring converts 19/20 and 1/20 separately to floats. Display rounding never defines a breach, tie or model selection. Null conditional severities and relative differences retain explicit denominator reasons.

QR implements mature two-block convex ADMM (Boyd et al., 2011), with 34 coefficients, an unpenalized intercept, penalty $\rho=1/W$, at most 10000 iterations, absolute tolerance 10^{-6} and relative tolerance 10^{-5}. Training checks finite values and linear-system solve residuals. Acceptance requires both primal and dual ADMM residual thresholds and a separate feasible-dual objective-gap condition. These numerical checks do not certify a statistical forecasting law.

$$J_\lambda=W^{-1}\sum_i \rho_\tau(y_i-X_i b)+(\lambda/2)\|\beta\|^2;\quad D_\lambda(v)=v^T y/W-\|Z^T v/W\|^2/(2\lambda).$$

Here y=G/a, X includes an intercept and Z its standardized slopes. The candidate dual vector satisfies $\tau-1\le v\le\tau$ and exact zero sum on an inward 2^{-40} dyadic grid; integer box limits use the exact binary-float representation of τ. The checksum checks the represented vector exactly, while objectives remain floating-point evaluations. A nonnegative Fenchel/stationarity decomposition evaluates the gap without subtracting similar objectives; the objective difference must agree within 256 machine epsilons times $\max(1,|J_\lambda|,|D_\lambda|)$, and the stable gap must be at most $10^{-6}(1+|J_\lambda|)$. This is not interval-arithmetic certification.

Equivalent peak parameterization

This is not generally ordinary zero-centered ridge on the peak. At a fixed refit κ, the standardized magnitude coordinate and transformed coefficients satisfy

$$B_i=\mu_{\kappa B}+s_{\kappa B}z_{i,\kappa,B};\quad \alpha_\kappa=\mu_{\kappa B}+a_\kappa\beta_{0,\kappa};\quad \theta_\kappa=a_\kappa\beta_\kappa+s_{\kappa B}e_B.$$

Multiplying the headroom objective by its positive response scale gives the equivalent unprojected peak objective with a shifted penalty:

$$W^{-1}\sum_{i\in T}\rho_\tau(P_{i,H}-\alpha_\kappa-z_{i,\kappa}^T\theta_\kappa)+[\lambda/(2a_\kappa)]\|\theta_\kappa-s_{\kappa B}e_B\|_2^2.$$

This real-arithmetic parameterization includes the scale factor and floor-centered penalty; the intercept remains unpenalized. The issued report is $\max\{B_t,\alpha_\kappa+z_{t,\kappa}^T\theta_\kappa\}$. Adding an issue-known B translates conditional quantiles of G into those of P. Raising a report below a known B≤P to B weakly reduces realized pinball loss. These elementary facts neither guarantee the fitted affine specification nor establish coverage.

Appendix D. Explicitly post-result report diagnostics

After viewing and publishing the complete primary results, we recorded a descriptive floor-binding/horizon-coherence plan and evaluated the already accepted forecasts. These checks are not part of the 98 before-execution contrast menu, and they modify no forecast or tuning choice. The original 90 configurations and 45 source/window/method pairs are retained in post-result-floor.csv and post-result-horizon.csv, with hashes and process resources in their separate audit. Every common realized target satisfies P(H6)≤P(H24). Those files cover the original QR/KNN/scaled-KNN study and its historical controls. The later AR diagnostics are separate files; the original tables and forecasts are unchanged.

Table D1. Original QR/KNN/scaled-KNN horizon-coherence diagnostics; the later AR addition is not included. A crossing means stored C(H6)>C(H24), with no tolerance used to redefine the event. Maximum gap is displayed in USD millions (FX divided by 10^6; Treasury already in that unit). The distinct financial quantities are not pooled.

Source	W	Method	Crossings	Origins	Cross %	Max gap
FX	720	QR	640	6625	9.660	18.742
FX	720	KNN	22	6625	0.332	1.930
FX	720	Scaled KNN	48	6625	0.725	1.775
FX	2880	QR	140	6625	2.113	5.303
FX	2880	KNN	0	6625	0.000	0.000
FX	2880	Scaled KNN	18	6625	0.272	1.772
Treasury	52	QR	91	277	32.852	139288.627
Treasury	52	KNN	39	277	14.079	8202.000
Treasury	52	Scaled KNN	72	277	25.993	124615.389
Treasury	260	QR	80	277	28.881	79417.646
Treasury	260	KNN	0	277	0.000	0.000
Treasury	260	Scaled KNN	25	277	9.025	18835.763

Historical short Treasury controls also cross: rolling F/S on 38/37 issues, frozen F/S on 39/39, and one-state F/S/K on 20/20/3. Treasury W260 one-state K crosses on 13; its other historical controls and all listed FX historical controls have zero recorded crossings. Their H-specific mature sets differ, so common-set construction monotonicity is no guarantee across these procedures. The two selected k=W=52 KNN reports equal their F counterparts, including the 39 crossings.

Exact floor equality describes a stored report, not an observed negative latent prediction. Treasury H24 QR W52 binds on 65 issues (58 breaches), with another 74 breaches off the floor; W260 binds on three (two breaches), with 82 others. The realized zero-headroom outcomes are 14/277 at H24 and 50/277 at H6. FX has 662/6,625 and 1,523/6,625 respectively. These descriptive event sets do not identify the cause of misspecification or effective independent sample size.

The separate AR addition retains all eight floor configurations and four same-source/window horizon pairs. FX W720/W2880 have no observed crossings over 6,625 origins each; Treasury W52/W260 have 59/26 over 277 (21.300%/9.386%), with maximum positive gaps 11784.473/5347.870 USD millions. Every realized target is ordered. Treasury W52 AR binds its floor on seven issues at each horizon, with five H6 and seven H24 breaches. The other six AR configurations never bind. Those empty-group breach rates are undefined, not zero. These inspected-sample diagnostics establish neither a cause nor a coverage guarantee.

The report-order problem does not by itself justify a particular repair. For example, B=0, P(H6)=P(H24)=1, C(H6)=10 and C(H24)=1 satisfy the target order. Raising the long report to the short one changes its .95 loss from zero to .45. This artificial real-arithmetic example is not source data; it distinguishes a learned report from a known lower bound. No isotonic reconciliation, horizon switching or post-result parameter search was run.

Complete execution, verification and arithmetic-readback records for this supplement are retained in REPRODUCIBILITY_V3.md and the corresponding result-directory README.

A later exploratory diagnostic uses the bounded strict hit $I=1\{P>C\}$, with equality a nonbreach, and the null that its conditional expectation given full issue information is at most $\alpha=1/20$ at every declared issue. It is not unconditional equal-loss testing or exact exceedance identification at a floor atom. Conditional-moment betting and delayed binary evidence are mature precedents (Casgrain et al., 2023, v2; Henzi and Ziegel, 2022, v3). For a fixed alternative u in $[\alpha,1]$, the hit/no-hit factor is u/α or $(1-u)/(1-\alpha)$; its null conditional expectation is at most one. Every globally anchored H+1 phase retains only issues whose previous same-phase target has resolved.

$$E_{n,k} = [1/(1-\alpha)] \int_\alpha^1 (u/\alpha)^k [(1-u)/(1-\alpha)]^{n-k}\, du.$$

$E_{cell} = [1/(H+1)] \sum_{\eta=0}^{H} E_{n(\eta),k(\eta)}$; $\alpha=1/20$.

The prior mixes each whole phase product, then all phases receive equal weight; empty phases have value one. Iterated expectation and nonnegative averaging justify a fixed-terminal e-value for a genuinely fixed causal procedure under the stated null, without independence, stationarity or a financial-magnitude moment assumption. This gives no full-filtration supermartingale or arbitrary-stopping claim. The complete matched family has 56 labels, including aliases, and excludes 24 rolling and 18 one-state reports. Its formal reference threshold is 56/(1/20)=1120. Because this diagnostic and family were chosen after inspecting results, the crossings do not obtain retrospective empirical five-percent familywise error control. The fixed-procedure proof also does not establish individually calibrated e-values or p-values for this same-sample, post-selected analysis: selection need not be measurable at the original issue times. No prior, phase or menu was retuned.

Table D2. All 56 late fixed-terminal evidence diagnostics, rounded to three significant figures; * marks an exact crossing of the fixed 1120 reference threshold. Treasury is the separately fitted Treasury aggregate; F/S/K are the matched frozen controls. A noncrossing is not coverage certification, and evidence magnitude is not a financial-quality ranking. All 896 phase rows and exact rational identities are retained in the supplement.

Source H/W	QR	KNN	Scaled	F	S	K	AR
FX 6/720	104	0.00905	0.00476	0.0112	0.00397	26.5	0.00527
FX 6/2880	0.0378	0.00923	0.00476	0.0183	0.00441	294	0.0125
FX 24/720	8.5e7*	1.18	0.171	0.146	0.0164	270	0.0558
FX 24/2880	0.734	0.153	0.0486	0.0318	0.0159	5.12	0.146
Treasury 6/52	1.31e8*	0.057	0.0537	0.057	0.057	0.38	0.242
Treasury 6/260	2.2e3*	0.637	0.305	0.919	0.919	0.401	1.95
Treasury 24/52	3.81e4*	19.7	0.949	19.7	19.7	5.18	2.46
Treasury 24/260	59.4	0.202	0.13	0.221	0.221	4.53	11

The four reference crossings are FX H24 W720 QR and Treasury H6 W52/W260 and H24 W52 QR. Treasury H24 W260 AR has 59/277 strict hits and evidence 11.0284, below 1120; this does not negate its adverse breach record. Two Treasury QR cells both have 84/277 hits but very different evidence: about 1.311×10^{8} at H6 W52 and 59.4235 at H24 W260. Their phase sizes and hit allocation differ, as do their forecasting tasks. No power calculation, equivalence or sharpness conclusion follows.

Ordinary empirical-rank ablation

After inspecting both source archives and the previous results, we fixed one further eight-cell supplement and published a 16-file execution freeze before its financial calculation. At each original evaluation-segment refit, the added intercept-only report uses the same W completed headrooms and original update schedule as frozen F. It selects the lower empirical .95 pinball quantile with duplicate preservation and no interpolation, then adds the actual current magnitude. There is no new tuning, selected model, source, evaluation subset or changed primary.

$C^{I}_{t} = B_t + G_{(\lceil 19W/20 \rceil),\kappa,H}$; $C^{F}_{t} = B_t + G_{(\lceil 19(W+1)/20 \rceil),\kappa,H}$.

Here $G_{(j),\kappa,H}$ denotes the jth ordered headroom in the original mature pool. For the six cells with W720, W2880 or W260, the two ranks are endpoints of the same ordinary empirical pinball minimizer interval. Only the two W52 cells have noninteger W times τ, yielding ordinary rank 50 versus corrected rank 51; ties can collapse distinctions in general. This is an endpoint/rank sensitivity, not a claim that corrected F is universally non-ERM. Dividing all headrooms by a positive training-response scale and multiplying the fitted intercept back by that scale preserves the rank and the minimizer interval in native units.

Table D3. Complete post-inspection ordinary empirical-rank ablation. Percentages are QR-minus-I and I-minus-F mean-loss changes relative to the right report; negative favors the left. QR means and counts are imported unchanged from the original selected fits. I/F counts are newly reconstructed; columns list I/F/QR strict breaches over 6,625 FX or 277 Treasury origins. H counts source records. Sources are not pooled.

Source	H	W	Ranks I/F	QR−I (%)	I−F (%)	Breaches I/F/QR
FX	6	720	684/685	−1.967	−0.153	337/332/531
FX	6	2880	2736/2737	−14.773	+0.008	365/360/376
FX	24	720	684/685	+31.508	−0.003	432/427/924
FX	24	2880	2736/2737	−3.783	+0.026	359/357/531
Treasury	6	52	50/51	+87.086	−1.995	14/11/84
Treasury	6	260	247/248	+46.058	+3.237	32/28/54
Treasury	24	52	50/51	+208.258	+6.213	37/33/132
Treasury	24	260	247/248	+105.020	+2.353	15/12/84

All eight reconstructed F summaries match every original summary metric exactly. The lower report I is strictly below F at every issue in this archive; its breach count and mean positive overrun increase in all eight cells, while mean unused allowance decreases. Its mean loss improves in only three cells. For example, Treasury H24 W52 I raises loss by 6.213% against F despite using the lower ordinary empirical minimizer. QR remains worse than I in all four Treasury cells, while the matched FX H6 W2880 improvement is 14.773%. The ablation supports these finite signs and tradeoffs under the stipulated convention, without removing the inspected-sample, model-universe or measurement limits.

Complete execution, verification and arithmetic-readback records for this supplement are retained in REPRODUCIBILITY_V3.md and the corresponding result-directory README.

Appendix E. Complete fixed-report preference frontiers

Rational boundaries are exact for saved decimal summaries of floating-point means; this does not assert exact unrounded-loss boundaries. Separately stored columns match the analytic normalized-loss identity within the recorded tolerance.

The main score fixes a shortfall/excess preference of 19:1. We now quantify how rankings of those already issued reports depend on that stipulated preference. For report m let U be its mean positive overrun and V its mean unused allowance, over the same origins. For $0 \le a \le 1$ define:

$J_m(a) = a\,U_m + (1-a)V_m$; $\omega = a/(1-a)$; $J_m(.95) = \text{Ł}_{m,.95}$.

Here a=0 and a=1 are limiting single-component objectives; finite positive ω corresponds to $0<a<1$. For two reports A and B, the difference is $a(U_A-U_B)+(1-a)(V_A-V_B)$. A coefficientwise reduction in both components is preferable for every nonnegative linear weighting. With opposing component differences, a unique crossing can reverse the ordering. This elementary affine geometry is not a new forecasting method. Changing a evaluates fixed actions under other hypothetical preferences; it neither refits them as other quantiles nor establishes dominance over all consistent scores for the .95 quantile (Gneiting, 2011).

Each row follows a from 0 to 1 through its maximal open intervals. NN=KNN; Sc=scaled KNN; F/S/K=matched frozen historical controls; AR=AR residual replay; I=ordinary empirical headroom quantile. A slash separates consecutive intervals. Adjacent displayed intervals tie at their common exact boundary; equals joins identical minimizing lines. Breakpoints are rounded for display. The public results.json retains exact rational boundaries and every endpoint/tie set, all 128 input summaries, 448 pair relations and coefficientwise dominance. Endpoints are evaluated separately and are not inferred from rounded interior thresholds. Ratios $\omega = a/(1-a)$ should be converted from the exact fractions, especially near 1.

Table E1. All eight full-origin frontiers among the same eight fixed reports; 6,625 FX or 277 Treasury origins per row. Units are source native and are never pooled.

Source	H	W	Preferred sequence	Interior a boundaries	At .95
FX	6	720	QR / NN / S	0.952882, 0.964505	QR
FX	6	2880	QR / Sc	0.972817	QR
FX	24	720	QR / NN / AR / S	0.845488, 0.889246, 0.938369	S
FX	24	2880	QR / S	0.957033	QR
Treasury	6	52	QR / AR / NN=F	0.717896, 0.972846	AR
Treasury	6	260	AR / K	0.960158	AR
Treasury	24	52	QR / K / S	0.423917, 0.897882	S
Treasury	24	260	K / AR / NN / S	0.778365, 0.970017, 0.995666	AR

Table E2. All eight existing globally anchored H+1 sparse-grid frontiers. Counts are FX 947/265 and Treasury 40/12 for H6/H24. These are descriptive sensitivities with shared training histories.

Source	H	W	Preferred sequence	Interior a boundaries	At .95
FX	6	720	QR / NN / Sc	0.950621, 0.984383	QR
FX	6	2880	QR / Sc	0.978149	QR
FX	24	720	QR / NN / AR / S	0.805398, 0.910344, 0.959437	AR
FX	24	2880	QR / S	0.966600	QR
Treasury	6	52	QR / K / NN=F	0.645837, 0.995462	K
Treasury	6	260	AR / K	0.937124	K
Treasury	24	52	QR / K / AR / NN=F=S	0.693667, 0.940131, 0.990484	AR
Treasury	24	260	K / AR	0.721694	AR

For full Treasury H6 W52, NN and F have identical component means and share their final region. For sparse Treasury H24 W52, NN, F and S share the final region. The complete mathematical comparison retains a report that is optimal only at a boundary even if it has no open interval; absence from a displayed sequence alone is not an endpoint claim. The .95 winner changes in four cells on the sparse grid. A complete envelope of inspected fixed reports is a hindsight descriptive object, not an estimated optimal decision rule, an independence argument or a confidence set.

Appendix F. Complete original model menus and exploratory replay

Table F1. Complete conditional loss menu and same-history frozen controls on 6,625 FX origins, in USD millions. Every selected QR/KNN/scaled cell is shown. Rolling and one-state reference results and all breach/severity quantities remain in the complete CSVs.

H	W	F frozen	S frozen	K frozen	QR	KNN	Scaled KNN
6	720	0.442615	0.450774	0.932433	0.433246	0.436445	0.466292
6	2880	0.452693	0.450801	0.972162	0.385845	0.410583	0.426458
24	720	0.787416	0.755301	1.205808	1.035483	0.836210	0.896753
24	2880	0.763381	0.759371	1.240039	0.734695	0.769768	0.848004

Table F2. Complete selected conditional loss menu and same-history frozen controls on 277 Treasury origins. Losses are in the source's USD-million units, not FX USD. H6/H24 mean six/24 weekly transitions. No monetary average is pooled across sources.

H	W	F frozen	S frozen	K frozen	QR	KNN	Scaled KNN
6	52	2779.054	2786.620	2574.098	5095.519	2779.054	4208.293
6	260	2758.665	2707.244	2287.758	4159.673	2663.561	3127.173
24	52	3715.693	3652.379	5017.741	12165.574	3715.693	5625.667
24	260	2480.306	2478.871	2301.476	5204.760	2257.963	4137.264

Table F3. Complete later AR comparison. Entries are within-reference mean-loss percentages; negative favors AR. S/F have matching source, H, W and refit schedule; QR is the original selected same-cell fit. Counts are AR / S strict breaches over 6,625 FX or 277 Treasury origins. Every row is exploratory; sources are not pooled.

Source	H	W	AR−S %	AR−F %	AR−QR %	Breaches AR / S
FX	6	720	-1.536	+0.278	+2.447	282 / 255
FX	6	2880	-0.546	-0.962	+16.197	338 / 268
FX	24	720	+2.386	-1.790	-25.318	401 / 289
FX	24	2880	+5.147	+4.595	+8.679	444 / 292
Treasury	6	52	-13.444	-13.208	-52.665	22 / 11
Treasury	6	260	-18.887	-20.399	-47.209	31 / 28
Treasury	24	52	+35.125	+32.823	-59.432	56 / 33
Treasury	24	260	-24.118	-24.162	-63.860	59 / 12

Table F4. Preference sensitivity of the four named fixed comparisons. Differences are left minus right. FX amounts are divided by 10^6; Treasury amounts already have USD-million source units. The sources and economic horizons are not pooled. ω is the stipulated shortfall/excess weight ratio; equality holds at a finite displayed crossing, before rounding. Reference definitions remain those of Section 5.

Comparison	dU	dV	Crossing ω	Left preferred
FX H6 primary	-0.010408	-0.920830	None	All $\omega \geq 0$
FX H6 matched	-0.029549	-0.775527	None	All $\omega \geq 0$
FX H24 legacy	+0.167864	-3.463190	20.630898	$\omega < 20.630898$
Treasury H6 primary	+1736.222894	-4968.077943	2.861429	$\omega < 2.861429$

AR estimation definition

After the conditional results had been inspected, we specified one additional path-propagation comparison without an order, volatility or tuning menu. At each original refit κ, put $i_0=\kappa-H-W+1$.

$N_s=c_\kappa+\varphi_\kappa N_{s-1}+\varepsilon_s$.

Fit on the unique transitions $i_0+1,\ldots,\kappa$. The W completed scenarios have W+H levels in their union and W+H−1 unique transitions; overlapping paths do not duplicate transitions in OLS. Coefficients and the historical pool remain fixed until the next original refit, while the query uses the actual N(t).

Using the same fitted coefficients for every residual in a block and replaying its original order from x=N(t) gives the replay identity

$M_{i,h}(x)=N_{i+h}+\varphi_\kappa^h(x-N_i)$.

Set h=0 explicitly to x and report the corrected empirical quantile of the W maximum absolute replay paths. The intercept cancels in this deterministic identity; it is still estimated in the mean recursion. At exact $\varphi_\kappa=1$ the arithmetic agrees with frozen S. Negative or above-one finite estimates are eligible, without clipping. Constant regressors, lost

dispersion and nonfinite computations fail without a substitute model.

This is a mean-filtered ordered-block historical comparison, not the full conditional-volatility filtered simulation of Hull and White (1998) or Barone-Adesi et al. (1999). It does not standardize/rescale innovations, fit GARCH, independently draw future steps or introduce Monte Carlo randomness. Conditional representativeness of historical residual blocks remains unverified; numerical OLS acceptance supplies no coverage or mean-reversion law. Appendix A.4 gives the path relationship, and Appendix C records numerical checks.

AR annual and sparse-grid sensitivity

Direct-S signs reverse for FX H24 W720 on the 265-origin grid (+2.386% overall, −1.884% sparse) and Treasury H24 W52 on only 12 origins (+35.125%, −14.100%). Treasury H24 W260 remains favorable on its sparse grid but is adverse in 2022 and 2023 (+9.928%, +20.224%). Every annual and sparse contrast remains. Observed fitted coefficients lie between zero and one, with refit means around .981–.985 for FX and .676–.850 for Treasury; these descriptions do not identify true mean reversion. Source, horizon, history and model family all matter, and neither average report reduction nor a below-one slope determines a loss ranking.

Appendix G. Reproducibility and pre-specification

The repository preserves the original source-version checks, execution freezes, complete validation menu, chronological selections, accepted forecasts and favorable/adverse outputs. The original and revised manuscript sources are separately retained. Reproduction commands, exact fingerprints, machine observations and consulted-edition records are in REPRODUCIBILITY_V3.md and the repository README.

The original execution and selection freezes are linked at https://github.com/ByronLi0207/financial-mathematics-paper-20261020/commit/1bd5b692fd9cb240763c032fb4e1c203871fad55 and the repository protocol directory. Later exploratory AR, rank and fixed-preference analyses retain their own execution records. New v3 inference has its own scripts, output identities and tests, leaving the archived scientific files intact.

Appendix H. Complete post-result inference and specification outputs

The complete machine-readable DM, HLN, doubled-bandwidth,72-family Holm, sparse-grid coverage and full-sample DQ-type results are retained in results/revision-20261023/inference/. Each NA retains an explicit reason. The accompanying scripts and known-answer tests are supplied together. The exact replicated six-table audit and source identities are in recovery/. Raw FX data and origin-level traces remain private.

The HAC covariance uses centered moment products with denominator n. DQ instruments are the intercept, four matured hit lags and standardized issue-known capacity; its chi-square reference has six degrees of freedom when full rank. The covariance must be finite, full rank and positive definite. The doubled bandwidth remains much shorter than the longest training history, so this limited check does not establish adequate long-run covariance calibration. Sparse grid tests remove overlapping target windows but retain shared training dependence. Treasury grid sizes of 40 or12 are particularly weak for asymptotic reference calibration. The equality-based POF, CC and DQ diagnostics require exact 5% moments; a quantile with atoms can validly be conservative, so rejection by low hit frequency differs from undercoverage. Missing transition-state rows render independence and CC unidentified. POF uses all grid hits, whereas the independence statistic uses transitions conditional on the first hit. The retained CC statistic is their sum, combining these two initialization conventions; it is not the finite-sample likelihood-ratio expression obtained by conditioning both components on the first observation as in Christoffersen (1998, p. 847). Its chi-square p value remains an asymptotic working reference subject to the dependence, boundary and small-sample limitations stated here. HLN was derived for mean-squared-error comparisons; pinball p values are approximate. Holm familywise control requires valid component p values: the adjustment does not validate the working DM references or remove post-inspection selection. All analyses are post-result and reuse inspected samples.

The DQ-type null is $E[Z_t(I_t-.05)]=0$ for the six raw instruments: the intercept, four mature hit lags and issued capacity. The implementation centers/scales capacity across diagnostic rows for numerical conditioning. With the intercept and a finite positive capacity standard deviation, this common invertible column transformation leaves the Wald statistic unchanged in exact arithmetic. Its sample constants are not forecast inputs or claimed available at issue. Rejection concerns these moments; nonrejection leaves other conditional departures untested. The original out-of-sample CAViaR theorem assumes growing estimation and evaluation samples with their evaluation-to-estimation ratio tending to zero. The fixed-window, overlapping-target HAC calculation here is an adaptation, not a direct application of that theorem.

Table H1. Murphy full-breakpoint audit for both original primaries. The extrema of left-minus-right elementary scores span zero in both cases. Integral agreement is an arithmetic identity check.

Source	Breakpoints	Minimum Δ	Maximum Δ	Sample ranking
FX	16146	-0.00443	0.00116	CROSSING
Treasury	640	-0.0184	0.0496	CROSSING

Table H2. Complete fixed 72-member common-history Holm family, including nonsignificant and identical-loss rows. Comparators F/S/K are frozen with the displayed common W. Negative t favors the conditional method. The .05 decision is based on unrounded normal-reference DM p values and Holm adjustment; it is an exploratory working-reference decision conditional on valid component tests. Identical losses retain their NA t/status and bookkeeping p=1.

Source	H	W	Conditional	Ref	DM t	DM p	Holm p	Decision / direction / status
FX	6	720	QR	F	-0.616	0.538	1	At least .05; Favorable; OK
FX	6	720	QR	S	-1.05	0.293	1	At least .05; Favorable; OK
FX	6	720	QR	K	-16.3	2.18e-59	1.46e-57	Below .05; Favorable; OK
FX	6	720	KNN	F	-0.719	0.472	1	At least .05; Favorable; OK
FX	6	720	KNN	S	-1.49	0.136	1	At least .05; Favorable; OK
FX	6	720	KNN	K	-18.3	7.04e-75	4.93e-73	Below .05; Favorable; OK
FX	6	720	Scaled KNN	F	1.24	0.213	1	At least .05; Adverse; OK
FX	6	720	Scaled KNN	S	0.874	0.382	1	At least .05; Adverse; OK
FX	6	720	Scaled KNN	K	-16.3	1.93e-59	1.32e-57	Below .05; Favorable; OK
FX	6	2880	QR	F	-5.87	4.47e-09	2.86e-07	Below .05; Favorable; OK
FX	6	2880	QR	S	-6.16	7.16e-10	4.72e-08	Below .05; Favorable; OK
FX	6	2880	QR	K	-21	3.15e-98	2.24e-96	Below .05; Favorable; OK
FX	6	2880	KNN	F	-3.62	0.0003	0.0162	Below .05; Favorable; OK
FX	6	2880	KNN	S	-4.09	4.35e-05	0.00256	Below .05; Favorable; OK
FX	6	2880	KNN	K	-21.8	2e-105	1.44e-103	Below .05; Favorable; OK
FX	6	2880	Scaled KNN	F	-1.26	0.209	1	At least .05; Favorable; OK
FX	6	2880	Scaled KNN	S	-1.29	0.196	1	At least .05; Favorable; OK
FX	6	2880	Scaled KNN	K	-18	8.12e-73	5.6e-71	Below .05; Favorable; OK
FX	24	720	QR	F	3.22	0.00127	0.0673	At least .05; Adverse; OK
FX	24	720	QR	S	3.05	0.00231	0.115	At least .05; Adverse; OK
FX	24	720	QR	K	-1.54	0.123	1	At least .05; Favorable; OK
FX	24	720	KNN	F	1.53	0.127	1	At least .05; Adverse; OK
FX	24	720	KNN	S	1.69	0.0909	1	At least .05; Adverse; OK
FX	24	720	KNN	K	-4.79	1.63e-06	9.79e-05	Below .05; Favorable; OK
FX	24	720	Scaled KNN	F	1.54	0.123	1	At least .05; Adverse; OK
FX	24	720	Scaled KNN	S	2.12	0.0343	1	At least .05; Adverse; OK
FX	24	720	Scaled KNN	K	-3.88	0.000105	0.00596	Below .05; Favorable; OK

Source	H	W	Conditional	Ref	DM t	DM p	Holm p	Decision / direction / status
FX	24	2880	QR	F	-0.504	0.614	1	At least .05; Favorable; OK
FX	24	2880	QR	S	-0.453	0.65	1	At least .05; Favorable; OK
FX	24	2880	QR	K	-5.56	2.69e-08	1.69e-06	Below .05; Favorable; OK
FX	24	2880	KNN	F	0.185	0.854	1	At least .05; Adverse; OK
FX	24	2880	KNN	S	0.288	0.773	1	At least .05; Adverse; OK
FX	24	2880	KNN	K	-6.1	1.08e-09	7.04e-08	Below .05; Favorable; OK
FX	24	2880	Scaled KNN	F	1.32	0.185	1	At least .05; Adverse; OK
FX	24	2880	Scaled KNN	S	1.51	0.131	1	At least .05; Adverse; OK
FX	24	2880	Scaled KNN	K	-3.91	9.07e-05	0.00526	Below .05; Favorable; OK
Treasury	6	52	QR	F	3.06	0.00224	0.114	At least .05; Adverse; OK
Treasury	6	52	QR	S	3.05	0.00231	0.115	At least .05; Adverse; OK
Treasury	6	52	QR	K	3.17	0.00152	0.0791	At least .05; Adverse; OK
Treasury	6	52	KNN	F	NA	NA	1	At least .05; Identical; IDENTICAL_LOSSES
Treasury	6	52	KNN	S	-1.06	0.29	1	At least .05; Favorable; OK
Treasury	6	52	KNN	K	0.37	0.711	1	At least .05; Adverse; OK
Treasury	6	52	Scaled KNN	F	3.68	0.000231	0.013	Below .05; Adverse; OK
Treasury	6	52	Scaled KNN	S	3.66	0.000254	0.014	Below .05; Adverse; OK
Treasury	6	52	Scaled KNN	K	2.85	0.00439	0.202	At least .05; Adverse; OK
Treasury	6	260	QR	F	2.67	0.00761	0.32	At least .05; Adverse; OK
Treasury	6	260	QR	S	2.74	0.0061	0.262	At least .05; Adverse; OK
Treasury	6	260	QR	K	2.8	0.00514	0.231	At least .05; Adverse; OK
Treasury	6	260	KNN	F	-0.442	0.658	1	At least .05; Favorable; OK
Treasury	6	260	KNN	S	-0.215	0.83	1	At least .05; Favorable; OK
Treasury	6	260	KNN	K	0.961	0.336	1	At least .05; Adverse; OK
Treasury	6	260	Scaled KNN	F	0.945	0.345	1	At least .05; Adverse; OK
Treasury	6	260	Scaled KNN	S	1.09	0.276	1	At least .05; Adverse; OK
Treasury	6	260	Scaled KNN	K	1.93	0.0538	1	At least .05; Adverse; OK
Treasury	24	52	QR	F	2.24	0.0252	0.983	At least .05; Adverse; OK

Source	H	W	Conditional	Ref	DM t	DM p	Holm p	Decision / direction / status
Treasury	24	52	QR	S	2.26	0.0235	0.942	At least .05; Adverse; OK
Treasury	24	52	QR	K	2.16	0.0306	1	At least .05; Adverse; OK
Treasury	24	52	KNN	F	NA	NA	1	At least .05; Identical; IDENTICAL_LOSSES
Treasury	24	52	KNN	S	1.05	0.294	1	At least .05; Adverse; OK
Treasury	24	52	KNN	K	-0.71	0.478	1	At least .05; Favorable; OK
Treasury	24	52	Scaled KNN	F	2.8	0.00516	0.231	At least .05; Adverse; OK
Treasury	24	52	Scaled KNN	S	3	0.00271	0.13	At least .05; Adverse; OK
Treasury	24	52	Scaled KNN	K	0.347	0.729	1	At least .05; Adverse; OK
Treasury	24	260	QR	F	1.67	0.0955	1	At least .05; Adverse; OK
Treasury	24	260	QR	S	1.67	0.0953	1	At least .05; Adverse; OK
Treasury	24	260	QR	K	2.32	0.0202	0.828	At least .05; Adverse; OK
Treasury	24	260	KNN	F	-1.94	0.0523	1	At least .05; Favorable; OK
Treasury	24	260	KNN	S	-1.92	0.0547	1	At least .05; Favorable; OK
Treasury	24	260	KNN	K	-0.0682	0.946	1	At least .05; Favorable; OK
Treasury	24	260	Scaled KNN	F	4.92	8.53e-07	5.29e-05	Below .05; Adverse; OK
Treasury	24	260	Scaled KNN	S	4.92	8.65e-07	5.29e-05	Below .05; Adverse; OK
Treasury	24	260	Scaled KNN	K	2.95	0.00319	0.15	At least .05; Adverse; OK

Appendix I. Post-result / exploratory selected-fit shrinkage diagnostics

The diagnostic replays only the four already selected Treasury C2 fits, without validation retuning. Every C2 capacity and every frozen-F capacity exactly matches the retained origin-level traces. All per-refit coefficient vectors, standardized training transforms, response scales, norm and native contribution summaries are retained in results/revision-20261023/round2/shrinkage/. The contribution uses evaluation queries, not centered training rows; aggregate means weight the actual number of queries in each block. The intercept-only counterfactual holds the fitted intercept fixed and removes slopes, whereas I refits an intercept by the lower empirical quantile. These are distinct comparisons.

Table I1. Treasury selected C2 coefficients and effects. Norms are in standardized/response-normalized coordinates. Mean absolute projected slope effect is in source USD millions. Loss changes are descriptive and all adverse F comparisons are retained.

H	W	λ	Refits	$\|\beta\|_2$ range	Mean |effect|	Δ I %	Δ F %
6	52	10	22	0.006541–0.01937	366.7	-0.6063	-2.589
6	260	10	22	0.004323–0.007924	136.8	-1.038	2.166
24	52	10	22	0.009692–0.01725	832.3	-0.2432	5.955
24	260	10	22	0.001497–0.01264	453	-2.192	0.1097

Table I2. Pointwise C2 comparisons with I and F. Max |Δ| is capacity difference in source USD millions; ρC includes the shared issue floor, ρG removes it. J is strict-breach-set Jaccard. Hits show C2/reference and XOR counts discordant breach occasions.

H	W	Ref	Max |Δ|	ρC	ρG	J	Hits	XOR
6	52	I	2096	0.9999	0.9992	0.9333	15/14	1
6	52	F	9113	0.9965	0.9776	0.7333	15/11	4
6	260	I	967.8	1	0.9997	1	32/32	0
6	260	F	3524	0.9995	0.9963	0.875	32/28	4
24	52	I	6126	0.9998	0.9982	0.9211	36/37	3
24	52	F	1.048e+04	0.9978	0.9905	0.9167	36/33	3
24	260	I	3082	0.9996	0.9791	1	15/15	0
24	260	F	4967	0.9987	0.926	0.8	15/12	3

I uses rank $\lceil 19W/20 \rceil$ and F uses $\lceil 19(W+1)/20 \rceil$ from the same frozen mature pool. At W=260 these distinct ranks are adjacent endpoints of the empirical pinball minimizer interval; I is not a unique minimizer. All four selected penalties remain at the menu upper endpoint. The diagnostic quantifies the executed fits; it does not identify a causal feature effect or establish statistical equivalence to a historical quantile.

Appendix J. Classical bound proof and exploratory L outputs

Proof of Proposition 1. Let $M=\max_{1\le h\le H}|S_h|$ and let η be the first h with $|S_h|>c$. Split each event $\{\eta=k\}$ by the sign of S_k. The remaining sum $R_k=S_H-S_k$ is independent of information through k and symmetric. On a positive first crossing, $R_k\ge 0$ ensures $|S_H|>c$; on a negative crossing, $R_k\le 0$ does so. Each half-axis has probability at least one half, including atoms at zero. Summing disjoint first-crossing events yields $\Pr(M>c)\le 2\Pr(|S_H|>c)$. This includes c=0 directly. The triangle inequality gives $\max_{0\le h\le H}|x+S_h|\le|x|+M$, while right continuity of the endpoint distribution gives $\Pr(|S_H|>q)\le 0.025$. These facts prove the stated corollary with strict breaches even for distributions with atoms.

The claim can fail without independence: take $X_1=Z$ and $X_2=-Z$, where Z takes −1 and1 equally. Both marginals are symmetric, but $S_2=0$ and M=1. At x=q=0 the peak exceeds the threshold with probability one. This exact counterexample is separate from the positive-AR simulation. A conditional use of the proof must establish the corresponding premises under the conditional law.

The simulation uses independently initialized stationary Gaussian increment paths with unit marginal variance and innovations of variance $1-\varphi^2$. The known endpoint variance is $H+2\sum_{j=1}^{H-1}(H-j)\varphi^j$, so the absolute-endpoint 0.975 quantile uses the normal probability 0.9875. Starting x=0 makes this an unconditional population-threshold experiment. It isolates serial dependence from empirical-quantile estimation error. Each cell has 100000 paths and a fixed recorded seed; Monte Carlo standard errors accompany observed rates. Positive φ violates independence but does not logically require violation of the numerical bound.

Table J1. Original positive-increment-AR oracle-quantile simulation, retained unchanged. Rates/SE are percentages; Δ5 is percentage points from 5%. Endpoint tails are evaluated at the same threshold. Every specified cell is retained.

H	φ	Oracle q	Max rate %	Endpoint %	Δ5 pp	MC SE %
6	0	5.49	3.035	2.454	-1.965	0.05425
6	0.7	10.09	2.597	2.573	-2.403	0.05029
6	0.85	11.63	2.55	2.55	-2.45	0.04985
6	0.98	13.19	2.557	2.557	-2.443	0.04992
24	0	10.98	3.819	2.504	-1.181	0.06061
24	0.7	24.6	2.772	2.475	-2.228	0.05191
24	0.85	33.39	2.619	2.525	-2.381	0.0505
24	0.98	49.78	2.572	2.572	-2.428	0.05006

Table J2. Every post-result / exploratory L comparison. Negative t/Δloss favors L. Holm uses only this new 16-comparison family; the original 72-member family is unchanged. Component p values are approximate working references with shared-training and post-inspection limitations.

Source	H	W	Ref	Δloss %	DM t	DM p	HLN p	Holm p
FX	6	720	F	30.93	7.891	2.988e-15	3.658e-15	4.483e-14
FX	6	720	QR	33.76	6.608	3.908e-11	4.37e-11	5.081e-10
FX	6	2880	F	30.39	7.79	6.695e-15	8.123e-15	9.373e-14
FX	6	2880	QR	52.98	12.83	1.089e-37	3.451e-37	1.743e-36
FX	24	720	F	10.85	1.848	0.06462	0.06562	0.5169
FX	24	720	QR	-15.71	-1.673	0.09434	0.09556	0.6604
FX	24	2880	F	7.941	2.058	0.03958	0.04032	0.3958
FX	24	2880	QR	12.16	1.53	0.1259	0.1273	0.7002
Treasury	6	52	F	3.37	0.9707	0.3317	0.3422	0.7002
Treasury	6	52	QR	-43.62	-2.778	0.005471	0.006887	0.06018
Treasury	6	260	F	-11.77	-1.345	0.1785	0.1884	0.7002
Treasury	6	260	QR	-41.49	-3.031	0.002441	0.003236	0.02929
Treasury	24	52	F	26.32	1.569	0.1167	0.1522	0.7002
Treasury	24	52	QR	-61.42	-1.991	0.04645	0.06948	0.418
Treasury	24	260	F	9.868	1.092	0.2746	0.3183	0.7002
Treasury	24	260	QR	-47.64	-1.516	0.1294	0.1664	0.7002

Corrected level mean reversion and negative-increment sensitivities

The fitted AR coefficient in the earlier replay describes position levels, not increments. Table J1 instead specifies a persistent increment process and remains an additional stress mechanism. The corrected level process below is closer to the fitted equation. Its centered Gaussian law and unit stationary variance are illustrative; they do not reproduce either financial source or its fitted intercept, innovation distribution and changing parameters.

$N_h=\varphi N_{h-1}+\varepsilon_h$, $\varepsilon_h \sim N(0,1-\varphi^2)$, $\mathrm{Var}(N_h)=1$ under stationarity.

Differencing a stationary level AR gives negatively correlated increments for positive φ, with the following reference correlation. This formula concerns the stationary random-start law; a fixed starting level changes conditional moments. A differenced level AR is not the negative-coefficient increment AR used as the second sensitivity family.

$X_h=N_h-N_{h-1}$, $\mathrm{Var}(X_h)=2(1-\varphi)$, $\rho_X(k)=-(1-\varphi)\varphi^{k-1}/2$, $k\geq 1$.

$D_H=N_H-N_0 \sim N(0,2(1-\varphi^H))$ under stationarity.

$D_H \mid N_0=x \sim N((\varphi^H-1)x,1-\varphi^{2H})$.

The primary level coefficients are 0.68, 0.85 and 0.98, with the original two horizons. Starts are either sampled from the stationary level distribution or fixed at its absolute-value 95th percentile. The same stationary-marginal endpoint threshold is held across starts, matching a population analogue of the paper's whole-training endpoint pool. At a fixed tail start it is not the conditional endpoint quantile. Positive and negative fixed starts have the same absolute-peak law by Gaussian sign symmetry.

$q_H^{\mathrm{marg}}=\Phi^{-1}(0.9875)\sqrt{(2(1-\varphi^H))}$, $L=|N_0|+q_H^{\mathrm{marg}}$.

The absolute-level 95th percentile uses the normal probability 0.975. Because a signed-level 95th percentile instead uses probability 0.95, every such alternative is retained separately in Table J4. For fixed starts, Table J4 also reports a different conditional-endpoint oracle threshold. It solves the folded-normal CDF using the displayed conditional mean and variance, and shares exactly the same paths with its marginal-threshold partner. Neither threshold restores independent increments or conditional zero-centered symmetry; neither is an oracle for the path maximum.

Table J3. Corrected primary simulation cases. Each row uses 100000 independent paths and includes the issue position in the peak. Rate/SE/endpoint are percentages; Δ5 is percentage points from 5%. Level rows use the stationary-marginal endpoint quantile; increment rows use their own known endpoint quantile with a zero starting position. Stat. denotes a stationary random start; Abs95 is the absolute-level 95th percentile. End measures the absolute change from the issue position exceeding the endpoint threshold. All seeds and exact counts are retained in the correction CSV.

Process/start	H	AR	q	Hits	Peak %	Δ5 pp	SE %	End %
Level/stat.	6	0.68	3.009	175	0.175	-4.825	0.01322	2.462
Level/Abs95	6	0.68	3.009	0	0	-5	0	10.45
Level/stat.	6	0.85	2.502	300	0.3	-4.7	0.01729	2.553
Level/Abs95	6	0.85	2.502	14	0.014	-4.986	0.003741	8.227
Level/stat.	6	0.98	1.071	913	0.913	-4.087	0.03008	2.434
Level/Abs95	6	0.98	1.071	372	0.372	-4.628	0.01925	3.623
Increment/zero	6	-0.3	4.244	3563	3.563	-1.437	0.05862	2.524
Increment/zero	6	-0.6	3.127	4439	4.439	-0.561	0.06513	2.558
Level/stat.	24	0.68	3.17	636	0.636	-4.364	0.02514	2.495
Level/Abs95	24	0.68	3.17	1	0.001	-4.999	0.001	11.42
Level/stat.	24	0.85	3.138	402	0.402	-4.598	0.02001	2.52
Level/Abs95	24	0.85	3.138	1	0.001	-4.999	0.001	11.16
Level/stat.	24	0.98	1.965	768	0.768	-4.232	0.02761	2.468
Level/Abs95	24	0.98	1.965	85	0.085	-4.915	0.009216	6.201
Increment/zero	24	-0.3	8.167	4281	4.281	-0.719	0.06401	2.476
Increment/zero	24	-0.6	5.701	5032	5.032	0.032	0.06913	2.516

All specified mean-reverting level cases have observed peak-breach rates below 5%, including some zero counts at fixed starts. Conservativeness here concerns only these mechanisms and thresholds. A large starting position can move toward zero in conditional mean, producing endpoint displacement without a larger absolute peak; innovations can still overshoot. Zero counts do not prove zero probability: for one prespecified 100000-path cell, the one-sided 95% binomial upper bound is 0.002996%, without a simultaneous-family claim. A fixed-start marginal threshold remains distinct from conditional endpoint calibration.

The negative-increment family keeps unit stationary increment variance, a zero starting position and coefficients −0.3 and −0.6; cumulative levels in that family are not stationary. The largest observed peak-breach rate is 5.032% at H=24 and increment coefficient -0.6. Its 0.032-percentage-point excess is smaller than its 0.06913-percentage-point Monte Carlo SE. This observation is not evidence of a population violation of the 5% threshold, and dependence precludes a general Lévy guarantee regardless of the observed direction.

Table J4. All prespecified anchor-definition and conditional-endpoint-oracle sensitivities. Each row has 100000 paths. Signed95 is the signed-level 95th percentile, while Abs95 is the absolute-level 95th percentile. Marginal/conditional thresholds on a given path cell share the same seed; these are not independent replications. Full seeds, endpoint rates, SE and exact parameters remain in simulation.csv.

H	φ	Start	Endpoint q	q	Hits	Peak %	Δ5 pp
6	0.68	Abs95	conditional	3.717	0	0	-5
6	0.68	Signed95	marginal	3.009	2	0.002	-4.998
6	0.68	Signed95	conditional	3.433	1	0.001	-4.999
6	0.85	Abs95	conditional	3.036	0	0	-5
6	0.85	Signed95	marginal	2.502	21	0.021	-4.979
6	0.85	Signed95	conditional	2.84	4	0.004	-4.996
6	0.98	Abs95	conditional	1.146	215	0.215	-4.785
6	0.98	Signed95	marginal	1.071	464	0.464	-4.536
6	0.98	Signed95	conditional	1.117	340	0.34	-4.66
24	0.68	Abs95	conditional	3.92	0	0	-5
24	0.68	Signed95	marginal	3.17	9	0.009	-4.991
24	0.68	Signed95	conditional	3.605	1	0.001	-4.999
24	0.85	Abs95	conditional	3.88	0	0	-5
24	0.85	Signed95	marginal	3.138	3	0.003	-4.997
24	0.85	Signed95	conditional	3.571	0	0	-5
24	0.98	Abs95	conditional	2.298	20	0.02	-4.98
24	0.98	Signed95	marginal	1.965	146	0.146	-4.854
24	0.98	Signed95	conditional	2.179	56	0.056	-4.944

All new paths use the prerecord of two independent random streams, fixed seeds and bounded batches. They use known population endpoint thresholds, so they do not test finite overlapping calibration pools, quantile estimation, non-Gaussian financial innovations or release-time availability. The original positive-increment table, empirical eight-cell L outputs and original primaries remain unchanged.

An analytical horizon boundary, not an additional simulation, makes the limit concrete. For the zero-coefficient member of the stable level-AR family, let all levels, including the initial level, be independent standard normals. For every positive horizon the stationary-marginal endpoint displacement is normal with variance two, so its absolute endpoint oracle below is constant across horizons. Conditional on any finite starting value, the peak-breach probability is:

$q=\sqrt{2}\,\Phi^{-1}(0.9875)$; $\Pr(\max_{0\le h\le H}|N_h|>|x|+q \mid N_0=x)=1-[2\Phi(|x|+q)-1]^H \to 1$.

The distribution function here is standard normal, and the displayed q is marginal rather than conditional. Bounded convergence gives the same unconditional limit under the stationary random start. Even the correct conditional endpoint threshold is finite and horizon-invariant at each fixed start, giving the same limit. Independent levels do not imply independent increments: adjacent increments share a level. This elementary example excludes a general guarantee uniform over stable level processes and horizons. It is neither a violation of the independent-increment inequality nor an adverse result for the specified positive-coefficient, finite-horizon simulation cells.

Appendix K. Complete exploratory aggregate-panel outputs and prospective protocol

The fourteen candidate keys were fixed from current official net-position metadata before financial outcomes. Twelve supply the complete original 706-date grid; PDPOSGSC-G11L21 and PDPOSGSC-G21 lack 457 required dates each. No series was spliced, interpolated or chosen by forecast performance. Each provenance file records actual URL, acquisition time, response hash and reporting-era metadata; raw input access code is supplied. The labels and reporting structure can change across eras, so continuity of a key and a complete calendar alone do not establish economic measurement invariance. There is no pooled independence-based significance claim across these related aggregate series.

Table K1. All 48 panel H/W cells, no favorable selection. F loss is in source USD millions; the next columns are percentage loss changes relative to F. Hits list F/QR33/C2/L, respectively. All methods complete on277 evaluation origins in every cell.

Series key	H	W	F loss	Δ QR %	Δ C2 %	Δ L %	Hits F/QR/C2/L
PDPOSMBS-TOT	6	52	921.7	171	3.238	7.174	21/101/27/11
PDPOSMBS-TOT	6	260	945.6	7.359	1.167	7.333	26/19/27/5
PDPOSMBS-TOT	24	52	1754	215.5	5.095	-17.87	39/145/43/27
PDPOSMBS-TOT	24	260	1612	26.79	1.038	-3.923	25/29/26/22
PDPOSCS-TOT	6	52	530.3	52.75	-5.787	18.28	16/48/23/11
PDPOSCS-TOT	6	260	399.7	1.902	-1.1	17.56	6/15/6/2
PDPOSCS-TOT	24	52	579.4	73.78	1.851	-5.065	16/63/24/11
PDPOSCS-TOT	24	260	569.9	27.46	-3.384	19.63	6/3/6/4
PDPOSFGS-TOT	6	52	400.7	107.2	-4.151	-4.743	16/77/24/13
PDPOSFGS-TOT	6	260	358.3	-1.081	4.883	16.77	6/14/9/0
PDPOSFGS-TOT	24	52	476.2	128.7	13.36	-3.723	36/64/37/37
PDPOSFGS-TOT	24	260	354	30.02	72.24	-4.06	9/21/26/12
PDPOSTIPS-L2	6	52	287.3	217.2	6.738	-1.891	26/125/29/12
PDPOSTIPS-L2	6	260	269.3	66.28	1.78	-7.423	24/45/25/19
PDPOSTIPS-L2	24	52	641	307.4	24.88	-34.6	48/181/69/30
PDPOSTIPS-L2	24	260	532.2	226.9	11.41	-4.519	47/147/59/46
PDPOSTIPS-G11	6	52	42.32	196.7	1.706	13.93	23/118/29/3
PDPOSTIPS-G11	6	260	41.76	90.31	63.18	27.71	14/38/29/1
PDPOSTIPS-G11	24	52	80.85	218.7	135.6	-16.39	61/123/103/16
PDPOSTIPS-G11	24	260	51.21	98.59	37.18	27.11	9/33/30/0
PDPOSTIPS-G2	6	52	181.3	208.6	8.58	19.45	16/96/21/4
PDPOSTIPS-G2	6	260	179.4	63.47	10.78	23	19/36/24/4
PDPOSTIPS-G2	24	52	377.9	277.8	11.77	6.363	31/131/42/27
PDPOSTIPS-G2	24	260	286	109.3	120.1	2.092	23/66/58/20
PDPOSTIPS-G6L11	6	52	133.6	59.3	-1.655	9.669	9/73/12/5
PDPOSTIPS-G6L11	6	260	114.9	-5.717	-1.109	36.27	4/14/4/0
PDPOSTIPS-G6L11	24	52	147	76.48	-0.9743	5.028	10/92/16/4
PDPOSTIPS-G6L11	24	260	133.8	-26.42	-1.297	12.63	2/1/2/0
PDPOSGSC-L2	6	52	1233	215.9	-3.846	20.65	15/129/23/6
PDPOSGSC-L2	6	260	1247	64.4	9.038	-2.638	15/40/20/6
PDPOSGSC-L2	24	52	2271	276.6	182.8	0.9259	46/165/96/27
PDPOSGSC-L2	24	260	2919	41.6	-0.1931	-22.23	33/55/33/24
PDPOSGSC-G2L3	6	52	792.4	121.1	145.3	17.31	24/93/112/15
PDPOSGSC-G2L3	6	260	1004	21.31	22.62	-14.81	28/44/59/12
PDPOSGSC-G2L3	24	52	2102	113	129.6	22.35	47/135/151/49
PDPOSGSC-G2L3	24	260	2677	11.6	39.73	-9.349	55/64/91/40
PDPOSGSC-G3L6	6	52	764.9	87.61	75.44	20.57	12/82/72/9
PDPOSGSC-G3L6	6	260	699.1	0.7876	1.907	3.661	20/15/21/8

Series key	H	W	F loss	Δ QR %	Δ C2 %	Δ L %	Hits F/QR/C2/L
PDPOSGSC-G3L6	24	52	977.4	544.3	299.1	-11.37	38/137/120/20
PDPOSGSC-G3L6	24	260	931	15.52	0.1286	-5.726	20/37/20/17
PDPOSGSC-G6L7	6	52	542.3	253.7	89	0.9519	16/142/79/12
PDPOSGSC-G6L7	6	260	564.9	31.61	71.05	-10.15	20/42/49/15
PDPOSGSC-G6L7	24	52	993.4	264.8	264.8	1.686	43/217/215/49
PDPOSGSC-G6L7	24	260	699	146.4	142.6	5.191	22/77/67/21
PDPOSGSC-G7L11	6	52	505.7	175.6	15.72	4.917	25/83/34/11
PDPOSGSC-G7L11	6	260	506.2	8.344	-0.06055	1.05	25/31/25/12
PDPOSGSC-G7L11	24	52	1654	94.95	62.4	-20.33	36/105/91/26
PDPOSGSC-G7L11	24	260	1003	230	202.8	1.881	32/94/99/26

The companion prospective-public-release-protocol.md and parameter-binding JSON freeze the four procedures, selected penalties, causal mature-pool schedule and code identities. They separate economic issue dates, actual public-information/forecast creation times and final target-publication times. At ordinary eight-day publication lag, the already published economic October 14 origin would mature for H6 economically November 25 and publicly December 3. The first planned post-freeze origin, economic October 21 released ordinarily October 29, instead has H6 economic completion December 2/public completion December 10; H24 completion April 7 2027/public completion April 15. Holiday and revision schedules require actual future receipts. Fifty-two consecutive future issue origins are planned, with no significance-based stopping; results remain pending and no automated future service is started here.

The separately dated prospective-vintage amendment preserves the accepted historical prefix, identifies the already-public bridge and freezes every future value at its first actual retained capture. This is an explicit join across acquisition vintages, not a reconstructed publisher first-release history. Later revisions are logged without rewriting the main path, mature labels or scores. Capture and forecast artifacts must both meet the recorded deadline and information boundary. Missing or late weeks consume their calendar origins; they cannot shorten the horizon, shift the refit phase, extend the plan or acquire hindsight forecasts. Every target includes the same captured issue anchor and requires all calendar components. The retrospective batch interface requires a full finite series with H outcome records beyond its last query, although its fitted pools and query slices do not use those future values. It is not a prefix-only live forecast interface. Batch functions abort failed cells. Capture, prefix-only issuance and missing-week continuation need a new adapter with its own prior timestamped binding, complete-data preservation, causal and missing-data fixture checks. That adapter is not implemented here. The amendment changes no retrospective result.

Appendix L. Post-result / exploratory mature-baseline methods and complete outputs

Every addition in this appendix is post-result / exploratory. No D1 result replaces either original primary, its parameter selection, the original 72-comparison Holm family or any archived result. The fixed protocol and code/version fingerprints precede the financial executions. The three new methods have 18 complete cells and 6 retained NA cells out of 24. There is no omitted-date scoring or fallback report.

Filter and simulated-path FHS

At refit κ, the W completed origins span $i_0=\kappa-H-W+1$ through $\kappa-H$. Fit a constant-mean Gaussian-QMLE GARCH(1,1) to increments in the same position envelope, containing $W+H-1$ increments. A training-only RMS conditions the fit numerically. A single optimization is allowed; failures remain NA. The coefficients and standardized residual pool are frozen between refits; newly observed increments update the variance state through issue t. In-sample historical filtered volatilities estimated at κ are not historical forecasts made at i. Finite fitted recursion and optimizer termination supply no independence, stationarity or coverage guarantee.

$\Delta_j=\mu_G+\varepsilon_j$, $\varepsilon_j=\sigma_j z_j$, $\sigma_j^2=\omega_G+\alpha_G\varepsilon_{j-1}^2+\beta_G\sigma_{j-1}^2$.

The subscript G marks local GARCH coefficients, distinguishing them from the regression intercept and fixed-report preference ratio used elsewhere. The POT shape and scale below carry hats and refer to a separate tail fit.

FHS draws 2048 iid residual paths with replacement, using a seed bound to source, H, W and κ; draws are shared within that refit block. Each path recursively updates GARCH variance and signed increments from actual N_t. Its simulated maximum includes h=0, and the empirical headroom quantile uses rank $\lceil 19(M+1)/20 \rceil$. The fitted mean equation is constant; resampling retains the stored standardized residuals without further empirical recentering or rescaling, so their finite resampling distribution need not have exactly zero mean or unit variance. This is a same-target adaptation, rather than an exact replica of the original derivative-portfolio application.

$P_{t,H}^{(s)}=\max_{0\le h\le H}|N_t+\sum_{\ell=1}^{h}\Delta_\ell^{(s)}|$, $G_{t,H}^{(s)}=P_{t,H}^{(s)}-B_t$.

Peak-headroom POT adaptation

Normalize each mature headroom by the frozen filter's one-step volatility at its historical starting origin. The threshold leaves $\lfloor 0.1W \rfloor$ potential upper observations; strict exceedances and ties determine the actual m. A minimum of 20 excesses, valid GPD support, positive scale and the declared shape regularity gate are required. Thus W52 is structurally ineligible. The single declared Nelder–Mead fit uses no alternate start, threshold or fallback. The quantile below uses the continuous exponential limit at zero shape.

$R_i=G_{i,H}/\hat{\sigma}_{i+1|\kappa}$, $\hat{q}_{0.95}=u+(\beta/\xi)[((m/W)/0.05)^{\xi}-1]$, $C_t=B_t+\hat{\sigma}_{t+1|\kappa}\max\{0,\hat{q}_{0.95}\}$.

McNeil and Frey (2000) fit tails of filtered one-step return innovations and simulate multi-period sums. The present headroom-tail scaling is not that original procedure. Dependence on the signed anchor relative to volatility, drift and the future variance path remains after a single-scale normalization; overlapping mature labels remain dependent. These are substantive approximation limits, not cured by successful GPD optimization.

Quantile-regression forest

The forest uses the same 33 causal predictors, training-only transforms and response scaling as QR. The fixed 100-tree regression partitions have max_features=11, minimum leaf 5, split 10, maximum depth 10, bootstrap enabled and one worker. All original training rows receive within-leaf mass, averaged over trees; bootstrap chooses partitions but does not multiply these CDF masses. The left inverse at 0.95 selects a positive-mass training response with no interpolation or F rank correction. This bounded fixed configuration is not claimed to optimize QRF or transfer the original consistency theorem to dependent fixed-window data.

$w_i(z)=(1/100)\sum_{b=1}^{100}1\{z_i\in L_b(z)\}/\#\{j:z_j\in L_b(z)\}$, $F(y|z)=\sum_i w_i(z)1\{G_{i,H}/a_\kappa\le y\}$.

Table L1. All 24 planned method cells. Loss is native FX USD or Treasury USD millions. Failed complete cells have NA, regardless of earlier issued reports; no partial average substitutes for the full evaluation.

Source	H	W	Method	Status	Loss	Hits	Δ F %	Δ QR %
FX	6	720	FHS	NA	NA	NA	NA	NA
FX	6	720	PEAK_POT	NA	NA	NA	NA	NA
FX	6	720	QRF	COMPLETE	4.683e+05	402	5.793	8.081
FX	6	2880	FHS	COMPLETE	4.47e+05	278	-1.258	15.85
FX	6	2880	PEAK_POT	COMPLETE	4.675e+05	363	3.269	21.16
FX	6	2880	QRF	COMPLETE	4.324e+05	354	-4.48	12.07
FX	24	720	FHS	NA	NA	NA	NA	NA
FX	24	720	PEAK_POT	NA	NA	NA	NA	NA
FX	24	720	QRF	COMPLETE	9.136e+05	779	16.02	-11.77
FX	24	2880	FHS	COMPLETE	8.111e+05	136	6.255	10.4
FX	24	2880	PEAK_POT	COMPLETE	8.444e+05	369	10.61	14.93
FX	24	2880	QRF	COMPLETE	8.194e+05	600	7.335	11.53
Treasury	6	52	FHS	COMPLETE	2727	7	-1.883	-46.49
Treasury	6	52	PEAK_POT	NA	NA	NA	NA	NA
Treasury	6	52	QRF	COMPLETE	2589	16	-6.83	-49.19
Treasury	6	260	FHS	COMPLETE	2530	4	-8.276	-39.17
Treasury	6	260	PEAK_POT	COMPLETE	2473	17	-10.35	-40.54
Treasury	6	260	QRF	COMPLETE	2930	33	6.198	-29.57
Treasury	24	52	FHS	COMPLETE	5955	0	60.26	-51.05
Treasury	24	52	PEAK_POT	NA	NA	NA	NA	NA
Treasury	24	52	QRF	COMPLETE	3875	44	4.279	-68.15
Treasury	24	260	FHS	COMPLETE	5582	0	125.1	7.249
Treasury	24	260	PEAK_POT	COMPLETE	3276	20	32.09	-37.05
Treasury	24	260	QRF	COMPLETE	3322	62	33.94	-36.17

Table L2. All 48 planned paired comparisons. Tests are post-result working references; NA component tests remain NA. A separate 48-slot Holm family reserves missing/degenerate slots as 1 for adjustment only. This is distinct from claiming an observed valid p value of 1. Negative Δ/t favors the new method; shared training and inspected outcomes limit interpretation.

Source	H	W	Method	Ref	Δloss %	DM t	DM p	HLN p	Holm p
FX	6	720	FHS	F	NA	NA	NA	NA	NA
FX	6	720	FHS	QR	NA	NA	NA	NA	NA
FX	6	720	PEAK_POT	F	NA	NA	NA	NA	NA
FX	6	720	PEAK_POT	QR	NA	NA	NA	NA	NA
FX	6	720	QRF	F	5.793	1.722	0.08512	0.08543	1
FX	6	720	QRF	QR	8.081	2.559	0.01048	0.01057	0.3984
FX	6	2880	FHS	F	-1.258	-0.6564	0.5116	0.5119	1
FX	6	2880	FHS	QR	15.85	5.8	6.64e-09	7.147e-09	3.054e-07
FX	6	2880	PEAK_POT	F	3.269	1.538	0.1241	0.1245	1
FX	6	2880	PEAK_POT	QR	21.16	6.305	2.885e-10	3.175e-10	1.356e-08
FX	6	2880	QRF	F	-4.48	-1.528	0.1264	0.1268	1
FX	6	2880	QRF	QR	12.07	4.672	2.978e-06	3.094e-06	0.000131
FX	24	720	FHS	F	NA	NA	NA	NA	NA
FX	24	720	FHS	QR	NA	NA	NA	NA	NA
FX	24	720	PEAK_POT	F	NA	NA	NA	NA	NA
FX	24	720	PEAK_POT	QR	NA	NA	NA	NA	NA
FX	24	720	QRF	F	16.02	1.953	0.05076	0.05163	1
FX	24	720	QRF	QR	-11.77	-1.872	0.06122	0.06219	1
FX	24	2880	FHS	F	6.255	1.062	0.2881	0.2898	1
FX	24	2880	FHS	QR	10.4	1.367	0.1717	0.1733	1
FX	24	2880	PEAK_POT	F	10.61	3.05	0.002286	0.002379	0.09144
FX	24	2880	PEAK_POT	QR	14.93	2.405	0.01618	0.01659	0.5987
FX	24	2880	QRF	F	7.335	1.08	0.2801	0.2819	1
FX	24	2880	QRF	QR	11.53	1.539	0.1237	0.1251	1
Treasury	6	52	FHS	F	-1.883	-0.2039	0.8384	0.8417	1
Treasury	6	52	FHS	QR	-46.49	-2.66	0.007819	0.009631	0.3049
Treasury	6	52	PEAK_POT	F	NA	NA	NA	NA	NA
Treasury	6	52	PEAK_POT	QR	NA	NA	NA	NA	NA
Treasury	6	52	QRF	F	-6.83	-0.9999	0.3174	0.3279	1
Treasury	6	52	QRF	QR	-49.19	-3.611	0.0003054	0.0004712	0.01313
Treasury	6	260	FHS	F	-8.276	-0.5508	0.5818	0.5897	1
Treasury	6	260	FHS	QR	-39.17	-2.277	0.02278	0.02642	0.7972
Treasury	6	260	PEAK_POT	F	-10.35	-1.265	0.206	0.2162	1
Treasury	6	260	PEAK_POT	QR	-40.54	-3.117	0.00183	0.002474	0.07503
Treasury	6	260	QRF	F	6.198	0.585	0.5585	0.5668	1
Treasury	6	260	QRF	QR	-29.57	-3.238	0.001205	0.001677	0.05062
Treasury	24	52	FHS	F	60.26	1.955	0.05059	0.0747	1
Treasury	24	52	FHS	QR	-51.05	-1.467	0.1423	0.1804	1
Treasury	24	52	PEAK_POT	F	NA	NA	NA	NA	NA
Treasury	24	52	PEAK_POT	QR	NA	NA	NA	NA	NA
Treasury	24	52	QRF	F	4.279	0.2932	0.7693	0.7886	1
Treasury	24	52	QRF	QR	-68.15	-2.395	0.0166	0.02921	0.5987
Treasury	24	260	FHS	F	125.1	11.21	3.542e-29	3.876e-21	1.7e-27
Treasury	24	260	FHS	QR	7.249	0.2367	0.8129	0.8287	1
Treasury	24	260	PEAK_POT	F	32.09	5.485	4.146e-08	9.315e-07	1.866e-06
Treasury	24	260	PEAK_POT	QR	-37.05	-1.219	0.2229	0.2657	1

Source	H	W	Method	Ref	Δloss %	DM t	DM p	HLN p	Holm p
Treasury	24	260	QRF	F	33.94	1.425	0.154	0.1931	1
Treasury	24	260	QRF	QR	-36.17	-1.736	0.08256	0.1133	1

Retained cell failures

fx_hourly H6 W720 FHS: FilteredFitError — single GARCH fit failed convergence or admissibility. Issued observations before failure: 240.

fx_hourly H6 W720 PEAK_POT: FilteredFitError — single GARCH fit failed convergence or admissibility. Issued observations before failure: 240.

fx_hourly H24 W720 FHS: FilteredFitError — single GARCH fit failed convergence or admissibility. Issued observations before failure: 240.

fx_hourly H24 W720 PEAK_POT: FilteredFitError — single GARCH fit failed convergence or admissibility. Issued observations before failure: 240.

nyfed_weekly H6 W52 PEAK_POT: FilteredFitError — POT structurally unavailable: top ten percent has fewer than 20 rows. Issued observations before failure: 0.

nyfed_weekly H24 W52 PEAK_POT: FilteredFitError — POT structurally unavailable: top ten percent has fewer than 20 rows. Issued observations before failure: 0.

The 4 short-history FX filter rejections have successful optimizer flags. Stored coefficients violate only the declared persistence tolerance after all other recorded admissibility conditions are verified. A fixed-coefficient readback invokes no optimizer and issues no new forecast; it reproduces the retained conditional-variance and standardized-residual hashes. No tolerance is relaxed and no rejected cell is rescored. These are numerical admissibility exclusions, not optimizer nonconvergence or evidence that FHS/POT predicts worse. The two short-history Treasury POT exclusions instead follow the minimum-tail-size rule.

$\alpha_G+\beta_G \le 1+10^{-10}$.

Table L3. Every retained NA classification. For boundary fits, δ is fitted persistence minus one and ε is excess over the declared numerical ceiling; optimizer flag 0 denotes success. Strict-tail structural exclusions have no fitted persistence. No missing full-cell loss is imputed.

Source	H	W	Method	Issued	Class	δ	ε	Flag
FX	6	720	FHS	240	Boundary tolerance	1.272e-10	2.721e-11	0
FX	6	720	PEAK_POT	240	Boundary tolerance	1.272e-10	2.721e-11	0
FX	24	720	FHS	240	Boundary tolerance	1.299e-10	2.986e-11	0
FX	24	720	PEAK_POT	240	Boundary tolerance	1.299e-10	2.986e-11	0
Treasury	6	52	PEAK_POT	0	Tail size	NA	NA	NA
Treasury	24	52	PEAK_POT	0	Tail size	NA	NA	NA

The separate 48-slot Holm adjustment concerns only these working-reference comparisons. It does not control selection across every supplement or validate the component reference distributions. Remaining alternatives include canonical innovation-tail EVT, CAViaR and sequential conformal procedures. Neither these added baselines nor the aggregate panel supplies compatible unseen-dealer validation.