

Recovery Summary

May 5, 2026 at 9:19 PM

This explains your doctor's instructions. It is not medical advice. Call your doctor if unsure.

Plan Overview

You were in the hospital for 5 days because of very high blood sugar, high blood pressure, a kidney infection, and kidney problems. Your diabetes was not well controlled and you had stopped taking your blood pressure medicines. The doctors gave you insulin, antibiotics, and other medicines to help you feel better. Your kidney function improved before you went home, and your blood sugar is now under better control.

Medications

Metformin

500 mg · twice daily with meals

Take after food. Watch for vomiting.

Glimepiride

2 mg · once daily before breakfast

Take before food. Do not skip meals.

Amlodipine

5 mg · once daily in morning

Do not stop suddenly.

Telmisartan

40 mg · once daily

Avoid foods high in potassium (bananas, oranges).

Furosemide (water pill)

20 mg · once daily in morning

Take in morning for 14 days. Watch for dizziness.

Nitrofurantoin (antibiotic)

100 mg · twice daily

Take after food for 5 days. Finish all pills.

Atorvastatin (cholesterol medicine)

40 mg · once daily at night

Avoid grapefruit juice.

Ecosprin (baby aspirin)

75 mg · once daily

Take after food.

Shelcal-CT (calcium and vitamin D)

1 tablet · once daily

Take after food.

Symptoms to Watch

- trouble breathing
- chest pain
- urinating much less than normal
- blood sugar above 300
- blood pressure above 180/110
- severe leg swelling
- confusion
- unable to take your medicines

Upcoming Follow-up

See kidney doctor (Dr. S. Bhandare)

2025-05-04 at 10:00 AM · Nephrology department - Tuesdays

See diabetes doctor (Dr. Meenakshi Iyer)

2025-05-08 · Diabetology department - Thursdays

See dietician (Ms. Padma Rao)

as soon as possible · Diet department - Mondays and Wednesdays