

**Problem Set 10**

Fall 2014

**Issued:** Saturday, November 15

**Due:** Friday, November 21

---

---

**Problem 10.1**

Read Chapter 14 of Keener.

**Problem 10.2**

Consider the general linear model with normality:

$$Y \sim N(X\beta, \sigma^2 I), \quad \beta \in \mathbb{R}^p, \quad \sigma^2 > 0.$$

If the rank  $r$  of  $X$  equals  $p$ , show that  $(\hat{\beta}, S^2)$  is a complete sufficient statistic.

**Problem 10.3**

*Inverse linear regression.* Consider the model for simple linear regression:

$$Y_i \sim \beta_1 + \beta_2(x_i - \bar{x}) + \epsilon_i, \quad i = 1, \dots, n,$$

studied in Section 14.5.

- (a) Derive a level- $\alpha$  test of  $H_0 : \beta_2 = 0$  versus  $H_1 : \beta_2 \neq 0$ .
- (b) Let  $y_0$  denote a “target” value for the mean of  $Y$ . The regression line  $\beta_1 + \beta_2(x - \bar{x})$  achieves this value when the independent variable  $x$  equals

$$\theta = \bar{x} + \frac{y_0 - \beta_1}{\beta_2}.$$

Derive a level- $\alpha$  test of  $H_0 : \theta = \theta_0$  versus  $H_1 : \theta \neq \theta_0$ . (Hint: You may want to find a test first assuming  $y_0 = 0$ . After a suitable transformation, the general case should be similar.)

- (c) Use duality to find a confidence region, first discovered by Fieller (1954), for  $\theta$ . Show that this region is an interval if the test in part (a) rejects  $\beta_2 = 0$ .

**Problem 10.4**

Consider a regression version of the two-sample problem in which:

$$Y_i = \begin{cases} \beta_1 + \beta_2 x_i + \epsilon_i & i = 1, \dots, n_1; \\ \beta_3 + \beta_4 x_i + \epsilon_i & i = n_1 + 1, \dots, n_1 + n_2 = n, \end{cases}$$

with  $\epsilon_1, \dots, \epsilon_n$  i.i.d. from  $N(0, \sigma^2)$ . Derive a  $1 - \alpha$  confidence interval for  $\beta_4 - \beta_2$ , the difference between the two regression slopes.

**Problem 10.5**

A variable  $Y$  has a log-normal distribution with parameters  $\mu$  and  $\sigma^2$  if  $\log Y \sim N(\mu, \sigma^2)$ .

- (a) Find the mean and density for the log-normal distribution.
- (b) If  $Y_1, \dots, Y_n$  are i.i.d. from the log-normal distribution with unknown parameters  $\mu$  and  $\sigma^2$ , find the UMVU for  $\mu$ .
- (c) If  $Y_1, \dots, Y_n$  are i.i.d. from the log-normal distribution with parameters  $\mu$  and  $\sigma^2$ , with  $\sigma^2$  a known constant, find the UMVU for the common mean  $\eta = EY_i$ .
- (d) In simple linear regression,  $Y_1, \dots, Y_n$  are independent with  $Y_i \sim N(\beta_1 + \beta_2 x_i)$ . In some applications this model may be inappropriate because the  $Y_i$  are positive; perhaps  $Y_i$  is the weight or volume of the  $i$ th unit. Suggest a similar model without this defect based on the log-normal distribution. Explain how you would estimate  $\beta_1$  and  $\beta_2$  in your model.