General layer of the GI tract:
• Mucosa
○ Epithelium
○ Lamina Propriae
○ Muscularis mucosa
• Submucosa
○ Glandular tissues
○ Submucosal plexus
Muscularis externa
○ Longitudinal muscle
Circular muscle
Myenteric plexus
• Serosa
○ Adventitia
○ Epithelium
• Oesophagus
○ Mucosa:
 Stratified epithelium (Protection), transition to simple squamous epithelium into the stomach
○ Submucosa
Muscularis externa
Contains both skeletal (upper part for peristalsis) and smooth muscles
○ Serosa
No visceral peritoneum
Stomach: Three parts (Fundus, body, antrum pylorus*)
○ Mucosa
Rugae folding in the internal surface to increase surface area
Squamous epithelium at surface, columnar epithelium in gastric pit.
○ Submucosa
Glands in the gastric pit, parietal cells secrete HCl, chief cells in gastric glands secrete pepsinogen that
activate in low pH,
 Invaginations comparing to smoother surface in the duodenum
Muscularis externa
Diaphragm contract create oesophageal-pyloric region
Outer to inner: Longitudinal, circular, oblique
 Ring of muscle at the pylorus region - pyloric sphincter, no oblique at pylorus
Brunner's gland at pyro-duodenum junction, secrete bicarbonates.
○ Serosa
Small intestines: Three parts (Duodenum, Jejunum, Ileum)
○ Mucosa:
 Villi structure lined by simple colunmar epithelial enterocytes, project outwards increase surface area,
containing artery and vein and lacteal duct for lipid.

• iviicrovilli p	present on surface of enterocytes, increase surface area.
∘ Submucosa:	
· Crypts of	ieberkuhn: Goblet cells (mucous), paneth cells (antimicrobial peptides), and stem cells
► Uniform v	lli surface compared to invaginations in the stomach
→ Greater for	ld in the duodenum & jejunum compared to the ileum
 Muscularis exte 	rna
∘ Serosa	
arge intestine: Cecu	m, colon, rectum
∘ Mucosa: Flat co	mparing to small intestine
○ Submucosa:	
Lymph ag	gregates
→ Goblet cel	Is (mucous)
 Muscularis exte 	rna:
► Thicker ci	cular and longitudinal
► Taenia Co	li - 3 strands of longitudinal muscle along the length up to the rectum, allow contraction
∘ Serosa	
	ion: Pyloric junction muscle, Brunner's gland secrete bicarbonate nsition from simple to stratified epithelium(from ectoderm), separated by the dentate line.
o-rectal junction: Tra	nsition from simple to stratified epithelium(from ectoderm), separated by the dentate line. er derived from epithelial tissues, causes colon cancer
o-rectal junction: Tra	nsition from simple to stratified epithelium(from ectoderm), separated by the dentate line. er derived from epithelial tissues, causes colon cancer
o-rectal junction: Tra	nsition from simple to stratified epithelium(from ectoderm), separated by the dentate line. er derived from epithelial tissues, causes colon cancer
o-rectal junction: Tra	nsition from simple to stratified epithelium(from ectoderm), separated by the dentate line. er derived from epithelial tissues, causes colon cancer
o-rectal junction: Tra	nsition from simple to stratified epithelium(from ectoderm), separated by the dentate line. er derived from epithelial tissues, causes colon cancer
o-rectal junction: Tra	nsition from simple to stratified epithelium(from ectoderm), separated by the dentate line. er derived from epithelial tissues, causes colon cancer
o-rectal junction: Tra	nsition from simple to stratified epithelium(from ectoderm), separated by the dentate line. er derived from epithelial tissues, causes colon cancer
o-rectal junction: Tra	nsition from simple to stratified epithelium(from ectoderm), separated by the dentate line. er derived from epithelial tissues, causes colon cancer
o-rectal junction: Tra	nsition from simple to stratified epithelium(from ectoderm), separated by the dentate line. er derived from epithelial tissues, causes colon cancer
o-rectal junction: Tra	nsition from simple to stratified epithelium(from ectoderm), separated by the dentate line. er derived from epithelial tissues, causes colon cancer
o-rectal junction: Tra	nsition from simple to stratified epithelium(from ectoderm), separated by the dentate line. er derived from epithelial tissues, causes colon cancer
o-rectal junction: Tra	nsition from simple to stratified epithelium(from ectoderm), separated by the dentate line. er derived from epithelial tissues, causes colon cancer
o-rectal junction: Tra	nsition from simple to stratified epithelium(from ectoderm), separated by the dentate line. er derived from epithelial tissues, causes colon cancer
o-rectal junction: Tra	nsition from simple to stratified epithelium(from ectoderm), separated by the dentate line. er derived from epithelial tissues, causes colon cancer
o-rectal junction: Tra	nsition from simple to stratified epithelium(from ectoderm), separated by the dentate line. er derived from epithelial tissues, causes colon cancer
o-rectal junction: Tra	nsition from simple to stratified epithelium(from ectoderm), separated by the dentate line. er derived from epithelial tissues, causes colon cancer
o-rectal junction: Tra	nsition from simple to stratified epithelium(from ectoderm), separated by the dentate line. er derived from epithelial tissues, causes colon cancer
o-rectal junction: Tra	nsition from simple to stratified epithelium(from ectoderm), separated by the dentate line. er derived from epithelial tissues, causes colon cancer
o-rectal junction: Tra	nsition from simple to stratified epithelium(from ectoderm), separated by the dentate line. er derived from epithelial tissues, causes colon cancer
o-rectal junction: Tra	nsition from simple to stratified epithelium(from ectoderm), separated by the dentate line. er derived from epithelial tissues, causes colon cancer
o-rectal junction: Tra	nsition from simple to stratified epithelium(from ectoderm), separated by the dentate line. er derived from epithelial tissues, causes colon cancer
o-rectal junction: Trai	nsition from simple to stratified epithelium(from ectoderm), separated by the dentate line. er derived from epithelial tissues, causes colon cancer