

Procrastination and Dementia

Cormac Monaghan^{1,2}, Rafael de Andrade Moral³, and Joanna McHugh Power²

¹Hamilton Institute, Maynooth University, Ireland

²Department of Psychology, Maynooth University, Ireland

³Department of Mathematics and Statistics, Maynooth University, Ireland

Author Note

Cormac Monaghan  <https://orcid.org/https://orcid.org/0000-0001-9012-3060>

Rafael de Andrade Moral  <https://orcid.org/https://orcid.org/0000-0002-0875-3563>

Joanna McHugh Power  <https://orcid.org/https://orcid.org/0000-0002-7387-3107>

Correspondence concerning this article should be addressed to Cormac Monaghan, Email:
cormacmonaghan@protonmail.com

Procrastination and Dementia

Dementia is a syndrome characterized initially by the progressive and typically irreversible decline of cognitive function, leading to memory loss, impaired reasoning, and difficulties with activities of daily living ([Prince et al., 2013](#); [Sanz-Blasco et al., 2022](#)). It arises from various conditions, including Alzheimer's disease, vascular dementia, and Lewy body dementia ([Cao et al., 2020](#)). The global burden of dementia is substantial, with projections suggesting that the number of affected individuals will rise from 57.4 million in 2019 to 152.8 million by 2050 ([Nichols et al., 2022](#)). Given this projection, identifying and addressing modifiable risk factors is crucial to mitigate the growing prevalence difficulties with activities of daily living of dementia worldwide. In fact, targeting these factors could prevent up to 50% of dementia cases worldwide ([Livingston et al., 2024](#)). The *Lancet* Commission highlights 14 key modifiable risks, emphasizing that dementia risk can be reduced at any age. Nevertheless, age remains the strongest non-modifiable risk factor, accounting for the largest proportion of dementia cases.

Mild cognitive impairment (MCI) is a condition characterized by cognitive changes — such as memory lapses or difficulty making decisions — that exceed typical age-related decline ([Gauthier et al., 2006](#)). Prior studies suggest that 46% of people with MCI transition to a dementia state within three years and 80% within six years ([Cooper et al., 2015](#); [Shigemizu et al., 2020](#); [Tschanz et al., 2006](#)). However, progression is neither uniform nor inevitable. Heterogeneity in outcomes underscores the importance of distinguishing between two classes of predictors: causal risk factors (e.g. smoking, hypertension, or diabetes) which may accelerate dementia onset and are potential targets for intervention ([Livingston et al., 2024](#)), and prodromal markers (e.g. memory loss, apathy, or struggles in daily living) that reflect early disease processes and may signal limited opportunities for prevention. While causal factors offer avenues for mitigating risk, prodromal features often emerge closer to irreversible pathological changes, highlighting the need for early detection and tailored interventions.

Apathy, defined as a significant loss of motivation, distinct from depression and cognitive impairment ([Fresnais et al., 2023](#); [Richard et al., 2012](#)) is among the most prevalent

neuropsychiatric symptoms in both MCI and dementia (Dalen et al., 2018; Donovan et al., 2014; Fresnais et al., 2023; Salem et al., 2023). Individuals with apathy exhibit reduced goal-directed behavior and a diminished desire to pursue rewards or pleasure (Fresnais et al., 2023).

Importantly, apathy has been identified as a significant predictor for the transition from MCI to dementia (Dalen et al., 2018; Ruthirakuhan et al., 2019; Salem et al., 2023). For instance, a meta-analysis by Dalen et al. (2018) found that apathy almost doubles the risk of transitioning to dementia. Additionally, levels of apathy measured using both the “*Neuropsychiatric Inventory Questionnaire*” (Kaufer et al., 2000) and “*Apathy Evaluation Scale*” (Marin et al., 1991) have shown to be correlated with higher levels of amyloid burden and neurofibrillary tangles in individuals with dementia, suggesting a potential connection to underlying neuropathology (Donovan et al., 2014; Marshall et al., 2013).

While apathy primarily reflects a lack of motivation, procrastination involves a delay in action despite an intention to complete such action (Steel, 2007). Although both behaviours indicate disruptions in goal-directed activity, they differ in their phenomenology: apathy is marked by diminished drive or emotional engagement, whereas procrastination typically involves intact intention but impaired follow-through. These differences suggest that procrastination may capture aspects of executive dysfunction and emotional regulation not fully accounted for by apathy alone.

Both behaviours have been linked with impaired executive function, particularly in goal-oriented behaviours, decision-making, and self-regulation (Gustavson et al., 2015; Kawagoe et al., 2017; Steel, 2010), domains which are frequently compromised in individuals with MCI and dementia (Kirova et al., 2015; Stopford et al., 2012). While apathy has been well-established as a prodromal marker and risk factor for dementia progression (Dalen et al., 2018; Ruthirakuhan et al., 2019; Salem et al., 2023), emerging evidence also suggests a potential link between procrastination and early cognitive decline. For instance, Fridén (2020) and Zhang et al. (2019) highlight that chronic procrastination is associated with dysfunction in both the dorsolateral and ventromedial prefrontal cortices, regions of the brain known to not only show early degeneration in dementia (Joseph et al., 2021), but also be linked to higher levels of apathy (Fahed & Steffens,

2021; Zhang et al., 2019).

Furthermore, procrastination may exacerbate existing cognitive decline by reinforcing patterns of inaction and passivity. Individuals who chronically delay tasks may inadvertently engage in fewer cognitively stimulating activities, such as physical activity, problem-solving, decision-making, and goal-setting. Such activities are known to build cognitive resilience and reduce dementia risk (Chowdhary et al., 2022; Livingston et al., 2024). By limiting engagement in such activities, procrastination may contribute to a self-reinforcing cycle of cognitive disengagement, potentially accelerating decline over time.

Given these behavioural and neuro-cognitive parallels with apathy, it is worth exploring whether chronic procrastination could serve as an early behavioural marker for cognitive impairment, or even a risk factor for dementia, especially in older adults. By examining procrastination alongside apathy, we may uncover a broader behavioural phenotype indicative of early neuropathological changes. Crucially, procrastination may identify individuals with motivational or executive dysfunction who do not yet meet the threshold for apathy, offering a potentially valuable and earlier behavioural marker for neurodegenerative risk.

Moreover, since age remains the strongest predictor of MCI and dementia, it is important to consider how behavioural predictors such as procrastination operate across the lifespan. Many established modifiable risk factors such as, hypertension, hearing loss, smoking, and social isolation, exert age-dependent effects, with certain factors carrying more weight at midlife than in late life (Livingston et al., 2020, 2024). In light of this, we sought to explore whether procrastination also demonstrates an age-dependent risk profile. Understanding how procrastination interacts with age may help clarify its role in the aetiology of cognitive decline and identify windows of opportunity for targeted intervention.

Although current research on procrastination in relation to dementia is non-existent, this possible association warrants exploration. Identifying procrastination as a potential risk factor could expand the scope of early interventions aimed at preventing or slowing the progression of dementia. As such, the purpose of this study was to (a) assess group differences in procrastination

levels across the normal cognition, MCI, and dementia groups, and b) test the hypotheses that higher levels of procrastination would be associated with an increased probability of transitioning from normal cognitive function or MCI to dementia.

Method

Data and study population

Analyses were conducted using a secondary data source; a multi-wave prospective cohort study called the Health and Retirement Study (HRS; [Fisher & Ryan, 2018](#)), which tracks the health, economic, and social well-being of over 18,000 American adults primarily aged > 50. The HRS is managed by the Institute for Social Research at the University of Michigan, with data collected every two years. Initial data collection of a participant is conducted through a face-to-face interview, with follow-up biennial interviews conducted either by phone or face-to-face. The average re-interview response rate ranges from 68.8% to 92.3% ([Health and Retirement Study, 2017](#)). At the time of writing, fifteen years of HRS data are currently archived.

For this study, we focused on four biennial waves of HRS data (from 2016 to 2022). Specifically our study sample consisted of respondents who participated in an experimental module assessing procrastination during the 2020 wave. These experimental modules, administered at the end of the core HRS interview, consist of concise questionnaires designed to explore new topics or supplement existing core survey data. Each respondent receives only one experimental module, with sample sizes for each module constituting approximately 10% of the core sample. As a result, while the core HRS sample includes approximately 18,000 respondents, our initial sample of interest consisted of 1,368 respondents. We excluded respondents with missing cognitive assessment data for any wave ($n = 445$), missing procrastination data ($n = 4$), and those under 50 years of age ($n = 16$). This resulted in a final analytic sample of 903 respondents.

Measures

Outcome: Cognitive Function and Cognitive Category

Cognitive function in the HRS is assessed using a series of tests adapted from the Telephone Interview for Cognitive Status (TICS; (Brandt et al., 1988; Fong et al., 2009)). These tests include an immediate and delayed 10-noun free recall test, a serial seven subtraction test, and a backward count from 20 test. Based on these assessments, Crimmins et al. (2011) developed both a 27-point cognitive scale and validated cut-off points to assess and classify cognitive status. Using these points, respondents who scored 12 – 27 were classified as having normal cognition, 7 – 11 as having MCI, and 0 – 6 as having dementia.

Predictor: Procrastination

Procrastination was measured using the Pure Procrastination Scale (Steel, 2010), which consists of 12 items rated on a Likert scale ranging from 1 (strongly disagree) to 5 (strongly agree). The total procrastination score ranges from 12 to 60, with higher scores indicating greater procrastination. The Pure Procrastination Scale conducted in 2020 (wave 3) had a Cronbach's α score of 0.91 in this sample, indicating high internal consistency. An example of a question from the scale includes "I delay making decisions until it's too late".

Confounders

To account for potential confounding, we controlled for variables with established associations with cognitive function and procrastination. These included baseline measures conducted in 2016 (wave 1) of age, sex, educational attainment, number of cardiovascular risk factors (history of hypertension, stroke, or heart disease, and a classification on "overweight" on a body mass index (BMI) scale ($BMI \geq 30$)), and depressive symptoms (Abner et al., 2012; Freedman & Cornman, 2024; Yu et al., 2013; ?). Educational attainment was classified into three categories: no formal education, GED (General Educational Development)/high school diploma, and college/further education. Depressive symptoms were measured using an eight-item version of the Center for Epidemiological Studies Depression (CES-D) scale (Briggs et al., 2018), with

scores ranging from 0 to 8, demonstrating good internal consistency ($\alpha = 0.82$).

References

- Abner, E. L., Kryscio, R. J., Cooper, G. E., Fardo, D. W., Jicha, G. A., Mendiondo, M. S., Van Eldik, L. J., Wan, L., & Schmitt, F. A. (2012). Mild cognitive impairment: Statistical models of transition using longitudinal clinical data. *International Journal of Alzheimer's Disease*, 2012(1), 291920.
- Brandt, J., Spencer, M., & Folstein, Marshal. (1988). The telephone interview for cognitive status. *Neuropsychiatry Neuropsychol Behav Neurol*, 1(2), 111–117.
- Briggs, R., Carey, D., O'Halloran, A. M., Kenny, R. A., & Kennelly, S. P. (2018). Validation of the 8-item centre for epidemiological studies depression scale in a cohort of community-dwelling older people: Data from the irish longitudinal study on ageing (TILDA). *European Geriatric Medicine*, 9, 121–126.
- Cao, Q., Tan, C.-C., Xu, W., Hu, H., Cao, X.-P., Dong, Q., & Lan Tan, and J.-T. Y. (2020). The prevalence of dementia: A systematic review and meta-analysis'. *Journal of Alzheimer's Disease*, 73(3), 1157–1166. <https://doi.org/10.3233/JAD-191092>
- Chowdhary, N., Barbui, C., Anstey, K. J., Kivipelto, M., Barbera, M., Peters, R., Zheng, L., Kulmala, J., Stephen, R., Ferri, C. P., Joannette, Y., Wang, H., Comas-Herrera, A., & Alessi, C. S. (2022). (Dy), k., mwangi. K. J., Petersen, R. C., Motala, A. A., Mendis, S., ... Dua, T. *Reducing the Risk of Cognitive Decline and Dementia: WHO Recommendations. Frontiers in Neurology*, 12. <https://doi.org/10.3389/fneur.2021.765584>
- Cooper, C., Sommerlad, A., Lyketsos, C. G., & Livingston, G. (2015). Modifiable predictors of dementia in mild cognitive impairment: A systematic review and meta-analysis. *American Journal of Psychiatry*, 172(4), 323–334.
- Crimmins, E. M., Kim, J. K., Langa, K. M., & Weir, D. R. (2011). Assessment of cognition using surveys and neuropsychological assessment: The health and retirement study and the aging, demographics, and memory study. *Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, 66(suppl_1), i162–i171.

- Dalen, J. W. van, Wanrooij, L. L. van, Charante, E. P. M. van, Brayne, C., Gool, W. A. van, & Richard, E. (2018). Association of apathy with risk of incident dementia: A systematic review and meta-analysis. *JAMA Psychiatry*, 75(10), 1012–1021.
- Donovan, N. J., Wadsworth, L. P., Lorus, N., Locascio, J. J., Rentz, D. M., Johnson, K. A., Sperling, R. A., Marshall, G. A., Initiative, A. D. N., et al. (2014). Regional cortical thinning predicts worsening apathy and hallucinations across the alzheimer disease spectrum. *The American Journal of Geriatric Psychiatry*, 22(11), 1168–1179.
- Fahed, M., & Steffens, D. C. (2021). Apathy: Neurobiology, assessment and treatment. *Clinical Psychopharmacology and Neuroscience*, 19(2), 181.
- Fisher, G. G., & Ryan, L. H. (2018). Overview of the health and retirement study and introduction to the special issue. *Work, Aging and Retirement*, 4(1), 1–9.
- Fong, T. G., Fearing, M. A., Jones, R. N., Shi, P., Marcantonio, E. R., Rudolph, J. L., Yang, F. M., Kiely, D. K., & Inouye, S. K. (2009). Telephone interview for cognitive status: Creating a crosswalk with the mini-mental state examination. *Alzheimer's & Dementia*, 5(6), 492–497.
- Freedman, V. A., & Cornman, J. C. (2024). Dementia prevalence, incidence, and mortality trends among US adults ages 72 and older, 2011–2021. *The Journals of Gerontology, Series A: Biological Sciences and Medical Sciences*, 79(Supplement_1), S22–S31.
- Fresnais, D., Humble, M. B., Bejerot, S., Meehan, A. D., & Fure, B. (2023). Apathy as a predictor for conversion from mild cognitive impairment to dementia: A systematic review and meta-analysis of longitudinal studies. *Journal of Geriatric Psychiatry and Neurology*, 36(1), 3–17.
- Fridén, I. (2020). *Procrastination as a form of self-regulation failure: A review of the cognitive and neural underpinnings*.
- Gauthier, S., Reisberg, B., Zaudig, M., Petersen, R. C., Ritchie, K., Broich, K., Belleville, S., Brodaty, H., Bennett, D., Chertkow, H., Cummings, J. L., de Leon, M., Feldman, H., Ganguli, M., Hampel, H., Scheltens, P., Tierney, M. C., Whitehouse, P., & Winblad, B. (2006). Mild cognitive impairment. *The Lancet*, 367(9518), 1262–1270.

[https://doi.org/10.1016/S0140-6736\(06\)68542-5](https://doi.org/10.1016/S0140-6736(06)68542-5)

Gustavson, D. E., Miyake, A., Hewitt, J. K., & Friedman, N. P. (2015). Understanding the cognitive and genetic underpinnings of procrastination: Evidence for shared genetic influences with goal management and executive function abilities. *Journal of Experimental Psychology: General*, 144(6), 1063.

Health and Retirement Study. (2017). *Sample sizes and response rates*.

https://hrs.isr.umich.edu/sites/default/files/biblio/ResponseRates_2017.pdf

Joseph, S., Knezevic, D., Zomorodi, R., Blumberger, D. M., Daskalakis, Z. J., Mulsant, B. H., Pollock, B. G., Voineskos, A., Wang, W., Rajji, T. K., et al. (2021). Dorsolateral prefrontal cortex excitability abnormalities in alzheimer's dementia: Findings from transcranial magnetic stimulation and electroencephalography study. *International Journal of Psychophysiology*, 169, 55–62.

Kaufer, D. I., Cummings, J. L., Ketchel, P., Smith, V., MacMillan, A., Shelley, T., Lopez, O. L., & DeKosky, S. T. (2000). Validation of the NPI-q, a brief clinical form of the neuropsychiatric inventory. *The Journal of Neuropsychiatry and Clinical Neurosciences*, 12(2), 233–239.

Kawagoe, T., Onoda, K., & Yamaguchi, S. (2017). Apathy and executive function in healthy elderly—resting state fMRI study. *Frontiers in Aging Neuroscience*, 9, 124.

Kirova, A. M., Bays, R. B., & Lagalwar, S. (2015). Working memory and executive function decline across normal aging, mild cognitive impairment, and alzheimer's disease. *BioMed Research International*, 2015(1), 748212.

Livingston, G., Huntley, J., Liu, K. Y., Costafreda, S. G., Selbæk, G., Alladi, S., Ames, D., Banerjee, S., Burns, A., Brayne, C., Fox, N. C., Ferri, C. P., Gitlin, L. N., Howard, R., Kales, H. C., Kivimäki, M., Larson, E. B., Nakasujja, N., Rockwood, K., ... Mukadam, N. (2024). Dementia prevention, intervention, and care: 2024 report of the Lancet standing Commission. *The Lancet*, 404(10452), 572–628. [https://doi.org/10.1016/S0140-6736\(24\)01296-0](https://doi.org/10.1016/S0140-6736(24)01296-0)

Livingston, G., Huntley, J., Sommerlad, A., Ames, D., Ballard, C., Banerjee, S., Brayne, C., Burns, A., Cohen-Mansfield, J., Cooper, C., et al. (2020). Dementia prevention, intervention,

- and care: 2020 report of the lancet commission. *The Lancet*, 396(10248), 413–446.
- Marin, R. S., Biedrzycki, R. C., & Firinciogullari, S. (1991). Reliability and validity of the apathy evaluation scale. *Psychiatry Research*, 38(2), 143–162.
- Marshall, G. A., Donovan, N. J., Lorus, N., Gidyczin, C. M., Maye, J., Pepin, L. C., Becker, J. A., Amariglio, R. E., Rentz, D. M., Sperling, R. A., et al. (2013). Apathy is associated with increased amyloid burden in mild cognitive impairment. *The Journal of Neuropsychiatry and Clinical Neurosciences*, 25(4), 302–307.
- Nichols, E., Steinmetz, J. D., Vollset, S. E., Fukutaki, K., Chalek, J., Abd-Allah, F., Abdoli, A., Abualhasan, A., Abu-Gharbieh, E., Akram, T. T., et al. (2022). Estimation of the global prevalence of dementia in 2019 and forecasted prevalence in 2050: An analysis for the global burden of disease study 2019. *The Lancet Public Health*, 7(2), e105–e125.
- Prince, M., Bryce, R., Albanese, E., Wimo, A., Ribeiro, W., & Ferri, C. P. (2013). The global prevalence of dementia: A systematic review and meta analysis. *Alzheimer's & Dementia*, 9(1), 63–75. <https://doi.org/10.1016/j.jalz.2012.11.007>
- Richard, E., Schmand, B., Eikelenboom, P., Yang, S. C., Ligthart, S. A., Moll van Charante, E. P., Gool, W. A. van, & Initiative, A. D. N. (2012). Symptoms of apathy are associated with progression from mild cognitive impairment to alzheimer's disease in non-depressed subjects. *Dementia and Geriatric Cognitive Disorders*, 33(2-3), 204–209.
- Ruthirakuhan, M., Herrmann, N., Vieira, D., Gallagher, D., & Lanctôt, K. L. (2019). The roles of apathy and depression in predicting alzheimer disease: A longitudinal analysis in older adults with mild cognitive impairment. *The American Journal of Geriatric Psychiatry*, 27(8), 873–882.
- Salem, H., Suchting, R., Gonzales, M. M., Seshadri, S., & Teixeira, A. L. (2023). Apathy as a predictor of conversion from mild cognitive impairment to alzheimer's disease: A texas alzheimer's research and care consortium (TARCC) cohort-based analysis. *Journal of Alzheimer's Disease*, 92(1), 129–139.
- Sanz-Blasco, R., Ruiz-Sánchez de León, J. M., Ávila-Villanueva, M., Valentí-Soler, M.,

- Gómez-Ramírez, J., & Fernández-Blázquez, M. A. (2022). Transition from mild cognitive impairment to normal cognition: Determining the predictors of reversion with multi-state Markov models. *Alzheimer's & Dementia*, 18(6), 1177–1185.
<https://doi.org/10.1002/alz.12448>
- Shigemizu, D., Akiyama, S., Higaki, S., Sugimoto, T., Sakurai, T., Boroevich, K. A., Sharma, A., Tsunoda, T., Ochiya, T., Niida, S., & Ozaki, K. (2020). Prognosis prediction model for conversion from mild cognitive impairment to alzheimer's disease created by integrative analysis of multi-omics data. *Alzheimer's Research & Therapy*, 12*, 1–12.
- Steel, P. (2007). The nature of procrastination: A meta-analytic and theoretical review of quintessential self-regulatory failure. *Psychological Bulletin*, 3(1).
- Steel, P. (2010). Arousal, avoidant and decisional procrastinators: Do they exist? *Personality and Individual Differences*, 48(8), 926–934.
- Stopford, C. L., Thompson, J. C., Neary, D., Richardson, A. M., & Snowden, J. S. (2012). Working memory, attention, and executive function in alzheimer's disease and frontotemporal dementia. *Cortex*, 48(4), 429–446.
- Tschanz, J. T., Welsh-Bohmer, K. A., Lyketsos, C. G., Corcoran, C., Green, R. C., Hayden, K., Norton, M. C., Zandi, P. P., Toone, L., West, N. A., & Breitner, J. C. S. (2006). And the cache county investigators. *Conversion to Dementia from Mild Cognitive Disorder: The Cache County Study*. **Neurology*, 67(2), 229–234.
- Yu, H. M., Yang, S. S., Gao, J. W., Zhou, L. Y., Liang, R. F., & Qu, C. Y. (2013). Multi-state Markov model in outcome of mild cognitive impairments among community elderly residents in mainland China. *International Psychogeriatrics*, 25(5), 797–804.
- Zhang, S., Liu, P., & Feng, T. (2019). To do it now or later: The cognitive mechanisms and neural substrates underlying procrastination. *Wiley Interdisciplinary Reviews: Cognitive Science*, 10(4), e1492.