

Sobol' Matrices for Multi-Output Gaussian Processes*

Robert A. Milton[†] and Solomon F. Brown[‡]

Abstract. Variance based global sensitivity analysis measures the relevance of inputs to a single output using Sobol' indices. A recent paper [R. A. Milton, A. S. Yearley, and S. F. Brown, *Submitted to SIAM/ASA Journal on Uncertainty Quantification*, 2024] extended the definition in a natural way to multiple outputs, directly measuring the relevance of inputs to the linkages between outputs in a correlation-like matrix of indices. Sobol' matrices and their standard errors were related to the moments of the multi-output model, to enable calculation. The present work carries through the explicit, analytic calculation of Sobol' matrices for the widely applicable, concrete case of multi-output Gaussian processes with an anisotropic radial basis function kernel. The calculations are extensively benchmarked numerically against test functions (with added noise), whose Sobol' matrices are calculated analytically.

Key words. Global Sensitivity Analysis, Sobol' Index, Surrogate Model, Multi-Output, Gaussian Process, Uncertainty Quantification

MSC codes. 60G07, 60G15, 62J10, 62H99

1. Introduction. This paper is concerned with analyzing the results of experiments or computer simulations in a design matrix of $M \geq 1$ input axes (columns) and $L \geq 1$ output axes (columns) over N samples (rows). Global Sensitivity Analysis (GSA) [24] examines the relevance of the various inputs to the various outputs. When pursued via ANOVA decomposition of a single output, this leads naturally to the well known Sobol' indices, which have by now been applied across most fields of science and engineering [26, 14]. A recent paper [21] extended the definition in a natural way to multiple outputs $L \geq 1$. This work implements the extended definition for the widely applicable, concrete case of Multi-Output Gaussian Processes (MOGPs) with an anisotropic Radial Basis Function (RBF) kernel.

Sobol' indices apportion the variance of a scalar output to the influence of various inputs. There is no generally accepted extension of Sobol' indices to multi-outputs $L > 1$. The Sobol' matrices developed in [21] apportion the correlation matrix of outputs – rather than the variance of a single output – to the influence of various inputs. The diagonal elements are identically the classical Sobol' indices, expressing the relevance of inputs to the output variables themselves. The off-diagonal elements express the relevance of inputs to the correlation between outputs. This may be of vital interest when outputs are, for example, yield and purity of a product, cost and efficacy of a process, or perhaps a single output measured at various times. The Sobol matrices reveal (amongst other things) which inputs it is worthwhile varying in an effort to alter the linkage between outputs.

*Submitted to the editors DATE.

Funding: This work was funded by the EPSRC under grant reference EP/V051458/1.

[†]Department of Chemical, Materials and Biological Engineering, University of Sheffield, Sheffield, S1 3JD, United Kingdom (r.a.milton@sheffield.ac.uk, <https://www.browngroupsheffield.com/>).

[‡]Department of Chemical, Materials and Biological Engineering, University of Sheffield, Sheffield, S1 3JD, United Kingdom (s.f.brown@sheffield.ac.uk, <https://www.browngroupsheffield.com/>).

Prior work on Sobol' indices with multiple outputs [12, 13, 30, 9] has settled ultimately on just the diagonal elements of the covariance matrix, so covariance between outputs remains unexamined. Previous studies have performed principal component analysis (PCA) on outputs prior to performing GSA on the resulting principal component outputs [7]. This has been used in particular to study synthetic multi-outputs which are actually the dynamic response of a single output over time [17, 34]. Hierarchical sensitivity analysis [33] employs prior mapping to independent subsystems. In all these cases, output covariance is essentially removed prior to GSA. Only very recently has GSA attempted to incorporate output covariance at all [19], as an ingredient in a total fluctuation. The Sobol' matrices [21] pursued in the present work constitute the first GSA performed directly on output covariance, in order to examine the influence of various inputs on the correlation between outputs.

Accurate calculation of Sobol' indices for a single output is computationally expensive and may require 10,000+ samples [18, 22]. A (potentially) more efficient approach is calculation via a surrogate model, such as a Gaussian Process (GP) [25, 23], Polynomial Chaos Expansion (PCE) [15, 32, 31], low-rank tensor approximation [10, 16], or support vector regression [11, 9]. Semi-analytic expressions for Sobol' indices are available for scalar PCEs [28], and the diagonal elements of multi-output PCEs [13]. Semi-analytic expressions for Sobol' indices of GPs have been provided in integral form by [22] and alternatively by [8]. These approaches are implemented, examined and compared in [20, 27]. Both [22, 20] estimate the errors on Sobol' indices in semi-analytic, integral form. Fully analytic, closed form expressions have been derived without error estimates for uniformly distributed inputs [29] using a GP with an RBF kernel. The present work performs the calculation of full Sobol' matrices and their standard errors for MOGPs with normally distributed inputs. This enhances analytical elegance and computational efficiency, and allows for rotation of the input basis. After all, an arbitrarily rotated multivariate normal distribution is still a multivariate normal distribution, whereas a multivariate uniform distribution can only be rotated by right angles if it is to remain uniform.

When data is limited, some form of surrogate is really indispensable to perform GSA. Accordingly Sobol' matrices and their error estimates are related to the moments of a surrogate model in [21]. From this starting point, the present work develops explicit, analytic formulae for Sobol' matrices and their standard errors from MOGP moments. An anisotropic RBF kernel is assumed, appropriate to outputs which vary smoothly over continuous inputs. The analytic formulae have implemented as an open source Python library [4]. This allows us to benchmark implementation against test functions with known Sobol' matrices. This is important, because practitioners are rarely privileged with an exact and reliable exogenous model linking outputs to inputs. Rather, GSA is a preliminary stage in reducing empirical observations to a tractable, analyzable representation. For example, it is frequently used to guide model reduction and the design of experiments, as well as or as part of optimization and risk assessment. Under these circumstances, the GSA itself must be robust and reliable, providing well benchmarked measures of its own uncertainty and domain of validity.

The contents of this paper are as follows. introduces the notation and context for performing GSA in [21].

2. Multi-output models with quantified uncertainty (MQUs). The purpose of this Section is to set the scene for this study and serve as a glossary of notation. The notation is not

quite standard, but is preferred for lightness and fluency once grasped. Regarding notation and other topics, the intention is to facilitate computation by combining efficient, tensorized operations such as `einsum` [1, 2, 3] into numerically stable calculations. As such, the devices employed herein should be familiar to practitioners of machine learning.

The preference throughout this paper is for uppercase constants and lowercase variables. Square bracketed quantities such as $[u]_{\mathbf{M} \times \mathbf{N}}$ are tensors subscripted by (the cartesian product of) their ranks, for bookkeeping. Each rank is expressed as an ordinal of axes, a tuple which is conveniently also a naive set and may be subtracted from others as such

$$\mathbf{0} := () \subseteq \mathbf{m} := (0, \dots, m-1) \subseteq \mathbf{M} := (0, \dots, M-1)$$

$$\mathbf{M} - \mathbf{m} := \{m' \in \mathbf{M} \mid m' \notin \mathbf{m}\} = (m, \dots, M-1)$$

GSA decomposes input space by families of axes, and this notation facilitates that. It does assume that input axes are expediently ordered already, which might impair convenience but not generality. Conventional indexing of components is permitted under this scheme, by allowing singleton axes to be written in italic without parentheses.

From a Bayesian perspective a determined tensor is a function of determined input(s), a random variable (RV) a function of undetermined input, and a stochastic process (SP) a function of both determined and undetermined inputs. In any case, all sources of (sans-serif) uncertainty may be gathered in a single input $[u]_M$, final to the definition of input space:

$$(2.1) \quad \begin{array}{ll} \mathbf{M} := (0, \dots, M-1) & \text{determined inputs on the unit interval} \quad [u]_{\mathbf{M}} \in [0, 1]^M \\ M =: \mathbf{M} + \mathbf{1} - \mathbf{M} & \text{an undetermined input on the unit interval} \quad [u]_M \sim \mathcal{U}(0, 1) \end{array}$$

Exponentiation is always categorical – repeated cartesian $\times$ or tensor $\otimes$ – unless stated otherwise. It is crucial to this work that all $\mathbf{M} + \mathbf{1}$ inputs vary completely independently of each other – they are in no way codependent or correlated. Uncertainty is distributed uniformly $[u]_M \sim \mathcal{U}(0, 1)$ in [21] to exploit the “universality of the uniform” [6, pp.224], better known as the inverse probability integral transform [5]. Let us take immediate advantage by defining

$$(2.2) \quad [z]_{\mathbf{M}+\mathbf{1}} := P^{-1}([u]_{\mathbf{M}+\mathbf{1}} \mid [0]_{\mathbf{M}+\mathbf{1}}, \langle 1 \rangle_{\mathbf{M}+\mathbf{1}^2}) \sim \mathcal{N}([0]_{\mathbf{M}+\mathbf{1}}, \langle 1 \rangle_{\mathbf{M}+\mathbf{1}^2})$$

where $\mathcal{N}([0]_{\mathbf{M}+\mathbf{1}}, \langle 1 \rangle_{\mathbf{M}+\mathbf{1}^2})$ is the $\mathbf{M} + \mathbf{1}$ dimensional standard normal distribution with mean $[0]_{\mathbf{M}+\mathbf{1}}$ and variance $\langle 1 \rangle_{\mathbf{M}+\mathbf{1}^2}$, and $P^{-1}([u]_{\mathbf{M}+\mathbf{1}} \mid [0]_{\mathbf{M}+\mathbf{1}}, \langle 1 \rangle_{\mathbf{M}+\mathbf{1}^2})$ is its quantile function, or inverse CDF. Henceforth, angle brackets always denote a diagonal tensor, identically 0 off the leading diagonal of the final two axes. The advantage of normally

$$(2.3) \quad [z]_{\mathbf{M}+\mathbf{1}} := P^{-1}([u]_{\mathbf{M}+\mathbf{1}} \mid [0]_{\mathbf{M}+\mathbf{1}}, \langle 1 \rangle_{\mathbf{M}+\mathbf{1}^2}) \sim \mathcal{N}([0]_{\mathbf{M}+\mathbf{1}}, \langle 1 \rangle_{\mathbf{M}+\mathbf{1}^2})$$

Throughout this work, expectations are subscripted by the input axes marginalized

$$\mathbb{E}_{\#}[\bullet] := \int_{[0]_{\#}}^{[1]_{\#}} [\bullet] \, d[u]_{\#} \quad \forall \# \subseteq \mathbf{M} + \mathbf{1} \text{ or } \# \in \mathbf{M} + \mathbf{1}$$

Covariances invoke the tensor product (summing the ranks of the arguments), and carry the subscript of their underlying expectation

$$\mathbb{V}_{\#}[[\bullet], [*]] := \mathbb{E}_{\#}[[\bullet] \otimes [*]] - \mathbb{E}_{\#}[\bullet] \otimes \mathbb{E}_{\#}[*] \quad \forall \# \subseteq \mathbf{M} + \mathbf{1} \text{ or } \# \in \mathbf{M} + \mathbf{1}$$

The covariance of anything with itself is expressed with a single argument $\mathbb{V}_{\#}[\bullet] := \mathbb{V}_{\#}[\bullet, \bullet]$, as is customary.

3. Sobol' matrices.

4. GP regression.

5. GP moments.

6. GP estimates.

7. Mitigating computational cost.

8. Test functions.

9. Benchmarking.

10. Conclusions.

REFERENCES

- [1] *NumPy*, 2023-08-19, <https://numpy.org>.
- [2] *PyTorch*, 2023-08-19, <https://pytorch.org>.
- [3] *TensorFlow*, 2023-08-19, <https://tensorflow.org>.
- [4] *rom-comma*, 2024-03-01, <https://github.com/C-O-M-M-A/rom-comma>.
- [5] J. E. ANGUS, *The probability integral transform and related results*, SIAM Review, 36 (1994), pp. 652–654, <https://doi.org/10.1137/1036146>.
- [6] J. K. BLITZSTEIN AND J. HWANG, *Introduction to Probability, Second Edition*, Taylor & Francis Group, 2019.
- [7] K. CAMPBELL, M. D. MCKAY, AND B. J. WILLIAMS, *Sensitivity analysis when model outputs are functions*, Reliability Engineering & System Safety, 91 (2006), pp. 1468–1472, <https://doi.org/10.1016/j.res.2005.11.049>.
- [8] W. CHEN, R. JIN, AND A. SUDJANTO, *Analytical variance-based global sensitivity analysis in simulation-based design under uncertainty*, Journal of Mechanical Design, 127 (2005), p. 875, <https://doi.org/10.1115/1.1904642>.
- [9] K. CHENG, Z. LU, AND K. ZHANG, *Multivariate output global sensitivity analysis using multi-output support vector regression*, Structural and Multidisciplinary Optimization, 59 (2019), pp. 2177–2187, <https://doi.org/10.1007/s00158-018-2184-z>.
- [10] M. CHEVREUIL, R. LEBRUN, A. NOUY, AND P. RAI, *A least-squares method for sparse low rank approximation of multivariate functions*, SIAM/ASA Journal on Uncertainty Quantification, 3 (2015), pp. 897–921, <https://doi.org/10.1137/13091899X>, <https://arxiv.org/abs/http://arxiv.org/abs/1305.0030v2>.
- [11] C. CORTES AND V. VAPNIK, *Support-vector networks*, Machine Learning, 20 (1995), pp. 273–297, <https://doi.org/10.1007/bf00994018>.
- [12] F. GAMBOA, A. JANON, T. KLEIN, AND A. LAGNOUX, *Sensitivity indices for multivariate outputs*, Comptes Rendus Mathématique, 351 (2013), pp. 307–310, <https://doi.org/10.1016/j.crma.2013.04.016>.
- [13] O. GARCIA-CABREJO AND A. VALOCCHI, *Global sensitivity analysis for multivariate output using polynomial chaos expansion*, Reliability Engineering & System Safety, 126 (2014), pp. 25–36, <https://doi.org/10.1016/j.res.2014.01.005>.
- [14] R. GHANEM, D. HIGDON, H. OWHADI, ET AL., *Handbook of uncertainty quantification*, vol. 6, Springer, 2017.
- [15] R. G. GHANEM AND P. D. SPANOS, *Spectral techniques for stochastic finite elements*, Archives of Computational Methods in Engineering, 4 (1997), pp. 63–100, <https://doi.org/10.1007/BF02818931>.

- [16] K. KONAKLI AND B. SUDRET, *Global sensitivity analysis using low-rank tensor approximations*, Reliability Engineering & System Safety, 156 (2016), pp. 64–83, <https://doi.org/10.1016/j.ress.2016.07.012>.
- [17] M. LAMBONI, H. MONOD, AND D. MAKOWSKI, *Multivariate sensitivity analysis to measure global contribution of input factors in dynamic models*, Reliability Engineering & System Safety, 96 (2011), pp. 450–459, <https://doi.org/10.1016/j.ress.2010.12.002>.
- [18] B. LAMOUREUX, N. MECBAL, AND J. R. MASSÉ, *A combined sensitivity analysis and kriging surrogate modeling for early validation of health indicators*, Reliability Engineering and System Safety, 130 (2014), pp. 12–26, <https://doi.org/10.1016/j.ress.2014.03.007>.
- [19] Q. LIU, N. TONG, AND X. HAN, *A global sensitivity analysis method for multi-input multi-output system and its application in structural design*, International Journal of Computational Methods, 19 (2021), <https://doi.org/10.1142/s021987622141005x>.
- [20] A. MARREL, B. IOOSS, B. LAURENT, AND O. ROUSTANT, *Calculations of sobol indices for the gaussian process metamodel*, Reliability Engineering & System Safety, 94 (2009), pp. 742–751, <https://doi.org/10.1016/j.ress.2008.07.008>.
- [21] R. A. MILTON, A. S. YEARDLEY, AND S. F. BROWN, *Submitted to SIAM/ASA Journal on Uncertainty Quantification*, 2024.
- [22] J. E. OAKLEY AND A. O'HAGAN, *Probabilistic sensitivity analysis of complex models: a bayesian approach*, Journal of the Royal Statistical Society: Series B (Statistical Methodology), 66 (2004), pp. 751–769, <https://doi.org/10.1111/j.1467-9868.2004.05304.x>.
- [23] C. E. RASMUSSEN AND C. K. I. WILLIAMS, *Gaussian Processes for Machine Learning (Adaptive Computation and Machine Learning series)*, The MIT Press, 2005.
- [24] S. RAZAVI, A. JAKEMAN, A. SALTELLI, C. PRIEUR, B. IOOSS, E. BORGONOVO, E. PLISCHKE, S. L. PIANO, T. IWANAGA, W. BECKER, S. TARANTOLA, J. H. GUILLAUME, J. JAKEMAN, H. GUPTA, N. MELILLO, G. RABITTI, V. CHABRIDON, Q. DUAN, X. SUN, S. SMITH, R. SHEIKHOESLAMI, N. HOSSEINI, M. ASADZADEH, A. PUY, S. KUCHERENKO, AND H. R. MAIER, *The future of sensitivity analysis: An essential discipline for systems modeling and policy support*, Environmental Modelling & Software, 137 (2021-03), p. 104954, <https://doi.org/10.1016/j.envsoft.2020.104954>.
- [25] J. SACKS, W. J. WELCH, T. J. MITCHELL, AND H. P. WYNN, *Design and analysis of computer experiments*, Statistical Science, 4 (1989), pp. 409–423.
- [26] A. SALTELLI, K. ALEKSANKINA, W. BECKER, P. FENNELL, F. FERRETTI, N. HOLST, S. LI, AND Q. WU, *Why so many published sensitivity analyses are false: A systematic review of sensitivity analysis practices*, Environmental Modelling & Software, 114 (2019-04), pp. 29–39, <https://doi.org/10.1016/j.envsoft.2019.01.012>.
- [27] A. SRIVASTAVA, A. K. SUBRAMANIYAN, AND L. WANG, *Analytical global sensitivity analysis with gaussian processes*, Artificial Intelligence for Engineering Design, Analysis and Manufacturing, 31 (2017), pp. 235–250, <https://doi.org/10.1017/s0890060417000142>.
- [28] B. SUDRET, *Global sensitivity analysis using polynomial chaos expansions*, Reliability Engineering & System Safety, 93 (2008), pp. 964–979, <https://doi.org/10.1016/j.ress.2007.04.002>.
- [29] Z. WU, D. WANG, P. O. N, F. HU, AND W. ZHANG, *Global sensitivity analysis using a gaussian radial basis function metamodel*, Reliability Engineering & System Safety, 154 (2016), pp. 171–179, <https://doi.org/10.1016/j.ress.2016.06.006>.
- [30] S. XIAO, Z. LU, AND F. QIN, *Estimation of the generalized sobol's sensitivity index for multivariate output model using unscented transformation*, Journal of Structural Engineering, 143 (2017), [https://doi.org/10.1061/\(asce\)st.1943-541x.0001721](https://doi.org/10.1061/(asce)st.1943-541x.0001721).
- [31] D. XIU, *Numerical Methods for Stochastic Computations: A Spectral Method Approach*, Princeton University Press, 2010.
- [32] D. XIU AND G. E. KARNIADAKIS, *The wiener-askey polynomial chaos for stochastic differential equations*, SIAM Journal on Scientific Computing, 24 (2002), pp. 619–644, <https://doi.org/10.1137/s1064827501387826>.
- [33] C. XU, P. ZHU, Z. LIU, AND W. TAO, *Mapping-based hierarchical sensitivity analysis for multilevel systems with multidimensional correlations*, Journal of Mechanical Design, 143 (2020), <https://doi.org/10.1115/1.4047689>.
- [34] K. ZHANG, Z. LU, K. CHENG, L. WANG, AND Y. GUO, *Global sensitivity analysis for multivariate output model and dynamic models*, Reliability Engineering & System Safety, 204 (2020), p. 107195,

211 <https://doi.org/10.1016/j.res.2020.107195>.