

The networks they are a-changin' - Investigating auditory system connectivities during cortical processing of voice, song and music in left-handers and musicians

A. Introduction

1. structures of auditory cortex

- only short talk on non-cortical auditory areas ("ear-structures" and maybe brainstem)
- primary and secondary areas within temporal lobe:
 - Heschel's gyrus: primary auditory cortex and its tonotopic fields (similarities according to cochlea-functions)
(Humphries, Liebenthal, & Binder, 2010), (Da Costa et al., 2011)
 - areas around HG: superior temporal gyrus (STG) with localisations of planum polare (PP) and planum temporale (PT): further labelled as secondary or non-primary auditory cortex
 - "higher processing": less tonotopic and more spectrotemporal representations of sound spectrograms with different degrees of spectral and temporal resolution
(Santoro et al., 2014)

2. processing speech singing and music - differences and similarities

- ongoing discussion: separated neuronal processing and overlaps
(Tervaniemi, 2006) (Norman-Haignere, Kanwisher, & McDermott, 2015)
(Rogalsky, Rong, Saberi, & Hickok, 2011)

a) neuronal processing of speech

- accurate processing requires fine-graded temporal resolution
- less important: spectral sampling (only sparse)
- proposed left-lateralisation of speech-processing
- pSTG/PT as mentioned neuronal correlates in auditory cortex
- term voice and not speech, because we are not interested in higher order processing like prosody,...etc.

b) neuronal processing of music

- short outlining regarding to music-properties: F0, harmonics and pitch
- vice versa to speech: finegraded spectral resolution is more important in music than temporal resolution

- proposed right-lateralisation of music-processing
- aSTG/PP as mentioned neuronal correlates in AC

c) singing and its properties

- properties combine both aspects mentioned in music- and speech processing
(Schön, Gordon, & Besson, 2005), (Kleber, Birbaumer, Veit, Trevorrow, & Lotze, 2007)

d) choice of stimuli - a gradient of musicality

- 3 stimuli music, song and voice generate a gradient according to spectrotemporal properties
- differences of music and voice and connection via song: sufficient to visualize a more general neuronal system in context of auditory stimuli-processing?

3. influence of handedness and musical training

a) musical training

- induced neuroplasticity according to musical training and its influence on processing speech and music

(Angulo-Perkins et al., 2014; Elmer, Hänggi, Meyer, & Jäncke, 2013; Meyer, Elmer, & Jäncke, 2012; Schlaug, Jäncke, Huang, & Steinmetz, 1995)

b) handedness

- short introduction according to handedness in general
- left-handers are often mentioned in context of lateralisation, but also often excluded in cognitive neuroscience due to varying neuronal processing strategies compared to right-handers
(Willems, Der Haegen, Fisher, & Francks, 2014)
- by including left-handers it is aimed to get a more reliable insight in neuronal auditory processing from an other "point of view"
- additional value by including left-handers referred to lateralisation and mentioned occurrence in music-, speech-, temporal-, spectral processing

4. aim of the study

- Neuronal connectivity differences within primary and secondary auditory cortex during the processing of voice, song and music shall be investigated; according to this findings the leading question will be further expanded to the influence of handedness and musical training on inter- and intrahemispheric connectivities

- To visualize the information flow of neuronal correlates in an optimal manner of spatial and temporal resolution, combined measurements with fMRI and EEG will be implemented
- ROI's: bilateral HG and STG (PP, PT)
- hypotheses:
 - 1. How does the application of the mentioned "gradient of musicality", manipulated through presentation of the 3 different auditory stimuli, alter connectivities between areas of auditory cortex?**
 - 2. Do left-handers, respectively musicians show other models of neuronal connectivity than right-handed people without any musical training?**

B. Material and Methods

1. participants

- 3 groups recruited: right-handed non-musicians (control group), right-handed musicians, left-handed non-musicians
- definition of "musician" (years of practice, weekly training)
- validation of experimental groups: musical ear test (MET), individual screening formular (reference to appendix), evaluation of handedness (oldfield)
(Wallentin, Nielsen, Friis-Olivarius, Vuust, & Vuust, 2010)
- amusia-test (MBEA)
(Peretz, Champod, & Hyde, 2003)

2. stimuli and experimental procedure

- music- and voice-stimuli from Angulo-Perkins et al. (2014; more details: ethic proposal)
- EEG-fMRI combined measurement
- 1 EKG-elektrode for measurement of physiological parameters

3. preprocessing and statistical analysis

- preprocessing as usual for fMRI-data and in addition, if application in python possible, EEG-assisted motion correction (usage of preprocessing pipeline in python)
(Caballero-Gaudes & Reynolds, 2017; Wong et al., 2016; Zotev, Yuan, Phillips, & Bodurka, 2012)
- GLM, DCM

(Friston, Harrison, & Penny, 2003), (Friston, Holmes, Worsley, Poline, & Frackowiak, 1995)

- adding physiological measurements as nuisance regressors to GLM (if possible)

C. Results

D. Discussion

Appendix

References

- Angulo-Perkins, A., Aubé, W., Peretz, I., Barrios, F. A., Armony, J. L., & Concha, L. (2014). Music listening engages specific cortical regions within the temporal lobes: Differences between musicians and non-musicians. *Cortex*, 59, 126–137. <https://doi.org/10.1016/j.cortex.2014.07.013>
- Caballero-Gaudes, C., & Reynolds, R. C. (2017). Methods for cleaning the BOLD fMRI signal. *NeuroImage*, 154(December 2016), 128–149. <https://doi.org/10.1016/j.neuroimage.2016.12.018>
- Da Costa, S., van der Zwaag, W., Marques, J. P., Frackowiak, R. S. J., Clarke, S., & Saenz, M. (2011). Human Primary Auditory Cortex Follows the Shape of Heschl's Gyrus. *Journal of Neuroscience*, 31(40), 14067–14075. <https://doi.org/10.1523/JNEUROSCI.2000-11.2011>
- Elmer, S., Hänggi, J., Meyer, M., & Jäncke, L. (2013). Increased cortical surface area of the left planum temporale in musicians facilitates the categorization of phonetic and temporal speech sounds. *Cortex*, 49(10), 2812–2821. <https://doi.org/10.1016/j.cortex.2013.03.007>
- Friston, K. J., Harrison, L., & Penny, W. (2003). Dynamic causal modelling. *NeuroImage*, 19(4), 1273–1302. [https://doi.org/10.1016/S1053-8119\(03\)00202-7](https://doi.org/10.1016/S1053-8119(03)00202-7)
- Friston, K. J., Holmes, A. P., Worsley, K. J., Poline, J., & Frackowiak, R. S. J. (1995). b. Statistical parametric maps in functional imaging: A general linear approach. *Human Brain Mapping*, 2, 189–210.
- Humphries, C., Liebenthal, E., & Binder, J. R. (2010). Tonotopic organization of human auditory cortex. *NeuroImage*, 50(3), 1202–1211. <https://doi.org/10.1016/j.neuroimage.2010.01.046>
- Kleber, B., Birbaumer, N., Veit, R., Trevorrow, T., & Lotze, M. (2007). Overt and imagined singing of an Italian aria. *NeuroImage*, 36(3), 889–900. <https://doi.org/10.1016/j.neuroimage.2007.02.053>
- Meyer, M., Elmer, S., & Jäncke, L. (2012). Musical expertise induces neuroplasticity of the planum temporale. *Annals of the New York Academy of Sciences*, 1252(1), 116–123. <https://doi.org/10.1111/j.1749-6632.2012.06450.x>
- Norman-Haignere, S., Kanwisher, N. G., & McDermott, J. H. (2015). Distinct Cortical Pathways for Music and Speech Revealed by Hypothesis-Free Voxel Decomposition. *Neuron*, 88(6), 1281–1296. <https://doi.org/10.1016/j.neuron.2015.11.035>

- Peretz, I., Champod, A. S., & Hyde, K. (2003). Varieties of Musical Disorders: The Montreal Battery of Evaluation of Amusia. *Annals of the New York Academy of Sciences*, 999, 58–75.
<https://doi.org/10.1196/annals.1284.006>
- Rogalsky, C., Rong, F., Saberi, K., & Hickok, G. (2011). Functional Anatomy of Language and Music Perception: Temporal and Structural Factors Investigated Using Functional Magnetic Resonance Imaging. *Journal of Neuroscience*, 31(10), 3843–3852. <https://doi.org/10.1523/JNEUROSCI.4515-10.2011>
- Santoro, R., Moerel, M., De Martino, F., Goebel, R., Ugurbil, K., Yacoub, E., & Formisano, E. (2014). Encoding of Natural Sounds at Multiple Spectral and Temporal Resolutions in the Human Auditory Cortex. *PLoS Computational Biology*, 10(1), e1003412. <https://doi.org/10.1371/journal.pcbi.1003412>
- Schlaug, G., Jäncke, L., Huang, Y., & Steinmetz, H. (1995). In Vivo Evidence of Structural Brain Asymmetry in Musicians. *Science*, 267(5198), 699–701.
- Schön, D., Gordon, R. L., & Besson, M. (2005). Musical and linguistic processing in song perception. *Annals of the New York Academy of Sciences*, 1060, 71–81. <https://doi.org/10.1196/annals.1360.006>
- Tervaniemi, M. (2006). From Air Oscillations to Music and Speech: Functional Magnetic Resonance Imaging Evidence for Fine-Tuned Neural Networks in Audition. *Journal of Neuroscience*, 26(34), 8647–8652. <https://doi.org/10.1523/JNEUROSCI.0995-06.2006>
- Wallentin, M., Nielsen, A. H., Friis-Olivarius, M., Vuust, C., & Vuust, P. (2010). The Musical Ear Test, a new reliable test for measuring musical competence. *Learning and Individual Differences*, 20(3), 188–196. <https://doi.org/10.1016/j.lindif.2010.02.004>
- Willems, R. M., Der Haegen, L. Van, Fisher, S. E., & Francks, C. (2014). On the other hand: Including left-handers in cognitive neuroscience and neurogenetics. *Nature Reviews Neuroscience*, 15(3), 193–201. <https://doi.org/10.1038/nrn3679>
- Wong, C. K., Zotev, V., Misaki, M., Phillips, R., Luo, Q., & Bodurka, J. (2016). Automatic EEG-assisted retrospective motion correction for fMRI (aE-REMCOR). *NeuroImage*, 129, 133–147. <https://doi.org/10.1016/j.neuroimage.2016.01.042>
- Zotev, V., Yuan, H., Phillips, R., & Bodurka, J. (2012). EEG-assisted retrospective motion correction for fMRI: E-REMCOR. *NeuroImage*, 63(2), 698–712. <https://doi.org/10.1016/j.neuroimage.2012.07.031>