



Influence of an optimal stomatal conductance scheme in ACCESS1.3b under current and future climate


Jatin Kala

(J.Kala@murdoch.edu.au / Jatin.Kala.JK@gmail.com)

26th August 2015



What is Stomatal Conductance (g_s)?




Schematic of a cross-section through a stomate (Pitman 2003)



Why does Stomatal Conductance (g_s) matter?


- ▶ Vegetation is a net Carbon Sink
- ▶ Carbon sequestration via re-vegetation is an option



Schematic of the terrestrial Carbon cycle (from IPCC AR4)

Why does Stomatal Conductance (g_s) matter?

- ▶ Vegetation is a net Carbon Sink
- ▶ Carbon sequestration via re-vegetation is an option
- ▶ Observations show that as atmospheric CO₂ ↑
 - ▶ Stomata act “optimally” to lose less water during photosynthesis
 - ▶ g_s schemes need to mimic this “optimal” behaviour accurately



Schematic of the terrestrial Carbon cycle (from IPCC AR4)



Why does Stomatal Conductance (g_s) matter?



Preferential cloud formation over native vegetation in Western Australia (Lyons 2002)

- ▶ g_s controls the amount of H_2O transpired by vegetation
- ▶ Influences the partitioning of available energy into latent and sensible heat
- ▶ Influences the boundary layer



Why does Stomatal Conductance (g_s) matter?



Preferential cloud formation over native vegetation in Western Australia (Lyons 2002)

- ▶ g_s controls the amount of H_2O transpired by vegetation
- ▶ Influences the partitioning of available energy into latent and sensible heat
- ▶ Influences the boundary layer
- ▶ Several papers have shown the response of vegetation to $\uparrow CO_2$ via g_s has both regional and global impacts



Default g_s parameterisation in CABLE (Leuning et al. (1995)):

$$g_s = g_0 + \frac{a_1 \beta A}{(C_s - \Gamma)(1 + \frac{D}{D_0})} \quad (1)$$



Default g_s parameterisation in CABLE (Leuning et al. (1995)):

$$g_s = g_0 + \frac{a_1 \beta A}{(C_s - \Gamma)(1 + \frac{D}{D_0})} \quad (1)$$

The Key Point here is:

- ▶ The fitted parameters g_0 and a_1 do Not vary by PFT but only by photosynthetic pathway (C3 versus C4)
 - ▶ No data to inform parameter choice = little theoretical meaning



Default g_s parameterisation in CABLE (Leuning et al. (1995)):

$$g_s = g_0 + \frac{a_1 \beta A}{(C_s - \Gamma)(1 + \frac{D}{D_0})} \quad (1)$$

The Key Point here is:

- ▶ The fitted parameters g_0 and a_1 do Not vary by PFT but only by photosynthetic pathway (C3 versus C4)
 - ▶ No data to inform parameter choice = little theoretical meaning
- ▶ But we know from observations that different PFTs differ in water-use strategies
 - ▶ New observations of g_s have recently lead to improved schemes



New g_s scheme in CABLE (Medlyn et al. (2011)):

$$g_s = g_0 + 1.6 \left(1 + \frac{g_1 \beta}{\sqrt{D}} \right) \frac{A}{C_s} , \quad g_1 \propto \sqrt{\frac{\Gamma^*}{\lambda}} \quad (2)$$

The key point here is:

- ▶ Same functional form as Leuning (1995) but:
 - ▶ g_1 varies per PFT and derived from 314 species at 56 sites
 - ▶ Comprehensive collation of observations of how g_s varies with CO₂.
 - ▶ Eq. 2 thoroughly tested, including data from elevated CO₂ experiments.



New g_s scheme in CABLE (Medlyn et al. (2011)):


$$g_s = g_0 + 1.6 \left(1 + \frac{g_1 \beta}{\sqrt{D}} \right) \frac{A}{C_s} , \quad g_1 \propto \sqrt{\frac{\Gamma^*}{\lambda}} \quad (2)$$

The key point here is:

- ▶ Same functional form as Leuning (1995) but:
 - ▶ g_1 varies per PFT and derived from 314 species at 56 sites
 - ▶ Comprehensive collation of observations of how g_s varies with CO₂.
 - ▶ Eq. 2 thoroughly tested, including data from elevated CO₂ experiments.
 - ▶ g_1 parameter provides a physical link to the plant's water-use strategy
 - ▶ relates to the marginal cost of water to the plant

Offline performance - Prescribed Meteorological forcing

- ▶ New Scheme results in a ↓ in evapotranspiration over Boreal forests in NH summer by up to 0.8 mm day^{-1}



Change in evapotranspiration between the default g_s scheme (LEU) and the new scheme (MED-P) (De Kauwe and Kala et al. 2015)



Coupled Land-Atmosphere ACCESS simulations

- ▶ AMIP style simulations (prescribed SSTs)
 - ▶ Historical simulations, 1950 to 2012 with observed SST
 - ▶ Compare with obs
 - ▶ RCP8.5 simulations, 2012 to 2100 with CMIP5 SST
 - ▶ Influence on future climate projections


Coupled Land-Atmosphere ACCESS simulations

- ▶ AMIP style simulations (prescribed SSTs)
 - ▶ Historical simulations, 1950 to 2012 with observed SST
 - ▶ Compare with obs
 - ▶ RCP8.5 simulations, 2012 to 2100 with CMIP5 SST
 - ▶ Influence on future climate projections
- ▶ Two experiments, 5 ensembles each, initialised a year apart
 - ▶ Control simulation with default Leuning (1995) g_s scheme (LEU)
 - ▶ Actual experiment with new Medlyn et al. (2011) g_s scheme (MED)

Coupled Land-Atmosphere ACCESS simulations

- ▶ AMIP style simulations (prescribed SSTs)
 - ▶ Historical simulations, 1950 to 2012 with observed SST
 - ▶ Compare with obs
 - ▶ RCP8.5 simulations, 2012 to 2100 with CMIP5 SST
 - ▶ Influence on future climate projections
- ▶ Two experiments, 5 ensembles each, initialised a year apart
 - ▶ Control simulation with default Leuning (1995) g_s scheme (LEU)
 - ▶ Actual experiment with new Medlyn et al. (2011) g_s scheme (MED)
- ▶ All results shown will be the ensemble mean

Evaluation under current climate - PFT distribution



1 - Evergreen needleleaf

2 - Evergreen broadleaf

3 - Deciduous needleleaf

4 - Deciduous broadleaf

5 - Shrub

6 - C3 Grassland


7 - C4 Grassland

8 - Tundra

9 - C3 Cropland


10 - Barren

Evaluation under current climate - TMAX






Evaluation under current climate - TXx







Evaluation under current climate - TMIN




Evaluation under current climate - DTR



Evaluation under current climate - ET/latent heat flux



Evaluation under current climate - PRECIP





Evaluation under current climate - Summary


- ▶ The new scheme reduces latent heat flux over boreal forests using NH summer by 0.5 to 1.0 mm day⁻¹






Evaluation under current climate - Summary

- ▶ The new scheme reduces latent heat flux over boreal forests using NH summer by 0.5 to 1.0 mm day⁻¹
 - ▶ Leads to warmer daily maximum and minimum temperatures by up to 1.0°C
 - ▶ Leads to warmer extreme maximum temperatures by up to 1.5°C
 - ▶ We improve the model's climatology by 10-20%


Influence on future climate (RCP8.5) - MED minus LEU



Influence on future climate (RCP8.5)




Influence on future climate (RCP8.5) - MED minus LEU





Damping effect towards 2099





Influence on future climate (RCP8.5) - Summary

- ▶ Large increases in warm extremes under RCP8.5 during summer with new scheme over NH up to 4-5°C!!!
 - ▶ Driven by reduction in latent heat flux (and associated increase in sensible heat)

Influence on future climate (RCP8.5) - Summary

- ▶ Large increases in warm extremes under RCP8.5 during summer with new scheme over NH up to 4-5°C!!!
 - ▶ Driven by reduction in latent heat flux (and associated increase in sensible heat)
- ▶ Magnitude of increase, decreases towards 2070-2099
 - ▶ At high leaf temperatures, photosynthesis and stomatal conductance (and thus ET) are reduced due to photosynthetic inhibition
 - ▶ This minimises the differences in ET between the two models

Take home message

- ▶ We introduced a new, more physically grounded g_s scheme in CABLE, based on best available data
 - ▶ We tested the scheme at single sites and globally offline
 - ▶ This showed improvements in CABLE's ET/latent heat flux simulations

Take home message

- ▶ We introduced a new, more physically grounded g_s scheme in CABLE, based on best available data
 - ▶ We tested the scheme at single sites and globally offline
 - ▶ This showed improvements in CABLE's ET/latent heat flux simulations
- ▶ We then tested the new scheme coupled within ACCESS
 - ▶ We show a general improvement in ACCESS's representation of warm extremes
 - ▶ This has serious implications for future climate
 - ▶ We are currently under-predicting warm extremes



Take home message

- ▶ Historical simulations and evaluation against observations currently under review:
 - ▶ Kala et al. (2015) Implementation of an optimal stomatal conductance model in the Australian Community Climate Earth Systems Simulator (ACCESS1.3b), *Geosci. Model. Dev. Discuss., under review*

Take home message

- ▶ Historical simulations and evaluation against observations currently under review:
 - ▶ Kala et al. (2015) Implementation of an optimal stomatal conductance model in the Australian Community Climate Earth Systems Simulator (ACCESS1.3b), *Geosci. Model. Dev. Discuss., under review*
- ▶ Impacts on future climate submitted to a journal yesterday
 - ▶ Watch this space!



Future work on g_s in CABLE/ACCESS

- ▶ Link changes in heat fluxes to changes in Carbon fluxes a bit more



Future work on g_s in CABLE/ACCESS

- ▶ Link changes in heat fluxes to changes in Carbon fluxes a bit more
- ▶ Suggestion that g_1 parameter varies with climate



Future work on g_s in CABLE/ACCESS

- ▶ Link changes in heat fluxes to changes in Carbon fluxes a bit more
- ▶ Suggestion that g_1 parameter varies with climate
- ▶ We use prescribed leaf area index
 - ▶ Prognostic LAI in CABLE is under-development
 - ▶ We expect differences to be larger



Future work on g_s in CABLE/ACCESS

- ▶ Link changes in heat fluxes to changes in Carbon fluxes a bit more
- ▶ Suggestion that g_1 parameter varies with climate
- ▶ We use prescribed leaf area index
 - ▶ Prognostic LAI in CABLE is under-development
 - ▶ We expect differences to be larger
- ▶ Do we see the same sign and magnitude of change if the new g_s scheme is implemented in other LSMs which use the Leuning (1995) scheme?



Influence of an optimal stomatal conductance scheme in ACCESS1.3b under current and future climate

Jatin Kala

Questions?

Evaluation under current climate - NPP

