

Collaboration for the Analysis of Photonic and Ionic Bursts and Radiation from Barcelona (CAPIBARA): Preliminary Report for the SPARK Program

Joan Alcaide-Núñez^{*1}, Martí Delgado Farriol², Lluc Soler Manich³, Anna Abadal Garrido⁴, Emma Massó Sala³, Christian Schaefer González¹, Hongda Zheng⁵, Alicia Lòpez ^{†3}, Susana De Mesa Sterthoff ^{‡1}, and Carles Fonseca Mauri⁶

¹Deutsche Schule Barcelona (DSB), Av. Jacint Esteve i Fontanet, 105, 08950 Esplugues de Llobregat, Barcelona, Spain.

²Col·legi Pureza de Maria Sant Cugat, Carrer de Mercè Vilaret, 21, 08174 Sant Cugat del Vallès, Barcelona, Spain

³CIC Escola de Batxillerats, Via Augusta, 205, 08021 Barcelona, Spain

⁴Salesians Sarrià, Passeig Sant Joan Bosco, 42, 08017 Barcelona, Spain

⁵Institut Les Corts, Travessera Les Corts, 131, 08028 Barcelona, Spain

⁶Jesuïtes Sarrià - Sant Ignasi, Carrasco i Formiguera, 32, 08017 Barcelona, Spain

November 10, 2024

Abstract

This report presents the preliminary concept for a satellite mission developed by the CAPIBARA Collaboration to detect, observe, and study high-energy astrophysics in both hadronic and photonic states. The missions aim to advance research in these fields while providing young students with first-hand experience in cutting-edge science and engineering. We discuss the mission's objectives, justification, scientific rationale, instrument design, future plans, and introduce the collaboration team.

^{*}Corresponding author: joanalnu@outlook.com

[†]Supporting Teacher

[‡]Supporting Teacher

Contents

1	Introduction	3
2	Mission Objectives	3
2.1	Primary CRs Detection	3
2.2	Photonic Detection	4
2.3	Deorbitation & Sustainability	4
3	Justification	4
3.1	Atmospheric Opacity	4
3.2	Application	5
3.3	Observational and Research Gap	5
4	Scientific Research	6
4.1	Gamma-Ray Burst Cosmology	6
4.2	Exoplanet Research from CR Interaction Radiation	7
4.3	Characterization of PCR Radiation from solar origin	8
4.4	Observation of Supernova Remnants	8
4.5	AGN Jet Dynamics and Cosmic Ray Interactions	9
5	Experiment Design	9
5.1	Primary Cosmic ray Detector	9
5.2	X-ray Observatory	10
5.3	γ -ray Observatory	11
6	Data Management Plan	12
6.1	Onboard Data Processing	12
6.2	Data Publication	12
7	Project Timeline	13
8	Team Presentation	14
8.1	Joan Alcaide-Núñez	14
8.2	Martí Delgado Farriol	14
8.3	Lluc Soler Manich	14
8.4	Anna Abadal Garrido	14
8.5	Emma Massó Sala	14
8.6	Christian Schaefer González	14
8.7	Hongda Zheng	14
8.8	Carles Fonseca	14
9	Conclusion	15

1 Introduction

Cosmic Rays (CRs) or cosmic radiation are charged particles coming from the Universe at high energy due to their high velocities (Lerner, 2024). These particles are mainly protons (87%) and Helium nuclei (10%) (de Espectroscopía Gamma y de neutrones, 2014), see table 1 for a complete composition. We refer to these particles as Primary Cosmic Rays (PCRs).

Type	Percentage
Protons	$\sim 87\%$
α particles	$\sim 10\%$
Electrons	$\sim 2\%$
Light elements (Li, Be, B, ...)	$\sim 0.25\%$
Antimatter	$\sim 0.01\%$

Table 1: Composition of primary cosmic rays from de Espectroscopía Gamma y de neutrones (2014).




Figure 1: Contour plot of the muon deficit as measured by IceCube in the region around the Moon's position, the so-called on-source region (credit: Ice-Cube Collaboration)

When these particles interact with a medium, they decay in lighter particles, called Secondary Cosmic Rays (SCRs). For instance, SCRs are produced as a consequence of PCRs decaying when hitting molecules in Earth's upper atmosphere. As a result, protons, alpha particles, pions, kaons, muons, electrons, and neutrinos are created (Morison, 2008), becoming a way of indirect detection (see Figure 1).

Furthermore, there is another radiation which helps trace CRs. However, it can not be categorized as cosmic rays due to its lack of electric charge. The interaction of both PCRs and SCRs emits light, for instance in accretion disks, nebulae, interstellar dust, quantum decays, atmospheres, among others. This type of light reaches us in high-energy wavelengths, mostly in X-rays and γ -rays, but also in the ultraviolet (UV) and infrared (IR) spectrum. These photons have different properties as compared with fermions and hadrons, meaning a totally different way to study the Universe.

Furthermore, CRs can also be divided by origin, which enables differentiation between galactic and extra-galactic CRs. Galactic CRs emerge mainly from the sun, other nearby stars, or other objects such as neutron stars (NSs). This category may be estimated by the source coordinates, where low latitudes more likely correspond to the position in the galactic plane, whereas high latitudes likely come from other galaxies. Knowing the distance to the source is also a useful tool for estimating the origin¹.

Although CRs are known to come from outside of Earth (Hess, 1912), we do not know specific sources of CRs. As these are charged particles, they interact with magnetic fields through their way to Earth. Therefore, when we detect cosmic rays, we cannot determine their provenance. However, there are probable sources of CRs discussed: supernovae explosions (SNe) and remnants (SNRs), neutron stars (NSs), microquasars (Quasi Stellar Object, QSO), active galactic nuclei (AGNs), and gamma-ray bursts (GRBs).

2 Mission Objectives

Our objective is to explore high-energy phenomena, both in the form of light (photons) and particles (mainly protons and alpha particles). In this section, the mission objectives and technological requirements of the mission are described. For further details on the experiment design, i.e. how we aim to accomplish these goals, see section 5

2.1 Primary CRs Detection

As explained before (see table 1), CRs mainly consist of protons (p^+) and α -particles, i.e. Helium nuclei, (α , α^{2+} , or He^{2+}); summing up to 97%. In order to explore the high-energy particles, providing useful data for the study of CRs, solar wind, astroparticle physics, and further topics of research, we aim to detect these particles using a variety of detectors. Each of the types of detectors, outlined in section 5 are capable of detecting a different property of these particles (e.g. velocity, energy, etc.). Therefore, all of them are needed to provide complete and reliable datasets.

¹This technique is only applicable to electromagnetic radiation, as charged particles interact with magnetic fields, therefore making their direction an unreliable source of origin information.

2.2 Photonic Detection

Moreover, we want to accomplish the observation of electromagnetic radiation emitted by CRs interactions, as they give us different information about CR sources and physical processes of their interactions due to the difference in nature of light. To do so, we have to be capable to detect photons in the X-ray and γ -ray range of the spectrum, as high-energies are where CRs are expected to emit on account of their initial energy and velocity.

A cosmic X-ray Background (CXB) was claimed with the launch of the first X-ray telescopes (e.g. ROSAT) in the late 20th century. [Gilli \(2013\)](#) showed a collection of different CXB measurements. However, more modern space telescopes have resolved the vast majority of this background into discrete, faint sources. As a consequence, with enough resolution, this 'background' radiation is negligible. For reference, see Figure 7 of [Tozzi et al. \(2001\)](#), where high-energies are expected to be filled by dust-obscured AGNs [Schneider \(2006\)](#).

The existence of a cosmic γ -ray background (CGRB) has been studied by the FERMI-LAT instrument in detail. It is believed that this radiation emerges from unresolved source (e.g. SNe, pulsars, or AGNs) and possible exotic processes like dark matter (DM) annihilation or decay. The background is far from negligible and must be accounted in γ -ray data analysis. common techniques to consider this radiation are image background subtraction (both instrumental background and template fitting), background modeling, and likelihood analysis (e.g. Markov Chain Monte Carlo (MCMC)).

Technical requirements to consider are that we expect to perform both photometric and spectroscopic analysis of observations. This is especially important, as both approaches provide different information and data about the phenomena, complementing each other.

Furthermore, light observations require the capability of pointing in the sky to localize sources. We intend to use a wide-field γ -ray camera to monitor events, this does not need a more precise accuracy than to localize sources in $5''$ to $1''$ (arcseconds). While a narrower field of view (FoV), should have a pointing accuracy between $0.5''$ and $1''$.

2.3 Deorbitation & Sustainability

A crucial aspect of modern space mission is ensuring they adhere to sustainability guidelines, including proper end-of-life procedures for deorbitation. Space debris represents a danger to space traffic and human spaceflight, as well as posing a threat to Earth and Space based astronomical observations. Therefore, we want our mission to incorporate a well-defined deorbitation plan to ensure that the satellite does not contribute to this growing problem, following international space mitigation standards. Applicable measures orbital decay monitoring, end-of-mission deorbit burn, sustainability compliance, and post-mission disposal.

Given that our satellite will operate in a Sun-Synchronous Orbit (SSO), as given by the Spark program, we aim to perform a controlled deorbitation maneuver at the end of the mission's operational lifetime. While natural orbital decay due to atmospheric is slower at the higher altitudes typical of SSO, we plan to initiate a controlled deorbitation sequence. This will involve reducing the satellite's altitude through propulsion-based maneuvers, eventually leading to a final deorbit burn. The objective is to ensure that the satellite safely re-enters Earth's atmosphere, where it will burn up, minimizing the risk of debris surviving re-entry and reaching the surface.

3 Justification

3.1 Atmospheric Opacity

The radiation we aim to detect, which consists of both particles and light, is not possible to study from the Earth's surface.

PCRs interact with Earth's upper atmosphere, creating particle showers, which are detectable from Earth, and so-called SCRs. However, the particles resulting from these interactions have lost part of the information inferable from PCRs detections. Furthermore, Earth's magnetic field also shapes how SCRs are detected. Therefore, the effect of magnetic paths should be smaller using space-based detectors.

Additionally, neither X-ray nor γ -ray radiation do penetrate Earth's atmosphere (see Figure 2), which means that their detections are only possible with space-based telescopes ([Brandt et al., 1992](#)). Therefore, the observation of this type of emission is of huge importance, as it is the only way to study high-energy phenomena ([Lyman Spitzer, 1990](#)).

All in all, high-energy particles and light with cosmic origin do not reach Earth's surface, meaning the development of space-based detection mission is crucial for the study of these phenomena.




Figure 2: Atmospheric electromagnetic opacity by wavelength. (credit: NASA)

3.2 Application

Furthermore, the fundamental research in cosmic rays has wide applications. The most obvious are particle physics, nuclear and atomic physics, as well as astronomy and astrophysics.

However, also geophysics and climate studies benefit, as CRs can help us study the formation of clouds, monitor the atmosphere (Kuwabara et al., 2006), and develop techniques for geologic dating (Blard et al., 2006). What is more, medicine benefits from both the best medical imaging methods, e.g. positron emission (PET); as of radiotherapy, the research of radiation for the development of medical treatments (Royon et al., 2019).

The improvement and development of new imaging techniques inspired by cosmic ray detection help to study the strength and durability of objects and substances, advancing in material science. Likewise, imaging methods allow the study of the interior of pyramids, caves, volcanoes, and other structures.

In addition, living beings in space suffer from the impact of radiation against their bodies, therefore, studying the nature of cosmic rays would help us innovate in potential solution for the protection against such radiation, which would allow very long duration crewed missions (Saganti et al., 2004).

3.3 Observational and Research Gap

Finally, given the current landscape in space observation, the study of cosmic rays and high-energy phenomena seemed particularly compelling. Although these topics are at the forefront of modern astrophysics, representing relatively new areas, many modern telescopes, such as the James Webb Space Telescope (JWST), Euclid, the Vera C. Rubin Observatory (LSST), or the Nancy Grace Roman Space Telescope (WFIRST), focus primarily on electromagnetic radiation in wavelengths close to the optical range (partially covering the UV and IR).

Furthermore, two of NASA's Great Observatories² were once dedicated to high-energies, Compton (CGRO) and Chandra (CXO). However, the Compton Observatory was deorbited in 2000, and the Chandra Observatory is facing funding issues without current plans for replacements³. Other γ -ray telescopes such as SWIFT (launched 2004) or FERMI (launched 2008) are also turning old.

Therefore, our objective of exploring the high-energy range by observing light as well as particles (CRs) align with the current gap in observational capabilities, which develops a gap in research.

All in all, these factors highlight a limitation in observational capabilities that develops a research gap, which we intend to address with our mission by exploring the high-energy range through the observation of both light and cosmic rays.

²Spitzer (IR), Hubble (NIR, Optical & UV), Chandra (X-ray), and Compton (γ -ray)

³Chandra Decommissioning




Figure 3: Figure showing the observational coverage of the high-energy range of the electromagnetic spectrum. Shaded in blue is the x-ray range, and in red the γ -ray one. Telescopes observational capabilities are plotted along with their age or predicted launch date. It can be seen that there's an observational gap in the γ -ray part of the spectrum, and that the x-ray part is more crowded, though the majority are old (<20 yrs) or short-mission telescopes.

4 Scientific Research

4.1 Gamma-Ray Burst Cosmology

- Author: Joan Alcaide-Núñez
- Field: Cosmology and Extragalactic Astrophysics
- Context: The Hubble-Lemaître constant ([Hubble, 1929](#)) ([Lemaître, 1927](#)) represents the local expansion rate of the Universe, sets its overall scale, and enables determination of its age and history. It is a fundamental measure in cosmology and a key stone for cosmological models of the Universe ([Freedman et al., 2010](#)). Since its first derivation in the 1920s, it has been recomputed multiple times ([Freedman, 2021](#)) ([Valentino et al., 2021](#)). However, its exact value is not yet known, as different methods result in different values ([Freedman et al., 2010](#)) ([Aghanim et al., 2020](#)) ([245 et al., 2017](#)). Not only that, but as techniques and instrumentation have become more precise, the discrepancy has gotten stronger ([Riess et al., 2024a](#)) ([Valentino et al., 2019](#)) ([Freedman, 2021](#)). This problem is called the Hubble Tension, often also referred to as the Crisis in Cosmology.

For reference, see the latest measurements for each method in Figure 4; Local Distance Ladder ([Freedman et al., 2024](#))⁴ ([Riess et al., 2024b](#))⁵; CMB ([Aghanim et al., 2020](#))⁶; BAO ([Collaboration et al., 2024](#))⁷; and Gravitational Waves ([245 et al., 2017](#))⁸.

⁴JAGB (J-region Asymptotic Giant Branch), TRGB (Tip of the Red Giant Branch), Cepheids, and Ia SNe (Type Ia Supernovae)

⁵Different Ia SNe catalogs

⁶Cosmic Microwave Background

⁷Baryonic Acoustic Oscillations

⁸Standard Siren method




Figure 4: Probability Density Function diagram for different measurements of the Hubble Constant. Data from: Freedman et al. (2024) (blue), Riess et al. (2024b) (red), Aghanim et al. (2020) (black), Collaboration et al. (2024) (orange), 245 et al. (2017) (green), Freedman et al. (2001) (purple).

Furthermore, relatively recently it was discovered that the expansion rate of the Universe is accelerating (Riess et al., 1998), this discovery having very important implications in our understanding of the Universe and the crafting of the Λ CDM model⁹ (Peebles et al., 2003). To do so, it was required to observe SNe in very distant galaxies. This has additional difficulties due to the reduction of spatial resolution (crowding effect (Freedman et al., 2024)) and the diminishing luminosity. However, only at this greater distance, the effect of an accelerating, neutral, or decelerating expansion rate are observable. More recently, some research has posed that dark energy, the substance responsible for accelerating the expansion, is evolving in function of time (Efstathiou, 2024)(DESI Collaboration et al., 2024). If true, this could be the first hint about the nature of dark energy since its discovery.

- **Goals:** Our aim is to use Gamma-Ray Bursts (GRBs) as cosmological distance indicators. GRBs are the most energetic events in the Universe, releasing in just a few seconds more energy than the Sun emits over its entire lifetime. Hence, they can be observed at great distances. For reference, GRBs have been observed up to $z > 8$, while the more distant Ia SN is at $z < 2$. The standardisation of their high luminosity and the use of GRBs as cosmological distance indicators will provide very valuable data for the study of the Hubble Tension, the nature of dark energy, and thus the validation of Λ CDM. Furthermore, the impact on the field is considerable, as greater distances will enable more precise fitting of cosmological parameters to the data, and new measurements.
- **Methods:** γ -ray photometric and X-ray photometric and spectroscopic analysis; identification of GRBs; study of new methods for the standardization of GRBs cosmological distances; cross-check with external databases for redshift data.

4.2 Exoplanet Research from CR Interaction Radiation

- **Author:** Joan Alcaide-Núñez
- **Field:** Earth and Planetary Sciences
- **Context:** Since the discovery of the first exoplanet in 1992 (Wolszczan et al., 1992) and the discovery of the first exoplanet around a star in 1995 (Mayor et al., 1995), the observation of new exoplanets has grown exponentially, as has the field (Editorial, 2024). IN 2001 the first exoplanetary atmosphere was confirmed via spectroscopy during planetary transits¹⁰ (Charbonneau et al., 2002). The first direct of an exoplanet was achieved by the VLT¹¹ (ESO)¹² in 2006 (?).

Moreover, Secondary Cosmic rays (SCRs) can be detected on Earth's surface (including at sea level) due to the decay of PCRs as they hit molecules in the upper atmosphere (). This event is also known as

⁹A cosmological model for a Universe with dark energy (Λ) and Cold Dark Matter (CDM).

¹⁰A planetary transit happens when a planet passes between its host star and the observer.

¹¹Very Large Telescope

¹²European Southern Observatory

particle shower, an example being shown in figure 5. Exoplanets with atmosphere are expected to express the same or similar γ -ray signatures based on these processes. The strength of this emission will depend on the source’s characteristics, the atmospheric composition, and the distance to the exoplanet.




Figure 5: Cosmic ray air shower scheme by the European Council for Nuclear Research (CERN, acronym in French). It is an example of the particles resulting from the decay of PCRs in Earth’s atmosphere.

- Goals: We aim to detect exoplanet SCRs signature coming from the interaction of PCRs with the exoplanetary atmospheres.
- Methods: Transit observations, X-ray & γ -ray astronomy (photometry and spectroscopy), exoplanets, host stars

4.3 Characterization of PCR Radiation from solar origin

- Author: Martí Delgado Farriol
- Field: Solar and Stellar Astrophysics, Earth and Planetary Sciences
- Context: Primary Cosmic Rays (PCRs) are thought to come from high-energy phenomena (Lerner, 2024). Nevertheless, we also know about stellar (and/or solar) wind, which is a flow of charged particles ejected from the Sun’s atmosphere the corona) (ESA). This charged particles can pose a danger to satellites (e.g. weather monitoring, communications, and GPS satellites) and human space flight.
- Goals: Our goal is to differentiate between solar PCRs and those coming from outside the solar system. We aim to characterize the sun’s radiation and understand the physical background phenomena of PCR sources by differentiating particle radiation (mainly photons) during orbiting Earth’s day, in which solar radiation will be present, with radiation detected on the night side orbit when solar radiation decreases. We expect solar PCRs to have slower velocities and lower energies. Furthermore, their interaction with magnetic fields should be easy to track due to the short distance. The understanding of CRs within the solar system is key to developing practical applications to protect long-duration missions.
- Methods: Particle detection, magnetic fields, orbit trajectories.

4.4 Observation of Supernova Remnants

- Author: Hongda Zheng
- Field: Supernova Remnants
- Context: Supernova Remnants (SNRs) are the material leftover of a supernova explosion (SNe). There are two criteria of classification of SNe which originates different subtypes:
 - Core Collapse (Type II SNe): They are the result of massive stars rich in iron that ran out of material to fusion, since no energy can be gained from nuclear fusion of iron. Then they collapse into proton-neutrons star (black holes for the most massive stars).
 - Thermonuclear explosion (Type I SNe): They are thought to be the explosion of Carbon/Oxygen White Dwarfs stars (no massive stars) that generate explosive nuclear burning. In both cases, the explosion leaves behind its nucleus and produces an explosion and the ejection of stellar materials such as gas or dust. Those rests are the SNR. GSFC (2021). The SNR consist of core and these materials around it, which are in constant expansion.

SNRs are very interesting astronomical phenomena since it allow us to study the creation of various chemical elements in the stars and what composition did they have. In addition, there is an ejection of heavy metals that contribute to the formation of new stars. SNRs also emit energy in different forms, such as X-Rays synchrotron energy, which allows us to know if there is or not a neutron star in it (Ishikawa, 2019).

Even with this classification of SNe, the supernova origin of SNRs is often difficult to establish. Because of that, SNRs have a classification of their own, which is based on their morphology. This classification recognizes three different classes: - Shell type SNRs: As the blast wave sweeps through the interstellar medium. the shockwave from the supernova explosion that plows through space, heats and stirs up any interstellar material it encounters and produces a big shell of hot material in space As the blast wave sweeps through the interstellar medium. - Crab type SNR or plerions: in this case, the nebulae is filled with high-energy electrons that are flung out from a pulsar in the middle, producing a blob form - Composite SNRs: they are the cross between the shell-type remnants and crab-like remnants.

- Goals: Our aim is to determine the category of a candidate of SNR and which its magnetic field strength is through analyzing which kinds and levels of radiation it emits and how they affect shock waves.
- Methods: X-ray Synchrotron Radiation Detection and Spectroscopic Analysis.

4.5 AGN Jet Dynamics and Cosmic Ray Interactions

- Author: Carles Fonseca Mauri
- Field: Active Galactic Nuclei
- Context: Active Galactic Nuclei (AGN) are among the most energetic astrophysical phenomena, with relativistic jets originating from their central supermassive black holes (SMBHs), Blandford et al. (1977). These jets are capable of injecting vast amounts of energy into the interstellar and intergalactic medium, impacting gas dynamics, star formation rates, and the broader evolution of galaxy clusters. Understanding the precise influence of AGN jets, particularly their role in quenching star formation and generating cosmic rays (CRs), remains an active research area. The CAPIBARA mission provides a novel opportunity to study these high-energy processes.
- Goals: We aim to compare the simulated high-energy outputs and particle distributions from AGN jets using the SWIFT simulation code (Schaller et al. (2024)) to CAPIBARA's expected observational data on CRs and associated photon emissions. Through these simulations, AGN jets are analyzed in terms of their jet power, spin dynamics, and interactions with the surrounding medium, with special attention to shock-induced CR production and X-ray cavity formation.
- Methods: This study employs SWIFT-based simulations of AGN jets, mirroring setups similar to those used by Huško et al (e.g. Huško et al. (2024)) to model jet feedback and black hole spin dynamics in galaxy groups and clusters.

By drawing comparisons between the simulated jet energetics and CAPIBARA's CR observations, this study aims to enhance our understanding on how AGNs contribute to the CR population and high-energy photon background. Furthermore, this context allows us to explore how AGN feedback mechanisms shape the evolution of galaxy clusters, offering theoretical support for CAPIBARA's objectives in mapping high-energy cosmic processes.

5 Experiment Design

You can see specifications on <https://spark-program.pldspace.com/es/descripcion.html> or ask for detail in the slack channel.

5.1 Primary Cosmic ray Detector

This experiment must be capable of detecting CR protons and, if possible, α -particles. The two most frequent CR particles (see 1).

When considering direct measurements, as this is the case, combining different detectors is essential to obtain a full characterization of the particles (Piera, 2012). Therefore, the detector should be "layer" built, basically stack the detectors together obtaining a relatively compact device (taking into consideration the distance between detectors needed to perform some detections). There are three types of measurements which need separate devices, which are the following:

- **Time-of-Flight (TOF):** Time-of-flight detectors work by pairs, and they trigger a clock when a particle hits the first one, which is stopped when the second one is hit. Then, through a series of calculations, the particle velocity can be obtained. In our case, and as many other satellites CR detectors (i.e. PAMELA¹³, AMS¹⁴), scintillators can be used. Scintillators are materials that exhibit scintillation, that is the physical process through which a material emits a flash of light when it is hit by a charged particle, for instance protons or *alpha* particles (Maddalena F, 2019). This light is then detected with photo-multipliers which are part of an electrical circuit the can recognize a difference in the voltage. The main disadvantage of scintillators is that the light produced is not linear, that means that the energy can not be inferred (Zeitlin, 2012).
- **Charge Detector:** Essentially, this detector determines whether the charge is positive or negative and the value itself. Among the available options, on that is fairly used in other detectors, are Silicon Detectors. They work as diodes to which a voltage is applied, acting as a parallel-plate ionization chamber, that is, when a charged particle passes through it generates a current (Zeitlin, 2012).
- **Energy Detector:** The principal function of this apparatus is to establish the total energy of the particle. In this case, an electromagnetic calorimeter is required. It works by the principle of "destruction", that means a particle loses all its energy passing through the detector, which is converted into a signal (Masciocchi, 2017).

5.2 X-ray Observatory

Electromagnetic detection in the X-ray range. **Pointing system with a pointing accuracy of about 5'' and 0.5'' in both right ascension (RA) and declination (DEC) coordinates.**

In astronomy, many X-ray detection methods have been discovered and evaluated, and one that has been widely applied within scientific purposes are Charged-Coupled Devices (CCDs), which are arrays of linked capacitors (Grant, 2011). This device converts the photons into photoelectrons and moves an electrical charge. This is negative, and can be shifted along the sensor to readout register where are converted into digital signal (process know as charge transfer, see figure 6. For a better understanding of their principle of working see (Teledyne, 2024a)). When these photons pass through the CCDs, they lose energy through different effects, i.e. Compton scattering, fluorescence or photoelectric effect (Teledyne, 2024b), generating electron-hole pairs¹⁵. Considering a silicon substrate, an average ionization energy of 3.65eV is required. Therefore, we can estimate the energy of the X-ray with the following formula $N_e = E_x/w$. Where N_e is the number of electrons, E_x is the energy of the X-ray photon and w the ionization energy (3.65eV).




Figure 6: Charge transfer process (credit: (Teledyne, 2024a))

The detector would work as a photon-counter, which basically means that it makes a recount of all the photons detected. To ensure high quality measurements, each photon should be collected within only one pixel. Nevertheless, it is possible that one photon interacts with two, or more, different pixels. To address this situation, an intensity-threshold is used to distinguish between single-pixel and multiple-pixel events. A grading system is used in X-ray telescopes with this technology, such as ASCA, which stands for Advanced Satellite for Cosmology and Astrophysics (see figure 7), to discriminate between X-ray and cosmic rays¹⁶ (Grant, 2011).

¹³<https://pamela.roma2.infn.it/>

¹⁴<https://ams02.space/>

¹⁵Electron-hole pairs refer to the excitation in a semiconductor where an electron is excited from the valence band to the conduction band, leaving behind a positively charged hole (Luiz H.G. Tizei, 2017).

¹⁶These grade codes were used in the ASCA mission to discriminate between X-ray and cosmic ray events. Generally, X-rays tend to be simpler events than other cosmic ray events, therefore, they can be differentiated. It is further explained in (Grant, 2011).




Figure 7: ASCA Grade Codes (credit: (Grant, 2011))

5.3 γ -ray Observatory

Wide-field camera for surveying γ -ray sources, less pointing capability is required.

γ -ray detection can be divided in two categories. The first one, includes what are called spectrometers or photometers in optical astronomy, which use scintillators or solid-state detectors. The second, includes detectors that perform γ -ray imaging. This ones rely on the processes that happen due to the nature of γ -rays, for example Compton scattering (GSFC, 2018).

In the first case, the operation of scintillators or solid-state detectors is done in the same way it is done in X-ray astronomy, so in our case these detectors would be very convenient. Nevertheless, a brief description of a specific γ -ray detector is provided:

- **Compton Scattering Detectors:** The so-called Compton Scattering is a phenomenon which occurs when a photon hits an electron and some of the photon energy is transferred to the charged particle, this is the basis of the detection. These telescopes have usually two levels of detectors (see figure). In the first layer, the γ -ray scatters off an electron, then this scattered photon is absorbed by a second layer. With the help of phototubes, an approximation is made to determine the interaction points and the deposited energy (GSFC, 2018).




Figure 8: The structure of a two-level Compton detector (credit: (GSFC, 2018))

6 Data Management Plan

We expect to collect 8 different types of data: (1) CR particle count, (2) CR time-of-flight, (3) CR charge, (4) CR energy, (4, 5) photometric and (6, 7) spectroscopic data from X-ray and γ -ray cameras, and (8) environmental data from the satellite. This data will be collected by: (1) combined CR detectors, (2) time-of-flight detector, (3) charge detector, (4) energy detector, (4, 5) X-ray and γ -ray CCDs, (6, 7) X-ray and γ -ray spectrometers, and (8) specialized sensors on the satellite¹⁷.

6.1 Onboard Data Processing

Onboard data processing is necessary, and as a consequence temporary onboard storage. Some high-energy phenomena only last for minutes or even seconds, therefore it is crucial to have onboard data processing to live monitor observations and identify and localize potential transient detections, as well as creating and sharing alerts via the General Coordinates Network¹⁸ (GCN) (Singer et al., 2023). Another consideration is to perform early-stage data reduction (e.g. instrumental background subtraction and image calibration) onboard, this way we would reduce the amount of data transmitted back to ground.

Regarding ground data processing, an automated or semi-automated data reduction pipeline will be developed by the collaboration and made available publicly. Afterwards, each research project will analyze the data to their convenience, focusing on their specific research requirements.

6.2 Data Publication

Data will be made public by steps, and differently depending on the data type. Firstly, regular observations will be reduced and made public internally in the collaboration as soon as possible. The data will be made totally public about 180 days (6 months) after the observation. The method for massive data publication is still to be determined.

For transient events¹⁹ data will follow the same path as regular data. However, due to the transient nature of these phenomena, some data will be shared via an alert through the GCN. This alerts enable quick follow-up observations with other telescopes and instruments covering diverse wavelengths, as well as cross-check observations.

Additionally, we expect to publish a formal data release paper, where we will extensively explain the data reduction pipeline and data accessing procedure.


We are exploring various data-sharing options include collaborative repository and public databases. Internal data-sharing protocols are still to be disused, with decision on storage solutions, licensing, and cybersecurity measures to be made as the project advances. Furthermore, storage capacity and data load expectations will be evaluated once the instrument specifications are finalized.

¹⁷Note that we do not have a plan for these sensors, but rely on the matchmaking for this aspect.

¹⁸<https://gcn.nasa.gov>

¹⁹As transient events we are considering SNe, GRBs, AGNs, TDEs, Kilonovae, Novae, M-dwarf flares, Variable stars and transient CR phenomena.

7 Project Timeline



Red text indicates deadline established by PLDSpace (from [Application Stages](#)). Also, note that the duration of the scientific operations is not yet decided. (last checked Aug 24, 2024).

8 Team Presentation

We are a group of students from various schools in the greater Barcelona metropolitan area, united by our interest in scientific research and space exploration. Motivated to participate in the Spark program by PLDSpace, we are excited to bring this project to life. Each team member has a unique background and area of expertise, contributing their knowledge and skills to the group. We are also grateful for the support of our science and technology teachers, who have guided us along the way. Below is a brief introduction to all participants. You can find more information about the project and our [team](#) on our website, capibara3.github.io.

8.1 Joan Alcaide-Núñez

- Contact: joanalnu@outlook.com
- Course: 12th grade, 2^o Bachillerato (DSB)
- Field of work: Astronomy & Astrophysics, Cosmology
- Experience: Joves i Ciència '23, Jugend Forscht '24, Explainers '22
- Hobbies: playing the piano, reading, traveling, coding
- Links: [LinkedIn](#), [Personal website](#)

8.3 Lluç Soler Manich

- Contact: llucsoler.m@gmail.com
- Course: 12th grade / 2^o Bachillerato (CIC Escola de Batxillerats)
- Field of work: Astrophysics, Engineering (Aerospace)
- Experience: CIC-MIT Hackathon '22, Bojos per la física '24, BIYSC (Cosmic Rays) '24
- Hobbies: cycling, skiing, playing the guitar, physics

8.5 Emma Massó Sala

- Contact: emmamassomail@gmail.com
- Course: 12th grade, 2^o Bachillerato (CIC Escola de Batxillerats)
- Field of work: Engineering
- Experience: Joves i Ciència '23, CIC-MIT Hackathon '22, '23 & '24, Explainers '22
- Hobbies: reading, traveling, singing & dancing.

8.7 Hongda Zheng

- Contact: zhenghongda7@gmail.com
- Course: 12th grade, 2^o Bachillerato
- Field of work: Engineering
- Experience: Explainers '24, Anem per més Mates
- Hobbies: Playing Tennis, Reading, Mathematics

8.2 Martí Delgado Farriol

- Contact: marti.delgado.farriol@gmail.com
- Course: 2^o Bachillerato (Pureza de María, Sant Cugat del Vallès)
- Field of work: Astronomy & Astrophysics
- Experience: Joves i Ciència '23
- Hobbies: Astrophotography, Chemistry, Garden agriculture and Politics.

8.4 Anna Abadal Garrido

- Contact: talaxar07@gmail.com
- Course: 2^o Bachillerat (Salesians of Sarrià)
- Field of work: Engineering
- Experience: Explainers '24, Dissabtes de la Física '24
- Hobbies: Building models, climbing, chemistry, football and reading.

8.6 Christian Schaefer González

- Contact: chrsch@tinet.org
- Course: 12th grade, 2^o Bachillerato (DSB)
- Field of work: Engineering
- Experience: Jugend Forscht '24
- Hobbies: CAD Design, 3D-printing, rocketry, skiing

8.8 Carles Fonseca

- Contact: carles.fonseca.m@gmail.com
- Course: 12th grade, 2^o Bachillerato (Jesuïtes Sarrià - Sant Ignasi)
- Field of work: Astrophysics, Cosmology, Computation
- Experience: Explainers '22, Joves i Ciència '23
- Hobbies: Playing the piano, basketball, skiing, coding, reading.

9 Conclusion

In conclusion, we have presented a preliminary concept for our mission covering the central aspects for further development and for the participation in the Spark Program. Describing our scientific and technical objectives and outlining the relevance of this mission. Additionally, we have shown potential applications and research within the collaboration, present preliminary instrument design plan, and future plans and timeline. We are confident that with continued refinement and support, this mission will create new opportunities for discovery and fostering a collaborative research environment that bridges educational and scientific achievements, contributing to a deeper understand of the Universe's most energetic phenomena.

References

- 245 D C H J K V R D T L S D V S Y S , 247 L C O C A I H G H D A M C P D V S . 2017. A gravitational-wave standard siren measurement of the hubble constant[J]. *Nature*, 551(7678): 85-88.
- AGHANIM N, AKRAMI Y, ASHDOWN M, et al. 2020. Planck 2018 results - vi. cosmological parameters[J/OL]. *Astronomy Astrophysics*, 641: A6. https://www.aanda.org/articles/aa/full_html/2020/09/aa33910-18/aa33910-18.html<https://www.aanda.org/articles/aa/abs/2020/09/aa33910-18/aa33910-18.html>. DOI: 10.1051/0004-6361/201833910.
- BLANDFORD R D, ZNAJEK R L. 1977. Electromagnetic extraction of energy from kerr black holes[J]. *Monthly Notices of the Royal Astronomical Society*, 179(3): 433-456.
- BLARD P H, BOURLÈS D, LAVÉ J, et al. 2006. Applications of ancient cosmic-ray exposures: Theory, techniques and limitations[J/OL]. *Quaternary Geochronology*, 1(1): 59-73. <https://www.sciencedirect.com/science/article/pii/S1871101406000781>. DOI: <https://doi.org/10.1016/j.quageo.2006.06.003>.
- BRANDT H V, OHASHI T, POUNDS K A. 1992. X-ray astronomy missions[J/OL]. *Annual review of astronomy and astrophysics*, 30: 291-427. <https://articles.adsabs.harvard.edu/full/1992ARA%26A..30..391B/0000391.000.html>.
- CHARBONNEAU D, BROWN T M, NOYES R W, et al. 2002. Detection of an extrasolar planet atmosphere [J/OL]. *The Astrophysical Journal*, 568: 377-384. <https://iopscience.iop.org/article/10.1086/338770><https://iopscience.iop.org/article/10.1086/338770/meta>. DOI: 10.1086/338770/FULLTEXT/.
- COLLABORATION D, ADAME A G, AGUILAR J, et al. 2024. Desi 2024 vi: Cosmological constraints from the measurements of baryon acoustic oscillations[A/OL]. *arXiv*: 2404.03002. <https://arxiv.org/abs/2404.03002>.
- DE ESPECTROSCOPIA GAMMA Y DE NEUTRONES G. 2014. Rayos cósmicos - la radiación del espacio[EB/OL]. http://webgamma.ific.uv.es/gamma/wp-content/uploads/2014/05/Rayos_Cosmicos_poster.pdf.
- DESI Collaboration, Adame A G, Aguilar J, et al. 2024. DESI 2024 VI: Cosmological Constraints from the Measurements of Baryon Acoustic Oscillations[A/OL]. *arXiv*: 2404.03002. *arXiv*: 2404.03002.
- EDITORIAL N A. 2024. The continued growth of exoplanets [J/OL]. *Nature Astronomy* 2024 8:7, 8: 797-798. <https://www.nature.com/articles/s41550-024-02333-5>. DOI: 10.1038/s41550-024-02333-5.
- EFSTATHIOU G. 2024. Evolving dark energy or supernovae systematics?[A/OL]. *arXiv*: 2408.07175. <https://arxiv.org/abs/2408.07175>.
- ESA. The solar wind[M/OL]. European Space Agency. https://www.esa.int/Science_Exploration/Space_Science/Smile/The_solar_wind.
- FREEDMAN W L. 2021. Measurements of the hubble constant: Tensions in perspective*[J/OL]. *The Astrophysical Journal*, 919: 16. <https://iopscience.iop.org/article/10.3847/1538-4357/ac0e95><https://iopscience.iop.org/article/10.3847/1538-4357/ac0e95/meta>. DOI: 10.3847/1538-4357/AC0E95.
- FREEDMAN W L, MADORE B F. 2010. The hubble constant[J/OL]. *Annual Review of Astronomy and Astrophysics*, 48: 673-710. <https://www.annualreviews.org/content/journals/10.1146/annurev-astro-082708-101829>. DOI: 10.1146/ANNUREV-ASTRO-082708-101829/CITE/REFWORKS.
- FREEDMAN W L, MADORE B F, GIBSON B K, et al. 2001. Final results from the hubble space telescope key project to measure the hubble constant[J]. *THE ASTROPHYSICAL JOURNAL*, 553: 47-72.
- FREEDMAN W L, MADORE B F, JANG I S, et al. 2024. Status report on the chicago-carnegie hubble program (cchp): Three independent astrophysical determinations of the hubble constant using the james webb space telescope[A].
- GILLI R. 2013. The cosmic x-ray background: abundance and evolution of hidden black holes[J/OL]. *Mem SAIt*, 84: 1. <https://arxiv.org/abs/1304.3665v1>.
- GRANT C. 2011. CCDs for X-ray Astronomy[M/OL]. https://cxc.cfa.harvard.edu/xrayschool/talks/ccd_talk.pdf.
- GSFC. 2018. Gamma-ray detectors - more information[R]. NASA.
- GSFC N. 2021. Supernova remnants[J/OL]. *Imagine the universe*. https://imagine.gsfc.nasa.gov/science/objects/supernova_remnants.html.
- HESS V F. 1912. Penetrating radiation in seven free balloon flights[J]. *Z Phys*, 13: 1084.
- HUBBLE E. 1929. A relation between distance and radial velocity among extra-galactic nebulae[J/OL]. *Proceedings of the National Academy of Sciences*, 15: 168-173. <https://www.pnas.org/doi/abs/10.1073/pnas.15.3.168>. DOI: 10.1073/PNAS.15.3.168.
- HUŠKO F, LACEY C G, SCHAYE J, et al. 2024. Winds versus jets: a comparison between black hole feedback modes in simulations of idealized galaxy groups and clusters[J]. *Monthly Notices of the Royal Astronomical Society*, 527 (3): 5988-6020.
- ISHIKAWA T. 2019. Accelerator-based x-ray sources: synchrotron radiation, x-ray free electron lasers and beyond [J]. *Philosophical Transactions of the Royal Society A*, 377 (2147): 20180231.
- KUWABARA T, BIEBER J W, CLEM J, et al. 2006. Real-time cosmic ray monitoring system for space weather[J/OL]. *Space Weather*, 4(8). <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2005SW000204>. DOI: <https://doi.org/10.1029/2005SW000204>.

- LEMAÎTRE G. 1927. Un univers homogène de masse constante et de rayon croissant rendant compte de la vitesse radiale des nébuleuses extra-galactiques[J/OL]. *Annales de la Société Scientifique de Bruxelles*, 47: 49-59. <https://articles.adsabs.harvard.edu/pdf/1927ASSB...47...49L>.
- LERNER L. 2024. Cosmic rays, explained[J/OL]. *uchicago news*. <https://news.uchicago.edu/explainer/what-are-cosmic-rays#:~:text=Cosmic%20rays%20are%20particles%20from,and%20made%20harmless%20to%20humans>.
- LUIZ H.G. TIZEI M K. 2017. Chapter four - quantum nano-optics in the electron microscope[J/OL]. Elsevier, 199. DOI: <https://doi.org/10.1016/bs.aiep.2017.01.002>.
- Lyman Spitzer J. 1990. Report to project rand: Astronomical advantages of an extra-terrestrial observatory[J/OL]. *Astronomy Quarterly*, 7(3): 131-142. <https://www.sciencedirect.com/science/article/pii/036492299090018V>. DOI: [https://doi.org/10.1016/0364-9229\(90\)90018-V](https://doi.org/10.1016/0364-9229(90)90018-V).
- MADDALENA F X A A Z S W H C P D W D C D C E A, Tjahjana L. 2019. Inorganic, organic, and perovskite halides with nanotechnology for high-light yield x- and -ray scintillators.[J/OL]. *Crystals*, 9(2). DOI: <https://doi.org/10.3390/cryst9020088>.
- MASCIOCCHI S. 2017. Electromagnetic calorimeters[M/OL]. https://www.physi.uni-heidelberg.de/~sma/teaching/ParticleDetectors2/sma_ElectromagneticCalorimeters.pdf.
- MAYOR M, QUELOZ D. 1995. A jupiter-mass companion to a solar-type star[J/OL]. *Nature* 1995 378:6555, 378: 355-359. <https://www.nature.com/articles/Art1>. DOI: [10.1038/378355a0](https://doi.org/10.1038/378355a0).
- Morison I. 2008. Introduction to Astronomy and Cosmology [M].
- PEEBLES P J E, RATRA B. 2003. The cosmological constant and dark energy[J/OL]. *Reviews of Modern Physics*, 75(2): 559-606. <http://dx.doi.org/10.1103/RevModPhys.75.559>. DOI: [10.1103/revmodphys.75.559](https://doi.org/10.1103/revmodphys.75.559).
- PIERA L G. 2012. COSMIC RAY DETECTION[C/OL]// https://www2.ung.si/~sstanic/teaching/APP/P_Ghia_CosmicRayDetection_I.pdf.
- RIESS A G, FILIPPENKO A V, CHALLIS P, et al. 1998. Observational evidence from supernovae for an accelerating universe and a cosmological constant[J]. *The astronomical journal*, 116(3): 1009.
- RIESS A G, ANAND G S, YUAN W, et al. 2024a. Jwst observations reject unrecognized crowding of cepheid photometry as an explanation for the hubble tension at 8 confidence [EB/OL]. <https://doi.org/10.3847/2041-8213/ad1ddd>.
- RIESS A G, SCOLNIC D, ANAND G S, et al. 2024b. Jwst validates hst distance measurements: Selection of supernova subsample explains differences in jwst estimates of local h_0 [EB/OL]. <https://arxiv.org/abs/2408.11770v1>.
- ROYON C, ISIDORI T, MINAFRA N. 2019. Cross-disciplinary research: from nuclear physics to cosmic ray detection and medical applications[J]. *Merrill Series on The Research Mission of Public Universities*: 54-60.
- SAGANTI P B, CUCINOTTA F A, WILSON J W, et al. 2004. Radiation climate map for analyzing risks to astronauts on the mars surface from galactic cosmic rays[J]. *2001 Mars Odyssey*: 143-156.
- Schaller M, et al. 2024. SWIFT: A modern highly-parallel gravity and smoothed particle hydrodynamics solver for astrophysical and cosmological applications[J/OL]. , 530(2): 2378-2419. DOI: [10.1093/mnras/stae922](https://doi.org/10.1093/mnras/stae922).
- SCHNEIDER P. 2006. Extragalactic astronomy and cosmology: An introduction[J/OL]. *Extragalactic Astronomy and Cosmology: An Introduction*: 1-459. DOI: [10.1007/978-3-540-33175-9/COVER](https://doi.org/10.1007/978-3-540-33175-9/COVER).
- Singer L, Racusin J. 2023. General Coordinates Network (GCN): NASA's Next Generation Time-Domain and Multimessenger Astronomy Alert System[C]//*American Astronomical Society Meeting Abstracts: Vol. 241 American Astronomical Society Meeting Abstracts*. 108.02.
- TELEDYNE. 2024a. Silicon-based ccids: The basics[R]. *Teledyne Vision Solutions*.
- TELEDYNE. 2024b. Direct Detection of X-rays (30 eV to 20 keV) Using Detectors Based on CCD Technology[R/OL]. *Teledyne Princeton Instruments*. <https://www.teledynevisionsolutions.com/es-ES/learn/learning-center/scientific-imaging/direct-detection-of-x-rays-30-ev-to-20-kev-using-detectors-based-on-ccd-t>
- TOZZI P, ROSATI P, NONINO M, et al. 2001. New results from the x-ray and optical survey of the chandra deep field-south: The 300 kilosecond exposure. ii.[J/OL]. *The Astrophysical Journal*, 562: 42-51. <https://iopscience.iop.org/article/10.1086/322492https://iopscience.iop.org/article/10.1086/322492/meta>. DOI: [10.1086/322492/FULLTEXT/](https://doi.org/10.1086/322492/FULLTEXT/).
- VALENTINO E D, MELCHIORRI A, SILK J. 2019. Planck evidence for a closed universe and a possible crisis for cosmology[Z].
- VALENTINO E D, MENA O, PAN S, et al. 2021. In the realm of the hubble tension—a review of solutions *[J/OL]. *Classical and Quantum Gravity*, 38: 153001. <https://iopscience.iop.org/article/10.1088/1361-6382/ac086dhttps://iopscience.iop.org/article/10.1088/1361-6382/ac086d/meta>. DOI: [10.1088/1361-6382/AC086D](https://doi.org/10.1088/1361-6382/AC086D).
- WOLSZCZAN A, FRAIL D A. 1992. A planetary system around the millisecond pulsar psr1257 + 12[J/OL]. *Nature* 1992 355:6356, 355: 145-147. <https://www.nature.com/articles/355145a0>. DOI: [10.1038/355145a0](https://doi.org/10.1038/355145a0).
- ZEITLIN C. 2012. Cosmic ray detectors: Principles of operation and brief overview of (mostly) u.s. flight instruments[M/OL]. *NASA*. <https://three.jsc.nasa.gov/articles/CosmicRayDetectors.pdf>.