

## Expert Opinion

## Nebivolol, a third generation beta-blocker

John Cruickshank

Independent Cardiovascular Consultant, Long Melford, Great Britain.

## Abstract

Nebivolol is a third generation beta-blocker, which is a cardiac beta-1 receptor antagonist and an endothelial beta-3 receptor agonist (beta-3 intrinsic sympathomimetic activity, or ISA). It is highly lipophilic and is metabolised in the liver. Like other beta-blockers with beta-2 or beta-3 ISA, nebivolol is a vasodilator and antioxidant as a result of endothelial nitric oxide (NO) release. Peripheral and aortic central pressures are reduced via a fall in total peripheral resistance. Hard end-point data exist only for heart failure. Unlike beta-blockers without ISA (bisoprolol, metoprolol, carvedilol), which reduce resting heart rate by about 14 beats per minute (bpm) and all-cause death by about 35% in patients with systolic heart failure (reduced ejection fraction), nebivolol (in common with other beta-blockers with ISA eg bucindolol and xamoterol) reduces resting heart rate by about 8–9 bpm and all cause mortality by about 17%. Nebivolol also reduces all-cause mortality in diastolic heart failure (normal ejection fraction) by about 17%. Thus, nebivolol behaves like a typical beta-blocker with beta-2 or beta-3 ISA.

**Keywords:** beta-3 intrinsic sympathomimetic activity; nitric oxide; anti-oxidant activity; systolic heart failure; diastolic heart failure.

**Received:** February 13, 2014; **Accepted:** June 5, 2014; **Published:** December 16, 2014

**Corresponding Author:** John Malcolm Cruickshank, BM, BCH, MA (Oxford), DM (Oxford), FRCP (London). **E-mail:** [Johndtl@aol.com](mailto:Johndtl@aol.com).

## Introduction

A whole 50 years has passed since the first beta-blocker, propranolol, appeared on the market [1]. Propranolol, a so-called first generation beta-blocker (BB) which blocks both beta-1 and beta-2 receptors (non-selective), was followed by second-generation BBs such as atenolol and metoprolol which preferentially block beta-1 (over beta-2 receptors), i.e. so-called cardioselective or beta-1 selective BBs. Then came the third-generation BBs, with additional vasodilatory effects i.e. carvedilol, labetalol, celiprolol, bucindolol, and nebivo-


lol. Nebivolol's particular claim was to be highly beta-1 selective, with vasodilatory and antioxidant properties, involving nitric oxide (NO) release i.e. it was unique[2].

The purpose of this review is to examine the clinical pharmacology of nebivolol, and to assess its role in cardiovascular medicine.

## Clinical Pharmacology

*Pharmacokinetics*

The pharmacokinetic properties of nebivolol have been described by Prisant [3] and Cruickshank [4].



**Figure 1.** Plasma levels of D-nebivolol (beta-1 blockade) in extensive (EM) and poor (PM) metabolisers, after 3 months therapy. Lefebvre J, *et al.* 2006.

### Metabolism

Nebivolol is highly lipophilic and, like metoprolol and timolol, is heavily metabolised in the liver via the cytochrome P4502D6 enzyme system. This enzyme system is vulnerable to genetic polymorphism, involving poor (PM) and extensive (EM) metabolisers, resulting in high variation in plasma concentrations of nebivolol [5]. Thus, mean steady state plasma concentrations of D- and L-nebivolol are 10- and 12-fold greater in PMs than EMs (**Figure 1**).

This highly significant difference ( $P < 0.0001$ ) in plasma levels was, surprisingly, not associated with differences in blood pressure control or adverse reactions (though beta-1 selectivity, vs the beta-2 receptor, will be markedly diminished at the high plasma levels observed in PMs).

Poor metabolisers (PM) account for about 8–10% of the UK white population [6], and possibly 30% of Chinese [7]. Nebivolol is rapidly absorbed, and food does not modify this process [3]. Peak plasma concentrations occur at about 2 hours, with a half-life of 11 hours. Thus once daily dosing is appropriate.

### Modifying Factors

The volume of distribution of lipid-soluble BBs is significantly decreased in obese patients [3], indicating that such agents diffuse less extensively into fatty, rather than lean, tissue. However, obesity does not result

in significant differences in the pharmacokinetic profile of nebivolol [3].

In patients with severe hepatic impairment, e.g. cirrhosis, metabolism of nebivolol is diminished, so that blood levels of the drug are markedly increased (with subsequent loss of beta-1 selectivity and the potential for increased adverse reactions [3]).

In patients with renal dysfunction, dosage reduction with nebivolol is generally not necessary [3].

### Drug Interactions

There is the potential for drug-interaction with agents that are metabolised via the same liver cytochrome P4502D6 system as nebivolol. Thus, the anti-ulcer drug cimetidine, when co-administered with nebivolol, induces a significant increase in plasma levels of unchanged nebivolol, but no change in heart rate or blood pressure [3]. Other drugs which are metabolised in a similar fashion to nebivolol, e.g. the ARB losartan, the ACE-I ramapril and furosemide, did not significantly modify plasma levels of nebivolol [3]. Likewise there are no kinetic interactions between nebivolol and digoxin, spironolactone and warfarin [3].

### Pharmacodynamics


#### *Nebivolol's d- and l- enantiomers*

Nebivolol is a racemic mixture of 2 enantiomers, d- and l-nebivolol, each associated with distinct pharmacological properties [8–10]. The d-enantiomer occupies the cardiac beta-1 receptor in a highly selective manner, while the l-enantiomer occupies sites(s) in the vascular endothelium, resulting in nitric oxide (NO) release, anti-oxidant effects, and vasodilation [8–9], and a fall in blood pressure via a reduction in total peripheral resistance (TPR) [10].

#### *Is Intrinsic Sympathomimetic Activity (ISA) involved in the Vasodilatory Action of Nebivolol?*

Early studies in animals [11] and human cardiac tissue [12–13] indicated that nebivolol possessed no ISA. Others suggested that nebivolol possessed beta-2 ISA, involving NO production [14].

However, later studies involving modern technology [15–18] indicated that vasodilatation induced by nebivolol was via beta-3 agonism i.e. beta-3 ISA.



**Figure 2.** Effect of nebivolol (on left) and atenolol on systemic vascular resistance in hypertension [23].

#### *Beta-3 ISA and Nitric Oxide Release*

The beta-3 receptor is widespread throughout the body [19], occurring in the brain, adipose tissue, gut, liver and partial circulation, urinary bladder, blood vessels (endothelium), myocardium and myometrium. Stimulation of the cardiac beta-3 receptor has negative inotropic effects, and stimulation of endothelial beta-3 receptors induces vasorelaxation [19]. Vasodilation (in humans) via beta-3 stimulation is associated with endothelial NO release [20]. Nebivolol induces vasodilation in this fashion [3, 21], resulting in a fall in forearm vascular resistance [22] (**Figure 2**). In hypertensive patients, compared to atenolol which tends to increase total peripheral resistance, nebivolol effects a significant 13% reduction in peripheral resistance [23].

#### *Beta-3 Stimulant Effects upon the Myocardium*

In hypertensive patients [23], compared to atenolol which causes a significant 24% reduction in cardiac output, nebivolol causes a significant 8% increase in ejection fraction and significant 21% increase in stroke volume. However, in both the non-failing [17] and failing [24] human heart, beta-3 stimulation induces an NO-dependent negative inotropic effect.

#### *Nebivolol and Antioxidant Effects*

Oxidative stress is a process whereby NO synthase (NOS) is uncoupled, so that instead of NO production there is production of peroxynitrite, a reactive oxygen species (ROS) which is toxic to the vasculature [25].

*In vitro* and *in vivo* studies have indicated that nebivolol scavenges ROS, resulting in improved endothelial function [26–27]. In hypertensive patients, in contrast to atenolol, nebivolol significantly lowered levels of ROS and superoxide concentration in the plasma and endothelial cells [28].

It has been suggested that the above action of nebivolol could lead to cardioprotective effects [3].

#### *How beta-1 Selective is Nebivolol?*

Literally speaking nebivolol is not beta-1 selective, as it occupies both the beta-1 and beta-3 receptors. However, beta-1 selectivity normally relates to the beta-1 and beta-2 occupancy. On that basis, nebivolol is highly beta-1 selective. In one study [13] nebivolol was somewhat less beta-1 selective than bisoprolol, but in 3 others was somewhat more selective than bisoprolol and other beta-blockers [12, 29–30]. The reason for these disparities could be explained by the use of different radioligands, or by experimental fluctuations.

#### *So how “unique” is the Pharmacological Profile of Nebivolol?*

##### *Haemodynamics*

There are BBs, other than nebivolol, that exert beta-blockade plus vasodilation.

Carvedilol exerts both beta- and alpha blockade. However, on chronic therapy there is evidence suggesting that alpha-blockade-induced vasodilatation is no longer present, due to tachyphylaxis [31–32].

Pindolol exerts both beta-1 and beta-2 blockade, and beta-2 ISA. Thus, on chronic therapy pindolol induces a significant increase in cardiac output and a significant fall in vascular resistance [33] i.e. a similar haemodynamic profile to nebivolol.

##### *Beta-3 ISA*

Pindolol has been reported to possess no beta-3 ISA in studies involving rat ileum [34] and human atrium [35]. However, others have demonstrated beta-3 agonism in human fat cells [36] and hamster ovaries [37].

Bucindolol has been shown to stimulate beta-3 receptors in hamster ovaries [37].

Oxprenolol has been shown to be an agonist of beta-3 receptors in human fat cells [36].

Alprenolol, at higher doses, possesses beta-3 agonist properties hamster ovarian cells [38].

Labetalol acts as a partial agonist on beta-3 adrenoceptors in the guinea-pig stomach [39].

### *Endothelial Nitric Oxide (NO) Release and beta-adrenoceptor Stimulation*

As already indicated [21], the vasodilatory effect of nebivolol is associated with NO release. However, nebivolol is not the only vasodilatory beta-blocker that is associated with this phenomenon.

Thus, pindolol (beta-2 and beta-3 ISA) [40], bopindolol (beta-2 ISA) [41], oxprenolol (beta-2 ISA) [42] and celiprolol (beta-2 ISA) [41, 43–45], are all beta-blockers with NO-dependent vasodilatory properties.

It is thus apparent that stimulation of beta-1, beta-2 and beta-3 receptors results in NO-release [40]. Hence, beta-stimulants such as isoprenaline [40], salbutamol [40, 46–48], clenbuterol [47] and albuterol [49], induced vasodilation involving NO-release. With salbutamol, tracheal relaxation is also associated with release of NO [50].

### *Reduction in Oxidative Stress*

As already noted [28], in hypertensive subjects nebivolol displays antioxidant properties. Such properties are not possessed by BBs without ISA, e.g. metoprolol [28] or atenolol [51].

However, nebivolol is not unique as a BB which displays anti-oxidant properties. Celiprolol, with beta-2 ISA, stimulates nitric oxide release and exhibits anti-oxidant properties [52]. Carvedilol, a non-selective beta-blocker with alpha-1 blocking vasodilating properties, also displays anti-oxidant properties in both animals [53–54] and man [55–56].

Is the possession of anti-oxidant properties, in addition to beta-blockade, useful clinically? There are suggestions that the answer may be yes [57–58]. However, this topic will be discussed later in the clinical section.

**Table 1.** Acute post-myocardial patients, randomised to placebo vs L-arginine; trial stopped due to excess deaths in L-arginine group.

|                  | Placebo | L-Arginine (%) | Pvalue |
|------------------|---------|----------------|--------|
| <b>Mortality</b> | 0.0     | 8.6            | 0.01   |

From Schulman SP, Becker LC, Kass DA. L-Arginine therapy in acute myocardial infarction. JAMA. 2006; 295: 58-64.

Certainly, on the general (non-BB) front, there are concerns about nitric oxide and anti-oxidants. While the dangers of oxidative stress are well recognised in diabetes [59], cancer progression [60], and post coronary artery interventions [61], there is recognition of an anti-oxidant paradox [62–63]. Thus, there are scenarios when an excess of NO aggravates the atheromatous process [64–65]; a worrying thought in view of the high level of antioxidant supplement intake in countries like the USA [66]. Likewise, the fact that in the post-myocardial infarction period, L-arginine (substrate for NO synthase) significantly increased mortality vs randomised placebo [67] (**Table 1**), is of concern.

The results of large randomised trials involving anti-oxidants give little reason for optimism. Meta-analyses of placebo-controlled trials involving antioxidants concluded that beta-carotene, vitamin A, and vitamin E, were all associated with significant increases in mortality [68]! In the randomised, double-blind, placebo-controlled Physicians' Health Study [69], vitamin E was associated with a significant increased risk of haemorrhagic stroke. Thus, while anti-oxidant vitamins do not decrease [70], and may increase [71], cardiovascular events and mortality, a further concern is the possibility that cancer-risk may be increased [72].

### *So is the Pharmacological Profile of Nebivolol Unique?*

The simple answer is no. Other vasodilating BBs, such as pindolol, increase cardiac output and lower peripheral resistance, as does nebivolol. Other vasodilating BBs, such as pindolol, bucindolol, oxprenolol, alprenolol and labetalol, have some beta-3 stimulatory properties associated with release of NO. Vasodilatory BBs like carvedilol and celiprolol, possess anti-oxidant properties.

**Table 2.** Morphological changes in heart failure with normal ejection fraction (HFNEF) and reduced ejection fraction (HFREF).

| Parameters  | HFNEF                         | HFREF               |
|---|-------------------------------|---------------------|
| <b>Microscopic and Neuroendocrine Features</b>  |                               |                     |
| Cardiac cell hypertrophy  | Increased                     | Minimal             |
| Resting tension of cardiomyocytes   | Increased                     | Decreased           |
| Myofilament density   | Preserved                     | Decreased           |
| Titin N28/N28A ratio  | Increased                     | Decreased           |
| Interstitial collagen   | Increased                     | Decreased           |
| MMP-1/TIMP-1 ratio  | Little changed                | Decreased           |
| Patterns of peptide growth factor induction   | Different                     | Different           |
| $\beta$ -Receptor down-regulation and/or myocardial $\beta$ -adrenergic receptor desensitization (postsynaptic) | Present                       | Present             |
| Norepinephrine  | Increased                     | Increased           |
| B-type natriuretic peptide  | Increased                     | More increased      |
| <b>Resting Echocardiographic Parameters</b>   |                               |                     |
| LV cavity size  | Normal or decreased           | Increased           |
| LV shape and geometry   | Little changed                | Spherical           |
| LV mass index   | Increased                     | Increased           |
| LV mass to cavity ratio   | Increased                     | Normal or decreased |
| Relative wall thickness   | Increased                     | Normal              |
| End-diastolic volume/wall stress  | Normal or decreased           | Increased           |
| End-systolic volume/wall stress   | Normal                        | Increased           |
| LV ejection fraction  | Normal                        | Decreased           |
| Longitudinal velocity/strain  | Decreased                     | More increased      |
| Radial strain   | Decreased                     | More increased      |
| LV twist (torsion)  | Normal or decreased           | Decreased           |
| LV twisting rate  | Normal or decreased           | Decreased           |
| LV untwisting rate  | Normal or decreased           | Decreased           |
|   | with/without<br>delayed onset |                     |

HFNEF, heart failure with normal ejection fraction; HF, heart failure with reduced ejection fraction; LV, left ventricular; MMP-1, matrix metalloproteinase-1; TIMP-1, tissue inhibitor of metalloproteinases-1. From Yip GW-K, Frenneaux M, Sanderson JE. *Heart failure with a normal ejection fraction: new developments. Heart. 2009; 95: 1549-1552.*

## Clinical Aspects

### Heart Failure

#### Types of Heart Failure

There are 2 basic types of heart failure, one termed systolic heart failure, with a reduced ejection fraction (HFREF), and diastolic heart failure, with a normal ejection fraction (HFNEF). **Table 2** shows the characteristics of patients with either type of heart failure [73]. HFREF comprises about 50% of all heart failure, and 65% of HFREF is associated with coronary artery disease [74]. In contrast, a typical case of HFNEF would be an elderly female, with a history of systolic hyper-

tension, concentric left ventricular hypertrophy – LVH (as opposed to eccentric LVH in HFREF), with a small left ventricular cavity [75].


#### Beta-blockers and Mortality in Heart Failure with Reduced Ejection Fraction (HFREF)

Not all BBs are equal in their ability to reduce all-cause mortality in patients with HFREF [4, 76] (**Figure 3**). It is apparent that the 3 BBs without ISA, i.e. bisoprolol [77], metoprolol slow release [78] and carvedilol [79], reduced all-cause mortality significantly by about 35%, accompanied by significant reductions of heart rate of 13–14 bpm. By contrast, the 3 BBs with ISA had no




significant benefit concerning reduction of all-cause death. Xamoterol, with a high level of beta-1 ISA, actually increased all-cause death vs placebo [80]. Bucindolol, with beta-2 ISA [81] and nebivolol, with beta-3 ISA [82], lowered heart rate by 8–9 bpm, and reduced all-cause death by a non-significant 10–12%.

The SENIORS trial has been heavily criticised [83–84]. After exclusion of patients with HFNEF (diastolic heart failure), it was concluded that compared to placebo nebivolol reduced the primary end-point by only 16–17%, being about half the efficacy of bisoprolol [77], metoprolol [78] and carvedilol [79] in HFREF.



**Figure 3.** Beta-blockers and hard end-point placebo-controlled trials in heart failure (on ACEIs); ISA (xamoterol, bucindolol, and nebivolol) reduces fall in heart rate (HR) and efficacy (all-cause death) compared to beta-blockers with no ISA.



**Figure 4.** Cardiovascular mortality in relation to changes in heart rate in chronic systolic heart failure.

Cook S, *et al.* 2006.

However, it has been pointed out that a) about one third of patients in SENIORS had HFNEF, and that higher doses of nebivolol resulted in a 25% reduction in all cause death or cardiovascular hospitalization [85], and b) in patients with an ischaemic aetiology of their heart failure, ischaemic events were reduced by about one third, with no benefit to non-ischaemic patients [86].

### *Heart Failure and the Heart-rate Factor*

In HFREF studies, there is a clear relationship between change in heart rate and all-cause mortality [87] (**Figure 4**), with best results arising at heart rates less than 71 bpm [88]. Such issues are relevant when considering the benefits of adding the sino-atrial blocker ivabradine to beta-blocker therapy in heart failure [89].

### *Beta-blockers and Heart Failure with Normal Ejection Fraction (HFNEF)*

It has been stated that no treatments benefit patient with HFNEF [90, 91]. However, nebivolol makes a claim to be the only beta-blocker to benefit patients with HFNEF [92], as does carvedilol at higher dosage [93]. Retrospective [94] and observational studies claim that BBs are effective in reducing death or hospitalization in HFNEF. Certainly in post myocardial infarction cases of HFNEF, BBs were protective [95, 96].

### *Hypertension*

There have been no “hard end-point” studies with nebivolol, in patients with essential hypertension.

### *Effect of Nebivolol upon Peripheral Blood Pressure, and Side Effects*

Age and race are important regarding antihypertensive efficacy of BBs. When plasma renin is low, as in elderly and Black subjects, traditional BBs are less effective in lowering blood pressure [96, 97]. While it has been stated that BBs like bisoprolol [98] and nebivolol [99] are equally effective in lowering blood pressure in elderly and Black subjects, others [100] state that nebivolol is less effective at lowering blood pressure in the elderly.

In younger/middle-aged hypertensive subjects, nebivolol is superior to randomised placebo in lowering blood pressure [101–102]. Nebivolol displayed an anti-

hypertensive action similar to atenolol [103], metoprolol [104], bisoprolol [105] and carvedilol [106].

In the elderly nebivolol lowered blood pressure to a similar degree as amlodipine [107].

**Side Effects** – The side effect profiles of highly beta-1 selective nebivolol and bisoprolol are similar [105]. However, compared to only moderately beta-1 selective atenolol [103, 108] and metoprolol [109], there is no metabolic disturbance involving lipids, blood glucose and insulin-sensitivity, with nebivolol. Compared to amlodipine [107] nebivolol is better tolerated. Sexual dysfunction is more likely with metoprolol than nebivolol [110]; in this respect nebivolol is similar to bisoprolol [111]. Occasional cases of skin lichenoid eruption occur with nebivolol [112]. Walking distance in hypertensive patients with peripheral arterial disease is not altered by nebivolol, and in this respect does not differ from metoprolol [113] or thiazide diuretics [114].

### *Nebivolol and Central Blood Pressure*

Central pressures are important, as they are possibly the best predictors of future cardiovascular events [115–117].

Not all BBs decrease central pressure. Only vasodilatory BBs, that decrease pressure wave reflection from the periphery, lead to a decrease in central aortic blood pressure [118]. This point was well illustrated in double-blind, randomised, crossover study in middle-aged hypertensive subjects where the BB dilevalol (with beta-2 ISA) was compared to atenolol [119]. While both BBs lowered brachial blood pressure to a similar degree, dilevalol was significantly more effective than atenolol in lowering central SBP and augmentation index. Likewise in elderly hypertensive subjects, atenolol has proved to be ineffective in lowering central aortic pressures, compared to ACE-inhibitors, diuretics and calcium blockers [120].

Nebivolol is relatively effective in reducing central aortic pressure. Under both open [121] and randomised, placebo-controlled conditions [122], nebivolol has been shown to be effective in lowering central systolic, diastolic, and pulse-pressure. Compared to atenolol [123, 124] and metoprolol [125–127], nebivolol has more favourable effects on central pressures. However, once diuretics are added to BB therapy, the advantage of nebivolol over metoprolol in reducing central aortic pressure disappears [128].


### *Is Nebivolol Unique, as a Beta-blocker in Lowering Central Aortic Pressures?*

The simple answer is no. As indicated above, dilevalol, a BB with beta-2 ISA, was effective in lowering central aortic pressures [119]. Also celiprolol, a BB with beta-2 ISA, had favourable effects upon central pressures [129]. Carvedilol, a vasodilating BB with alpha-blocking properties, is superior to atenolol in reducing central aortic pressure [130]. Indeed, inhalation of the beta-2 stimulant salbutamol, results in fall in central systolic pressure [131].


Interestingly, in young/middle-aged hypertensive subjects, even a highly beta-1 selective BB with no ISA or alpha-blocking properties i.e. bisoprolol, was, unlike atenolol, found to possess positive arterial distensibility/compliance properties leading to a significant fall in central arterial systolic pressure [132].

### *Heart Rate*

Nebivolol reduces resting heart less than BBs without ISA. For example, in middle-aged hypertensive subjects, nebivolol was similar to atenolol in lowering blood pressure, but reduced heart rate significantly less [108]. This may have prognostic implications, as the Framingham Group have shown that in untreated male hypertensive subjects followed-up for 36 years, the best results in terms of all-cause death, coronary heart disease and cardiovascular events were at lower heart rates [133] (**Figure 5**).



**Figure 5.** Framingham: Effect of resting heart rate on all-cause death, CHD and CVD events in untreated male hypertensives, followed-up for 36 years. Gillman MW, *et al.* 1993.



**Figure 6.** Beta-blockers, oral 4 days post MI – best mortality reduction with low heart rates.

### Ischaemic Heart Disease

There are no hard endpoint data in this area regarding nebivolol.

Nebivolol may not be the ideal BB to prescribe in the post myocardial infarction period. As indicated earlier, in the post-infarction period, L-arginine (substrate for NO synthase) significantly increased mortality vs randomised placebo [67] (**Table 1**). Also, nebivolol lowers resting heart rate less than a classic BB like atenolol [134]. This may be important, as reduction in mortality post myocardial infarction relates to reduction in heart rate, with BBs possessing ISA i.e. pindolol and oxprenolol being the least effective [135] (**Figure 6**). A similar story holds for non-fatal re-infarction [136].

### Summary and Conclusions

Nebivolol is a third-generation beta-blocker (BB) with beta-3 intrinsic sympathomimetic activity (ISA). Its d-isomer occupies the cardiac beta-1 receptor, effecting beta-1 blockade, while the l-isomer occupies cardiac and vascular beta-3 receptors, the latter effecting vasodilatation. Nebivolol is metabolised via the hepatic P4502D6 enzymatic system, so that in slow-metabolisers (comprising about 10% of the UK population) blood levels are increased 10–20 fold (with loss of beta-1 selectivity vs the beta-2 receptor).

Resting heart rate is lowered to a lesser degree with nebivolol than with BBs without ISA e.g. atenolol, metoprolol, bisoprolol (this may have prognostic implications). Blood pressure decreases via a fall in total peripheral resistance (as with other vasodilating BBs with ISA, like pindolol and celiprolol). In common with

other BBs with ISA, nebivolol decreases central aortic pressure, is associated with nitric oxide (NO) release, and possesses anti-oxidant properties. Thus, as is sometimes claimed, nebivolol does not possess a unique pharmacological profile.

Hard endpoint data with nebivolol exist only for heart failure. For patients with systolic heart failure (low ejection fraction) nebivolol, like other BBs with ISA (i.e. xamoterol and bucindolol) was associated with a non-significant reduction in all-cause death. These results compare poorly with the results of BBs without ISA, i.e. carvedilol, metoprolol, and bisoprolol, that reduce the risk of all-cause death by a significant 34–35%. Patients with diastolic heart failure (normal ejection fraction) benefited to a degree from nebivolol, with a fall in all-cause death of 17–18%.

Thus, nebivolol is similar to other BBs with ISA in terms of its pharmacodynamic profile, its effect on central blood pressure, and in its relatively poor results (vs BBs without ISA) in reducing all-cause death in patients with systolic heart failure.

### Disclosure

There are no conflicts of interest.

### References

1. Black JW, Crowther AF, Shanks RG, Smith LH, Dornhorst AC. A new adrenergic beta-receptor antagonist. *Lancet*. 1964; 1: 1080-1081.
2. Ignarro LJ. Experimental evidences of nitric oxide-dependent vasodilatory activity of nebivolol, a third-generation beta-blocker. *Blood Pressure*. 2004; 13(1): 2-16.
3. Prisant LM. Nebivolol: Pharmacologic profile of an ultrasensitive, vasodilatory beta-1 blocker. *J Clin Pharmacol*. 2008; 48: 225-239.
4. Cruickshank JM. The modern role of beta-blockers in cardiovascular medicine. People's Medical Publishing House-USA. Shelton, Connecticut 2010.
5. Lefebvre J, Poirier L, Poirier P, Turgeon J, Lacourciere Y. The influence of CYP2D6 phenotype on the clinical response of nebivolol in patients with essential hypertension. *Br J Clin Pharmacol*. 2007; 63: 575-582.
6. Evans DA, Mahgoub A, Sloan TP, et al. A family and population study of the genetic polymorphism of debrisoquine oxidation in a white British population. *J Med Genet*. 1980; 17: 102-105.
7. Karlow W. The metabolism of xenobiotics in different populations. *Canad J Physiol Pharmacol*. 1982; 60: 1-12.
8. Kuroedov A, Cosentino F, Luscher TF. Pharmacological mechanisms of clinically favourable properties of a selective beta-1 adrenoceptor antagonist, nebivolol. *Cardiovasc Drug Rev*. 2004; 22: 155-168.



9. Van Nueten L, De Cree J. Nebivolol: comparison of the effects of dl-nebivolol, d-nebivolol, l-nebivolol, atenolol and placebo on exercise-induced increases in heart rate and systolic blood pressure. *Cardiovasc Drugs Ther.* 1998; 12: 339-334.
10. Xhonneax R, Wouters L, Reneman RS, Janssen PA. The l-enantiomer of nebivolol potentiates the blood pressure lowering effect of the d-enantiomer. *Eur J Pharmacol.* 1990; 181: 261-265.
11. Janssens WJ, Van de Water A, Xhonneax R, Reneman RS, Van Nueten JM, Janssen PA. Nebivolol is devoid of intrinsic sympathomimetic activity. *Europ J Pharmacol.* 1989; 159L89-95.
12. Brixius K, Bundkirchen A, Bolek B, Mehlhorn U, Schwinger HG. Nebivolol, bucindolol, metoprolol and carvedilol are devoid of intrinsic sympathomimetic activity in human myocardium. *Br J Pharmacol.* 2001; 133: 1330-1338.
13. Maack C, Tyroller P, Schnabel P, et al. Characterisation of beta-1 selectivity, adrenoceptor-Gs-protein interaction and inverse agonism of nebivolol in human myocardium. *Br J Pharmacol.* 2001; 132: 1817-1826.
14. Broeders MA, Doevendans PA, Bekkars CA, et al. Nebivolol a third-generation beta-blocker that augments vascular nitric oxide release. *Circulation.* 2000; 102: 677-684.
15. Dessy C, Saliez J, Ghisda P et al. Endothelial beta-3 adrenoceptors mediate nitric oxide-dependent vasorelaxation of coronary microvessels in response to the third-generation beta-blocker nebivolol. *Circulation.* 2005; 112: 1198-1205.
16. Rozec B, Quang TT, Noireaud J, Gauthier C. Mixed beta-3 adrenoceptor agonist and adrenoceptor antagonist properties of nebivolol in rat thoracic aorta. *Brit J Pharmacol.* 2006; 147: 699-706.
17. Rozec B, Erfanian M, Laurent K, Trochu J-N, Gauthier C. Nebivolol, a vasodilating selective beta-1 blocker, is a beta-3 adrenoceptor agonist in the non-failing transplanted human heart. *J Am Coll Cardiol.* 2009; 53: 1532-1538.
18. Feng MG, Prieto MC, Navar LG. Nebivolol induced vasodilatation of renal afferent arterioles involves beta-3 adrenergic receptor and nitric oxide synthase activation. *Am J Physiol Renal Physiol.* 2012; 303: 775-782.
19. Ursino MG, Vasina V, Raschi E, Crema F, De Ponti F. The beta-3 adrenoceptor as a therapeutic target: current perspectives. *Pharmacol Res.* 2009; 59: 221-234.
20. Dessy C, Moniotte S, Ghisda P, Havaux X, Noirhomme P, Balligand JL. Endothelial beta-3 adrenoceptors mediate vasorelaxation of human coronary micro-arteries through nitric oxide and endothelium-dependent hyperpolarization. *Circulation.* 2004; 110: 948-954.
21. Bakris GL, Basile JN, Giles TD, Taylor AA. The role of nitric oxide in improving endothelial function and cardiovascular health: focus on nebivolol. *Am J Med.* 2010; 123: S 2-8.
22. Stoleru L, Wijns W, van Eyck C et al. Effects of d-nebivolol on left ventricular systolic and diastolic function: comparison with d-l nebivolol and atenolol. *J Cardiovasc Pharmacol.* 1993; 22: 183-190.
23. Kamp O, Sieswerda GT, Visser CA. Comparison of effects on systolic and diastolic left ventricular function of nebivolol versus atenolol in patients with uncomplicated essential hypertension. *Am J Cardiol.* 2003; 92: 344-348.
24. Napp A, Brixius K, Pott C, et al. Effects of the beta-3 adrenergic agonist BRL 37344 on endothelial nitric oxide synthase phosphorylation and force of contraction in human failing myocardium. *J Card Fail.* 2009; 15: 57-67.
25. Cai H, Harrison DG. Endothelial dysfunction in cardiovascular diseases: the role of oxidant stress. *Circ Res.* 2000; 87: 840-844.
26. Mollnau H, Schulz E, Daiber A, et al. Nebivolol prevents vascular NOS 111 uncoupling in experimental hyperlipidemia and inhibits NADPH oxidase activity in inflammatory cells. *Arterioscler Thromb Biol.* 2003; 6: 615-621.
27. de Groot AA, Mathy M-J, Van Zwieten PA, Peters SL. Antioxidant activity of nebivolol in the rat aorta. *J Cardiovasc Pharmacol.* 2004; 43: 148-153.
28. Fratta Pasini A, Garbin U, Nava MC, et al. Nebivolol decreases oxidative stress in essential hypertensive patients and increases nitric oxide by reducing its oxidative inactivation. *J Hypertens.* 2005; 23: 589-596.
29. Bundkirchen A, Brixius A, Bolck B, Nguyen Q, Schwinger RG. Beta-adrenoceptor selectivity of nebivolol and bisoprolol. A comparison of (3H) CGP 12,177 and (125) iodocyanopindolol binding studies. *Eur J Pharmacol.* 2003; 460: 19-26.
30. Bristow MR, Nelson P, Minobe W, Johnsson C. Characterisation of beta-1 adrenergic receptor selectivity of nebivolol and various other beta-blockers in human myocardium. *Am J Hypertens.* 2005; 18(2): 51A-52A.
31. Kubo T, Azevedo ER, Newton GE, Parker JD, Floras JS. Lack of evidence for peripheral alpha-1 adrenoceptor blockade during long-term treatment of heart failure with carvedilol. *J Am Coll Cardiol.* 2001; 38: 463-469.
32. Hryniewicz K, Androne AS, Hudaihed A, Katz SD. Comparative effects of carvedilol and metoprolol on regional vascular responses to adrenergic stimuli in normal subjects and patients with chronic heart failure. *Circulation.* 2003; 108: 971-976.
33. Man in't Veld AJ, Schalekamp AD. Effects of ten different beta-blockers on hemodynamics, plasma renin activity and plasma norepinephrine in hypertension. *J Cardiovasc Pharmacol.* 1983; 5: S 30-34.
34. Hoey A, Jackson C, Pegg G, Sillence M. Atypical responses of rat ileum to pindolol, cyanopindolol, and iodocyanopindolol. *Br J Pharmacol.* 1996; 117: 712-716.
35. Joseph SS, Lynham JA, Molenaar P, Grace AA, Colledge WH, Kaumann AJ. Intrinsic sympathomimetic activity of (-) - pindolol mediated through a (-) - propranolol-resistant site of the beta-1 adrenoceptor in human atrium and recombinant receptors. *Naunyn Schmiedergs Arch Pharmacol.* 2003; 368: 496-503.
36. Emorine LJ, Feve B, Pairault J et al. The human beta-3 adrenergic receptor: relationship with atypical receptors. *Am J Clin Nutr.* 1992; 55(1): 215S-218S.
37. Blin N, Nahmias C, Drumare MF, Strosberg AD. Mediation of most atypical effects by species homologues of the beta-3 adrenoceptor. *Br J Pharmacol.* 1994; 112: 911-919.
38. Baker JG. Evidence for a secondary state of the human beta-3 adrenoceptor. *Mol Pharmacol.* 2005; 68: 1645-1655.
39. Horinouchi T, Asai S, Fukushima M, Koike K. Partial agonist activity of labetalol, the aryethanolamine, on beta-3 adrenoceptors in the guinea gastric fundus. *Auton Autacoid Pharmacol.* 2002; 22: 29-35.
40. Figueroa XF, Poblete I, Fernandez R, Pedemonte C, Cortes V, Huidobro-Toro JP. NO production and e NOS phosphorylation induced by epinephrine through the activation of beta-adrenoceptors. *Am J Physiol Heart Circ Physiol.* 2009; 297: H134-143.
41. Kakoki M, Hirata Y, Hayakawa H, et al. Effects of vasodilatory beta-adrenoceptor antagonists on endothelium-derived nitric oxide release in rat kidney. *Hypertension.* 1999; 33: 467-471.
42. Szilvassy Z, Nagi I, Szilvassy J, Jakab I, Csati S, Lonovics J. Impaired relaxation of the sphincter of Oddi of hyperlipidaemic rabbits. *Eur J Pharmacol.* 1996; 301: R17-18.
43. Heusser K, Schobel HP, Adamidis A, Fischer T, Frank H. Cardiovascular effects of beta-blockers with and without intrinsic sympathomimetic activity. A comparison between celiprolol and metoprolol. *Kidney Blood Press Res.* 2002; 25: 34-41.
44. Kobayashi N, Mori Y, Nakano S, Tsubokou Y, Shirataki H, Matsuoka H. Celiprolol stimulates endothelial nitric oxide synthase expression and improves remodelling in deoxycorticosterone acetate-salt hypertensive rats. *J Hypertens.* 2001; 19: 795-801.
45. Liao Y, Asakura M, Takashima S, et al. Celiprolol, a vasodilatory beta-blocker, inhibits pressure overload-induced cardiac hypertrophy and prevents the transition to heart failure via ni-

- tric oxide-dependent mechanisms in mice. *Circulation*. 2004; 110: 692-699.
46. Chang HY. Role of nitric oxide in vasodilator response induced by salbutamol in rat diaphragmatic microcirculation. *Am J Physiol*. 1997; 272: H173-179.
  47. Figueira ME, de Vale FM, Barroso FM, Rico JM, Castro M. Dietary beta-adrenoceptor agonists have a persistent effect on nitric oxide synthase in rat cultured smooth muscle cells. *Eur J Pharmacol*. 1998; 362: 261-266.
  48. Puri R, Liew GY, Nicholls SJ et al. Coronary beta-2 adrenoceptors mediate endothelium dependent vasoreactivity in humans: novel insights from an in vivo intravascular ultrasound study. *Eur Heart J*. 2012; 33: 495-504.
  49. Huang B, Brennan KM, Budinger TF, Maltz JG. Assessment of endothelial function in the radial artery using inhaled albuterol. *Conf Proc IEEE Eng Med Biol Soc*. 2007; 3629-3631.
  50. Montalvo F, Cantres-Fonseca O, Santos G, et al. Nitric oxide is involved in the response of the isolated intact and epithelium-denuded rat trachea to the beta-2 adrenergic receptor agonist salbutamol. *Respiration*. 2010; 80: 426-432.
  51. Khan MU, Zhao W, Zhao T et al. Nebivolol, a multifaceted anti-oxidant and cardioprotectant in hypertensive heart disease. *J Cardiovasc Pharmacol*. 2013; 62: 445-451.
  52. Kobayashi N, Mita S-I, Yoshida K, et al. Celiprolol activates e NOS through the PI3K-Akt pathway and inhibits VCAM-1 NF-kB induced by oxidative stress. *Hypertension*. 2003; 42: 1004-1013.
  53. Feuerstein GZ, Yue TL, Cheng HY, Ruffolo RR Jr. Myocardial protection by the novel vasodilating beta-blocker carvedilol: potential relevance of anti-oxidant activity. *J Hypertens Suppl*. 1993; 11(4): S41-S48.
  54. Hayashi T, De Velasco MA, Saitou Y, et al. Carvedilol protects tubular epithelial cells from ischemia-reperfusion injury by inhibiting oxidative stress. *Int J Urol*. 2010; 17: 989-995.
  55. Zapida RJ, Castillo R, Rodrigo R, et al. Effect of carvedilol and nebivolol on oxidative stress-related parameters and endothelium function in patients with essential hypertension. *Basic Clin Pharmacol Toxicol*. 2012; 111: 309-316.
  56. Ozaydin M, Icli A, Yucel H, et al. Metoprolol vs carvedilol or carvedilol plus N-acetyl cysteine on post-operative atrial fibrillation: a randomised, double-blind, placebo-controlled study. *Europ Heart J*. 2013; 34: 597-604.
  57. Goldhammer E, Maor I, Schnitzer S, Lanir A, Abinader EG. The early antioxidant effect of carvedilol predicts the clinical course in congestive heart failure patients. *J Cardiovasc Med*. 2007; 8: 453-456.
  58. Li H, Horke S, Forstermann U. Oxidative stress in vascular disease and its pharmacological prevention. *Trends Pharmacol Sci*. 2013; 34: 313-319.
  59. Paneni F, Beckman JA, Creager MA, Cossentino F. Diabetes and vascular disease: pathophysiology, clinical consequences and medical therapy part 1. *Europ Heart J*. 2013; 34: 2436-2443.
  60. Borek C. Dietary antioxidants and human cancer. *Intergr Cancer Ther*. 2004; 3: 333-341.
  61. Juni RP, Duckers HJ, Vanhoutte PM, Virmani R, Moens AL. Oxidative stress and pathological changes after coronary artery interventions. *J Amer Coll Cardiol*. 2013; 61: 1471-1481.
  62. Halliwell B. The antioxidant paradox. *Lancet*. 2000; 355: 1179-1180.
  63. Forstermann V, Munzel T. Endothelial nitric oxide synthase in vascular disease. From marvel to menace. *Circulation*. 2006; 113: 1708-1714.
  64. Bloodsworth A, O'Donnell VB, Freeman BA. Nitric oxide regulation of free radical- and enzyme-mediated lipid and lipoprotein oxidation. *Arterioscler Thromb Vasc Biol*. 2000; 20: 1707-1715.
  65. Mechanisms of the pro- and anti-oxidant actions of nitric oxide in atherosclerosis. *Cardiovasc Res*. 2000; 47: 465-474.
  66. Harman D. Anti-oxidant supplements: effects on disease and aging in the United States. *J Am Aging Assoc*. 2000; 23: 25-31.
  67. Schulman SP, Becker LC, Cass DA. L-Arginine therapy in acute myocardial infarction. *JAMA*. 2006; 295: 58-64.
  68. Bjelakovic G, Nikolova D, Gluud LL, Simonetti RG, Gluud C. Mortality in randomised trials of anti-oxidant supplements for primary and secondary prevention. Systematic Review and Meta-analysis. *JAMA*. 2007; 297: 842-857.
  69. Sesso HD, Buring JE, Christen WG, et al. Vitamins E and C in the prevention of cardiovascular disease in men. The Physicians Health Study 11 Randomised Controlled Trial. *JAMA*. 2008; 300: 2123-2133.
  70. Myung S-K, Ju W, Cho B et al. Efficacy of vitamin and antioxidant supplements in prevention of cardiovascular disease: systematic review and meta-analysis of randomised controlled trials. *BMJ*. 2013; 346: 12.
  71. Bjelakovic G, Nikolova D, Gluud C. Anti-oxidant supplements to prevent mortality. *JAMA*. 2013; 310: 1178-1179.
  72. Dolara P, Bigagli E, Collens A. Antioxidant vitamins and mineral supplementation, life span expansion and cancer incidence: a critical commentary. *Eur J Nutr*. 2012; 51: 769-781.
  73. Jessup M, Brazena S. Heart failure. *N Engl J Med*. 2003; 348: 2007-2008.
  74. Gheorghiade M, Sopko G, De Luca L, et al. Navigating the crossroads of coronary artery disease and heart failure. *Circulation*. 2006; 114: 1202-1213.
  75. Aurigemma GP, Gaasch WH. Diastolic heart failure. *N Engl J Med*. 2004; 351: 1097-1105.
  76. Cruickshank JM. Are we misunderstanding beta-blockers? *Int J Cardiol*. 2007; 120: 10-27.
  77. CIBIS-11 investigators and committee. The Cardiac Insufficiency Bisoprolol Study 11 (CIBIS 11): a randomised trial. *Lancet*. 1999; 353: 9-13.
  78. The MERIT-HR Investigators. Effect of metoprolol CR/XL in chronic heart failure. Metoprolol CR/XL Randomised Intervention Trial in Congestive Heart Failure (MERIT-HR). *Lancet*. 1999; 35: 2001-2007.
  79. Packer M, Coats AJ, Fowler MB et al. Effect of carvedilol on survival in severe chronic heart failure. *N Engl J Med*. 2001; 344: 1651-1658.
  80. Xamoterol in Severe Heart Failure Study Group. Xamoterol in severe heart failure. *Lancet*. 1990; 336: 1-6.
  81. The Beta-blocker Evaluation of Survival Trial Investigators. A trial of the beta-blocker bucindolol in patients with advanced chronic heart failure. *N Engl J Med*. 2001; 344: 1659-1667.
  82. Flather MD, Shibata MC, Coats AJ et al. Randomised trial to determine the effect of nebivolol on mortality and cardiovascular hospital admissions in elderly patients with heart failure (SENIORS). *Eur Heart J*. 2005; 26: 215-225.
  83. Mc Murray J. Making sense of SENIORS. *Eur Heart J*. 2005; 26: 203-206.
  84. Wikstrand J, Wedel H, Castagno D, Mc Murray JJ. The large-scale placebo-controlled beta-blocker studies in systolic heart failure re-visited: results from CIBIS II, COPENICUS, and SENIORS-SHF compared with stratified subsets from MERIT-HF. *J Intern Med*. 2013; Oct 1. doi:10.1111/joim.12141.
  85. Dobre D, van Veldhuisen DJ, Mordenti G, et al. Tolerability and dose-related effects of nebivolol in elderly patients with heart failure. *Am Heart J*. 2007; 154: 109-115.
  86. Ambrosio G, Flather MD, Bohm M, et al. Beta-blockade with nebivolol for prevention of acute ischaemic events in elderly patients with heart failure. *Heart*. 2011; 97: 209-214.
  87. Cook S, Togni M, Schaub MC et al. High heart rate: a cardiovascular risk factor. *Eur Heart J*. 2006; 2: 2387-2393.
  88. Kato N, Kinugawa K, Imamura T, et al. Trend of clinical outcome and surrogate markers during titration of beta-blocker in heart failure patients with reduced ejection fraction: relevance of achieved heart rate and beta-blocker dose. *Circ J*. 2013; 77: 1001-1008.

89. Hardman SM. Ivabradine in heart failure: NICE guidance. *Heart*. 2014; 100: 93-95.
90. Jong P, McKelvie R, Yusuf S. Should treatment of heart failure with preserved ejection fraction differ from that for heart failure with reduced ejection fraction? *BMJ*. 2010; 341: 453-455.
91. Borlaug BA, Paulus WJ. Heart failure with preserved ejection fraction: pathophysiology, diagnosis, and treatment. *Eur Heart J*. 2011; 32: 670-679.
92. Lipsic E, Van Veldhuisen DJ. Nebivolol in chronic heart failure: current evidence and future perspectives. *Expert Opin Pharmacother*. 2010; 11: 983-992.
93. Yamamoto K, Origasa H, Hori M: J-DHF Investigators. Effects of carvedilol on heart failure with preserved ejection fraction: the Japanese Diastolic Heart Failure Study (J-DHF). *Eur J Heart Fail*. 2013; 15: 110-118.
94. Cohen SA, Leurs I, Assayag P et al. Optimization of heart Failure medical Treatment after hospital discharge according to ventricular ejection fraction: the FUTURE survey. *Arch Cardiovasc Dis*. 2012; 105: 355-356.
95. Nevzorov R, Porath A, Henken Y, Kobal SL, Jotkowitz A, Novack V. Effect of beta-blocker therapy on survival of patients with heart failure and preserved systolic function following hospitalisation with acute decompensated heart failure. *Eur J Intern Med*. 2012; 23: 374-378.
96. Richardson DW, Freund J, Gear AS et al. Effect of propranolol on elevated arterial blood pressure. *Circulation*. 1968; 37: 534-42.
97. Buhler FR. Age and cardiovascular response adaptation. *Hypertension*. 1983; 5: 94-100.
98. Prisant LM, Mensah GA. Use of beta-adrenergic blockers in Blacks. *J Clin Pharmacol*. 1996; 36: 867-873.
99. Giles TD. The role of vasodilating beta-blockers in patients with complicated hypertension: focus on nebivolol. *Am J Med*. 2010; 123: S16-S20.
100. Germino FW, Lin Y, Pejovic V, Bowen L. Efficacy and tolerability of nebivolol: does age matter? A retrospective analysis of three randomised, placebo-controlled trials in stage 1-11 hypertension. *Ther Adv Cardiovasc Dis*. 2012; 6: 185-199.
101. Giles TD, Khan BV, Lato J, Brenner L, Ma Y, Lukie T. Nebivolol monotherapy in younger adults (younger than 55 years) with hypertension: a randomised, placebo-controlled trial. *J Clin Hypertens*. 2013; 15: 687-693.
102. Lewin A, Punzitt, Luo X, Stapff A. Nebivolol monotherapy for patients with systolic stage II hypertension: results of a randomised, placebo-controlled trial. *Clin Ther*. 2013; 35: 142-152.
103. Badar VA, Hiware SK, Shrivastava MP, Thawani VR, Hardas MM. Comparison of nebivolol and atenolol on blood pressure, blood sugar and lipid profile in patients of essential hypertension. *Indian J Pharmacol*. 2011; 43: 437-440.
104. Fici F, Celik T, Balta S, et al. Comparative effects of nebivolol and metoprolol on red cell distribution width and neutrophil/lymphocyte ratio in patients with new diagnosed hypertension. *J Cardiovasc Pharmacol*. 2013; 62: 388-393.
105. Czuriga I, Rieckensky I, Bodnar J et al. Comparison of the new cardioselective beta-blocker nebivolol with bisoprolol in hypertension: Nebivolol, Bisoprolol Multicentre Study (NEBIS). *Cardiovasc Drugs Ther*. 2003; 17: 257-263.
106. Erdogan O, Ertem B, Altun A. Comparison of antihypertensive efficacy of carvedilol and nebivolol in mild-to moderate primary hypertension: a randomised trial. *Anadolu Kardiyol Derg*. 2011; 11: 310-313.
107. Mazza A, Gil-Extremera B, Maldonado A, Toutouzas T, Pessina A. Nebivolol vs Amlodipine as first-line treatment of essential arterial hypertension in the elderly. *Blood Pressure*. 2002; 11: 182-188.
108. Poirier L, Cleroux J, Nodeau A, Lacourciere Y. Effects of nebivolol and atenolol on insulin sensitivity and haemodynamics in hypertensive patients. *J Hypertens*. 2001; 19: 1429-1435.
109. Ayers K, Byrne I, Dematteo A, Brown N. Differential effects of nebivolol and metoprolol on insulin sensitivity and plasminogen activator inhibitor in the metabolic syndrome. *Hypertension*. 2012; 59: 893-898.
110. Basile JN. One size does not fit all: the role of vasodilating beta-blockers in controlling hypertension as a means of reducing cardiovascular and stroke risk. *Am J Med*. 2010; 123: S9-S15.
111. Broekman CP, Haensel SM, van den Ven LL, Slob AK. Bisoprolol and hypertension: effects on sexual dysfunction in men. *J Sex Marital Ther*. 1992; 18: 325-331.
112. Bodmer M, Egger S, Hohenstein E, Beltraminelli H, Krahenbuhl S. Lichenoid eruption associated with the use of nebivolol. *Ann Pharmacother*. 2006; 40: 1688-1690.
113. Lane DA, Lip GY. Treatment of hypertension in peripheral arterial disease. *Cochrane Database Syst Rev* 2013; Dec 4th.
114. Diehm C, Pittrow D, Lowall H, et al. Effect of nebivolol vs hydrochlorothiazide on the walking capacity in hypertensive patients with intermittent claudication. *J Hypertens*. 2011; 29: 1448-1456.
115. Wilkinson IB, Franklin SS, Hall I, et al. Pressure amplification explains why pulse pressure is unrelated to risk in young subjects. *Hypertension*. 2001; 38: 1461-1466.
116. Williams B, Lacy PS. Central aortic pressure and clinical outcomes. *J Hypertens*. 2009; 27: 1123-1125.
117. Benetos A, Gautier S, Labat C, et al. Mortality and cardiovascular events are best predicted by low central/peripheral pulse pressure amplification, but not by high blood pressure levels in elderly nursing homes subjects. The PARTAGE Study. *J Am Coll Cardiol*. 2012; 60: 1503-1511.
118. Pedersen ME, Cockcroft JR. The vasodilatory beta-blockers. *Curr Hypertens Rep*. 2007; 9: 269-277.
119. Kelly R, Daley J, Avolio A, O'Rourke M. Arterial dilation and reduced wave reflection. Benefit of diltiazem in hypertension. *Hypertension*. 1989; 14: 14-21.
120. Morgan T, Lauri J, Bertrum D, Anderson A. Effect of different anti-hypertensive drug classes on central aortic pressure. *Am J Hypertens*. 2004; 17: 118-123.
121. Soanker R, Naidu MU, Raju SB, Prasad AK, Rao TR. Effect of beta-1 blocker nebivolol on central aortic pressure and arterial stiffness in patients with essential hypertension. *Indian J Pharmacol*. 2012; 44: 407-411.
122. Davis JT, Pasha DN, Khandrika S, Fung MM, Milic M, O'Connor DT. Central haemodynamics of prehypertension: effect of the beta-adrenergic antagonist nebivolol. *J Clin Hypertens*. 2013; 15: 69-74.
123. Dhakam Z, Yasmin, McEniery CM, Burton T, Brown MJ, Wilkinson IB. A comparison of atenolol and nebivolol in isolated systolic hypertension. *J Hypertens*. 2008; 26: 351-356.
124. Redon J, Pascual-Izuel JM, Rodilla E, et al. Effects of nebivolol and atenolol on central aortic pressure in hypertensive patients: a multicentre, randomised, double-blind study. *Blood Press*. 2014; 23(3): 181-188.
125. Briasoulis A, Oliva R, Kalaitzidis R et al. Effect of nebivolol on aortic compliance in patients with diabetes and maximal renin-angiotensin system blockade: the EFFORT study. *J Clin Hypertens*. 2013; 15: 473-479.
126. Studinger P, Tabak AG, Chen CH, et al. The effects of low dose carvedilol, nebivolol, and metoprolol in central arterial pressure and its determinants: a randomised clinical trial. *J Clin Hypertens*. 2013; 15: 910-917.
127. Kampus P, Serg M, Kals J et al. Differential effects of nebivolol and metoprolol on central aortic pressure and left ventricular mass. *Hypertension*. 2011; 57: 1122-1128.
128. Schattenkerk DW, van den Bogaard B, Cammenga M, Westerhof BE, Stroe ES, van den Born B-J. Lack of difference between nebivolol/hydrochlorothiazide and metoprolol/hydrochlorothiazide on aortic wave augmentation and central blood pressure. *J Hypertens*. 2013; 31: 2447-2454.

## Nebivolol

129. Podzolkov VI, Mazharova LG. Use of selective long-acting beta-blocker with vasodilating activity celiprolol in patients with essential hypertension. *Kardiologia*. 2002; 42: 22-26.
130. Shah NK, Smith SM, Nichols WW et al. Carvedilol reduces aortic wave reflection and improves left ventricular/vascular coupling: a comparison with atenolol (CENTRAL Study). *J Clin Hypertens*. 2011; 13: 917-924.
131. Tahvanainen AM, Tikkanen AJ, Leskinen MH et al. Supine and upright haemodynamic effects of sublingual nitroglycerin and inhaled salbutamol: a double-blind, placebo-controlled, randomised study. *J Hypertens*. 2012; 30: 297-306.
132. Zhou W-J, Wang R-Y, Li Y, et al. A randomized controlled study on the effects of bisoprolol and atenolol on sympathetic nervous activity and central aortic pressure in patients with essential hypertension. *Plos one*. 2013; 8: 1-7.
133. Gillman MW, Kannel WB, Balanger A, et al. Influence of heart rate on mortality among persons with hypertension: the Framingham Study. *Am Heart J*. 1993; 125: 1148-1154.
134. Sahana GN, Sarala N, Kumar TN, Lakshmaiah V. A comparative study of nebivolol and (S) atenolol on blood pressure and heart rate on essential hypertensive patients. *Indian J Pharmacol*. 2010; 42: 401-405.
135. Kjekshus J. Comments – beta-blockers: heart rate reduction a mechanism of benefit. *Eur Heart J*. 1985; 6(suppl): 9-30.
136. Kjekshus JK. Importance of heart rate in determining beta-blocker efficacy in acute and long-term acute myocardial infarction intervention trials. *Am J Cardiol*. 1986; 57: 43F-49F.

**Copyright:** 2014 © John Cruickshank. This is an Open Access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.