

Cardio-Selective Beta-Blocker: Pharmacological Evidence and Their Influence on Exercise Capacity

Dennis Ladage,^{1,3} Robert H.G. Schwinger² & Klara Brixius¹

¹ Department of Molecular and Cellular Sport Medicine, Institute of Cardiology and Sport Medicine, German Sport University Cologne, Cologne, Germany

² Clinic II of Internal Medicine and Cardiology, Weiden, Germany

³ Heart Center, Department III of Internal Medicine, University of Cologne, Cologne, Germany

Keywords

Beta-blocker; Hypertension; Cardiovascular disease.

Correspondence

Prof. Dr. Klara Brixius, Department of Molecular and Cellular Sport Medicine, Institute of Cardiology and Sport Medicine, German Sport University Cologne, Carl-Diem-Weg 6, 50933 Cologne, Germany.

Tel.: +49-221-49825220;

Fax: +49-221-49828370;

E-mail: Brixius@dshs-koeln.de

SUMMARY

For the past 40 years, beta-blockers have been widely used in cardiovascular medicine, reducing morbidity as well as mortality. Beta-blockers are currently used in a number of cardiovascular conditions such as systolic heart failure, postmyocardial infarction, and in prevention and treatment of arrhythmias. They are not recommended as the first line antihypertensive therapy, particularly in the elderly, unless there are specific indications. Despite the benefits of beta-blockers, tolerability concerns in patients with co-morbidities have limited their use. Some of these problems were overcome with the discovery of cardioselective beta-blockers. The third generation beta-blockers have additional properties of vasodilatation and advantages in terms of minimizing the adverse effects of beta-blockers. Some of the advantages include improvement of insulin resistance, decrease in cholesterol as well as alleviation of erectile dysfunction. Acute treatment with beta-blockers modifies local muscular metabolic properties and impairs endurance exercise capacity whereas the influence of chronic is debated controversially.

doi: 10.1111/j.1755-5922.2011.00306.x

Introduction

β -adrenoceptor blocking agents (beta-blockers) were first discovered in 1962 by Sir James Black at the Imperial Chemical Industries in the United Kingdom [1]. The development of propranolol is considered one of the most important contributions to clinical medicine and pharmacology and Sir Black was consecutively awarded the Nobel prize in 1988. In combination with ACE inhibitors, antiplatelets and statins beta-blockers were proven to reduce cardiovascular mortality and morbidity by as much as 90% in acute coronary syndromes [2].

Beta-blockers have evolved ever since from their origin as treatment for angina and cardiac arrhythmia to be very successful therapeutics in large variety of diseases as hypertension, glaucoma, migraine just to name a few. However, the mechanism of action of beta-blockers probably differ within the class and are not fully understood yet. Newer third generation beta-blockers exhibit additional properties including the improvement of insulin resistance, decrease in cholesterol as well as alleviation of erectile dysfunction.

Furthermore, cardiovascular conditions are common in the general population and thus are also common in athletes or patients who are physically active. When choosing a drug for these patients, providers should choose an agent that has favorable effects on blood pressure and minimal detrimental hemodynamic change during exercise. This review discusses the pharmacological properties of cardioselective beta-blockers, their clinical use in cardiovascular medicine as well as their influence on exercise capacity.

Mechanism of Action of Beta-Blockers

β -adrenergic receptors are essential regulators of cardiovascular homeostasis. Three different types of beta-receptors have been identified by molecular pharmacology: β_1 , β_2 , and β_3 are variably distributed in different tissues [3, 4]. β_1 receptors are predominantly located in the heart and make up to 75% of all b-ARs, while β_2 receptors are found in vascular and bronchial smooth muscle [5]. β_3 receptors are located in the adipocytes where they are presumed to be involved in the fatty acid metabolism [6, 7]. In addition, β_3 -adrenoceptors have been described in




Figure 1 Comparison of the affinity ratio to the β_1 and β_2 -adrenoceptor for currently available beta blockers and intracellular signal transduction. Under agonist action β_1 and β_2 receptors promote the formation of cAMP by interacting with adenylate cyclase via activation of the Gs protein. Moreover β_2 receptors have also been found to couple with Gi and decrease adenylcy-

clase activity. cAMP facilitates the PKA phosphorylation which then activates the calcium release channel (ryanodine receptor) as well as the SERCA activity. Increased cytosolic calcium results in an increased contraction of the heart muscle cell.

cardiomyocytes, where they make up a population of about 5% of the β -AR [8], as well as in endothelial cells [9]. Furthermore, a putative β_4 receptor has been reported to exist in the heart [10], however there is now evidence that this putative receptor is another activation status of the β_1 -AR [11].

All beta-blockers cause a competitive inhibition of the β receptor and counter the effects of catecholamines. The β_2 -adrenoceptor, like the β_1 -adrenoceptor, is a G-protein coupled receptor that couples to G α s-proteins. If activated it stimulates an increase in intracellular cAMP via the adenylyl cyclase (Figure 1) [12]. Moreover the cyclic AMP Q1 (cAMP) response element binding protein (CREB) is activated by beta-adrenoceptor signaling [13]. cAMP, the second messenger, activates protein kinase A (PKA) which phosphorylates the membrane calcium channel and hence increases calcium entry into the cytosol. In addition, PKA

enhances calcium release from the sarcoplasmic reticulum. The increase in calcium loading leads to an positive inotropic effect. The lusitropic effect is caused by phosphorylation of troponin I and phospholamban by PKA, which increase the calcium reuptake by sarcoplasmic reticulum [14].

Chronically increased noradrenaline levels induce downregulation of β_1 -adrenoceptors and alter signal transduction [15]. In failing human hearts noradrenaline concentration is increased in left ventricular biopsies [16], which leads to a reduction of myocardial β_1 -AR as a hallmark of CHF. β_1 -AR downregulation is associated with a reduction in the re-uptake of [11C]-hydroxyephedrine by neural terminals in the myocardium [17]. Thus the less efficient disposal of catecholamines from the synaptic cleft contributes to myocardial β -AR downregulation. In parallel with the alterations of the β_1 (as well as the β_2)-AR an upregulation of Gi-proteins has

been described [18]. All these reactions result in a massive depression of cardiac contractility. β_3 -AR expression has been shown to be upregulated and this further contributes to the depression of cardiac force by the release of nitric oxide, which has been shown to impair cardiac force of contraction [19].

Clinical Pharmacology

Beta-Blockers are in general well absorbed after oral administration [20]. The plasma half-life of beta-blockers is significantly shorter than their biologic half-life with esmolol having the shortest half-life of about 10 min [21]. The water-soluble beta-blockers (e.g., Atenolol) have in general longer half-lives, tend to be non-metabolized, and excreted unchanged via the kidney [22]. In contrast, Lipid-soluble beta-blockers (e.g., metoprolol) have shorter half-lives and are metabolized mainly in the liver [23].

Currently, beta-blockers can be categorized based on their receptor affinity and hemodynamic properties: the absence or presence of cardioselectivity and vasodilatory effects. The mechanisms of vasodilation are vasodilation mediated either via nitric oxide (carvedilol, nebivolol) or via α receptor blockade (labetalol, carvedilol) [24]. Carvedilol and nebivolol are in addition antiproliferative, antioxidant and block the expression of several genes involved in myocardial damage [25]. Nebivolol reduces interleukin-1 α , cyclooxygenase-2 (COX-2), and tumor-necrosis-factor (TNF)- α -induced protein, while the beta-blocker metoprolol increases inflammatory genes [26]. Tumor necrosis factor- α (TNF- α) is a pluripotent mediator of inflammation involved in atherogenesis inducing a pro-inflammatory activation of macrophages [27]. COX-2 is an inducible enzyme that is expressed in response to inflammatory cytokines and is responsible for the synthesis of proinflammatory PGs such as PGE₂, which is chemoattractant for leukocytes [28].

Another criterion that distinguishes a smaller group is the presence of intrinsic sympathomimetic activity (ISA). These beta-blockers have lately been suspected to increase peripheral vascular resistance and attenuate the decreases in heart rate and cardiac output that may negate any benefit in a cardiovascular disease population [29]. Examples of beta-blockers with this property are acebutol and pindolol.

Noncardioselective beta-blockers such as propranolol cause an equal blockade of β_1 - and β_2 -adrenergic receptors while vasodilating effects are absent [30]. Beta-blocker such as atenolol and metoprolol have a higher affinity for the β_1 receptor and are therefore deemed cardioselective. Nevertheless, with increasing doses some β_2 effects may appear [30]. These beta-blockers have no vasodilatory properties. Noncardioselective beta-blocker as carvedilol exerts equal blockade of the β_1 - and β_2 -adrenergic receptors, while exhibiting vasodilating properties due to α_1 -receptor blockade [31]. One of the newest agents is the cardioselective and vasodilating beta-blocker Nebivolol. Nebivolol has demonstrated to have the highest cardioselectivity of any currently available beta-blocker [32]. In addition, nebivolol exhibits vasodilatory properties not related to α_1 -receptor blockade but instead the vasodilatation mediated by nitric oxide (NO) [33].

A recent pharmacological study has shown that several of beta-adrenoceptor antagonists of the first and second generation cur-

Table 1 Use of beta-blockers in AHA/ESC guidelines

Strong evidence	
Secondary prevention post-MI	- improve survival (I A) - prevention of sudden death (I A) - prevention of re-infarction (I A)
Chronic heart failure (CHF)	- prolong survival (I A)
Some evidence	
Acute myocardial infarction (AMI)	- relief of ischemic pain (I B) - control arrhythmia (I B) - prevention of sudden death (I C) - limit infarct size (IIa A)
Weak evidence	
Hypertension (HTN)	- recommendation as first-line therapy in uncomplicated HTN withdrawn - (currently guidelines are debated)

rently in clinical use for hypertension, heart disease and, more recently, heart failure have poor selectivity between adrenoceptor subtypes 1 and 2 in whole-cell assays [34].

Clinical Uses and Comparisons

The benefit and clinical indications of beta-blockers have been clearly defined in many cardiovascular conditions and clinical settings (Table 1). In patients at risk for coronary vascular disease a meta-analysis of 18 randomized long-term, placebo-controlled trials found treatment with beta-blockers associated with a reduction in the risk of stroke by 29%, coronary heart disease by 7%, and heart failure by 29% [35]. In a large meta-analysis in post-MI patients beta-blockers proved to reduce significantly morbidity and mortality, regardless whether selective or nonselective beta-blockers were used [36].

Beta-blockers are among the standard therapy for heart failure patients [37], although only bisoprolol, metoprolol, and carvedilol have proven themselves in large randomized trials [38–40]. Some studies suggest that beta-1 selective blockers have a slightly greater antihypertensive effect than the nonselective blockers. Evidence that beta-2 blockade does not contribute to the antihypertensive effect of beta-blockade originates from a study with the beta-2 selective blocker ICI 188 551 [41]. Support is drawn from early observations that adrenaline in presence of beta-2 blockade produces a greater rise in blood pressure compared to its beta-2 vasodilator action [42]. Beta-2 blockade may antagonize the modest background vasodilator effect of circulating adrenaline that explains the 2–3 mmHg greater fall in blood pressure seen with beta-1 blockade as opposed to nonselective block [43]. Beta-blockers with ISA seem to have a reduced clinical benefits in post-MI patients, so that these agents should be avoided in this population [36]. Furthermore, these agents have also been observed to have a negative effect on heart failure in high-risk patients [44].

The anti-arrhythmic effects of beta-blockers are mainly related to their capacity to inhibit beta-receptor-mediated adrenergic neural activation [45]. The ability of the beta-blockers to slow conduction and increase atrioventricular node refractoriness is useful

Table 2. Comparison of beta-blockade affecting exercise capacity

Study	Drug	Condition	Results
Dubach 2002	Bisoprolol	Heart failure	Exercise capacity ↑
Issa 2007	Bisoprolol	Heart failure	Exercise capacity unchanged
Nodari 2003	Atenolol	Heart failure	Exercise capacity unchanged
	Nebivolol	Heart failure	Exercise capacity ↑
Patrianakos 2005	Nebivolol	DCM	Exercise capacity ↑
Diehm 2011	Nebivolol	Heart failure	Exercise capacity unchanged
Dalla Libera 2010	Nebivolol	Heart failure	Prevents muscle wastage thus improves exercise capacity
Patrianakos 2005	Nebivolol	Heart failure	Exercise capacity ↑
	Carvedilol	Heart failure	Exercise capacity ↑
Suazzi 1999	Carvedilol	Heart failure	No effect, pulmonary dysfunction improved
Agostini 2002	Carvedilol	Heart failure	Exercise capacity ↑
Volterrani 2011	Carvedilol	Heart failure	Exercise capacity unchanged
Nessler 2008	Carvedilol	Heart failure	Exercise capacity ↑
Witte 2005	all	Heart failure	Exercise capacity ↑
Beloka 2008	Bisoprolol	Healthy	Exercise capacity ↓
Vankees 2000	Bisoprolol	Healthy	Exercise capacity ↓
Van Bortel 1992	Nebivolol	Healthy	Exercise capacity unchanged

in the prevention and treatment of supraventricular arrhythmias [46]. These drugs are also effective in arrhythmias related to excessive cardiac adrenergic stimulation such as those associated with thyrotoxicosis, pheochromocytoma, exercise, and emotional stress [47–49]. Beta-blockers can prevent exercise-induced augmented ventricular ectopy in patients with coronary artery disease and treat arrhythmias associated with the mitral valve prolapse syndrome [50]. Furthermore, they are effective in reducing the number and complexity of ventricular contractions after myocardial infarction. Several studies have demonstrated a reduction in the incidence of sudden death after myocardial infarction [51].

Effects on Glucose and Lipid Metabolism

Diabetes and adipositas are growing concerns in the western societies but also in developing countries. In patients with diabetes and/or adipositas differences among beta-blockers may have significant clinical implication. Some studies suggest that beta-blockers even raise the risk of occurrence of diabetes as well as of dyslipidemia [52]. Metabolic changes involve blood sugar, HbA1c, insulin-sensitivity, triglycerides, VLDL, and HDL, and are closely associated with beta-2 blockade [53]. Various mechanisms are discussed to explain these negative metabolic effects of beta-blockers. For once beta-blockers produce unopposed α 1-adrenergic receptor activity, which may induce vasoconstriction, decrease skeletal blood flow, and reduce insulin-stimulated peripheral glucose uptake [54, 55]. Furthermore, beta-blockers may inhibit insulin secretion from pancreatic beta-cells [56]. Weight gain is another proposed mechanism of insulin sensitivity deterioration with beta-blockers, as beta-blockers are associated with increased body weight in numerous large studies [57] while weight gain is closely connected to insulin sensitivity reduction [58].

The nonselective beta-blockers (e.g., timolol and propranolol) are the most susceptible to cause these changes, followed by partially beta-1 selective agents (e.g., metoprolol and atenolol). In

contrast in highly beta-1 selective agents (e.g., nebivolol [59] or bisoprolol [60]) the metabolic disturbances involving blood sugar, insulin-sensitivity, and lipids are almost completely absent [61]. This also applies for agents triggering alpha-blockade (e.g., carvedilol) [62] or beta-2 intrinsic sympathomimetic activity (e.g., nebivolol) [63]. Current guidelines for the management of hypertension emphasize beta-blockers as preferred agents for patients with concomitant diabetes [64].

Lately, there has been interest in the beta-3 receptor and especially in its role in obesity and diabetes. The beta-3 adrenoceptors are found in adipose tissue and in the skeletal muscle [65, 66]. Agonists at the beta-3 receptor lead to increased fat oxidation, lipolysis, and insulin sensitivity. On the other hand, blockade of beta-3 receptors are associated with weight increase and unfavorable metabolic changes [53].

Thus, beta-blockers differ regarding their influence on glucose or fat metabolism with respect to their molecular pharmacological mechanisms. These differences should be taken into account when treating cardiovascular patients with additional metabolic diseases.

Impact on Skeletal Muscle Adaptation and Exercise Capacity

The above-described influence of β -blockers on glucose and fat metabolism may also influence skeletal muscle adaptation in situations of acute and chronic exercise and thus exercise capacity. The effects of beta-blockade on adaptations to aerobic exercise have been studied in healthy and hypertensive subjects under both acute and chronic conditions (Table 2, Figure 2).

Acute treatment with beta-blocking drugs modifies local muscular metabolic properties [67] and impairs endurance exercise capacity resulting in an increase in perceived exertion, lower VO₂max, and lower work rate [68, 69]. Especially the




Figure 2 Treatment with beta-blockers affects physical activity. Beta-blockers reduce heart rate both in rest and under exertion. Acute treatment with beta-blocking drugs modifies local muscular metabolic properties and impairs endurance exercise capacity resulting in an increase in

perceived exertion, lower VO_{2max} and lower work rate. Nonselective beta-blockers can increase lactate levels during exercise through β_2 -associated peripheral vasoconstriction.

nonselective inhibitors, which also antagonize β_2 -receptors, can restrict aerobic training effects [70].

In normal subjects, beta-blocker treatment with metoprolol is associated with a decrease in sub maximal exercise \dot{V}_E and in maximum O_2 -uptake (VO_{2max}) [69]. However, as described above, differences may exist depending on the underlying molecular pharmacological effects of the respective β -blocking agent. Nebivolol, for example, does not impair exercise capacity in healthy persons [71].

Depression of exercise capacity by at least some beta-blocking agents is due to limitation of oxygen convection to the muscles by decreased chronotropy and inotropy [72]. However, by a relative decrease in cardiac output and O_2 -transport the muscle metaboreflex during exercise may be enhanced, thus leading to increased chemosensitivity and activation of the sympathetic nervous system [72, 73].

Recent studies reported a marked increase of MSNA after acute β -blockade associated with unchanged values of arterial blood pressure and baroreflex sensitivity. Acute β -blockade also alters the oscillatory characteristics of MSNA, thus decreasing its effectiveness on peripheral vasoconstriction [74]. This reaction may counteract the above-mentioned suppression of the cardiopulmonary system and may contribute to an upregulation or at least compensation of exercise capacity when treated with β -blocking agents. Data from animal studies confirm that β_2 -blockade may

impair the adaptation to endurance exercise to a greater extent than β_1 -blockade [75]. In accordance metoprolol compared with carvedilol provided a greater effect in LV ejection fraction at rest and in LV stroke volume and stroke work during exercise. In contrast, carvedilol produces greater decreases in mean pulmonary artery pressure and pulmonary wedge pressure, both at rest and during exercise [25].

Aside from the different physiological effects, β -blocker treatment may also alter skeletal muscle and resting conditions as well as skeletal muscle adaptation toward exercise. It has been shown, for example, that one of the suggested mechanisms are mitochondrial adaptations. Mitochondrial function (succinate dehydrogenase and citrate synthase activity) can be increased by endurance exercise, while the response to training can be attenuated when animals or healthy subjects are treated with selective or nonselective beta-blockers. The impaired mitochondrial response may be due to a β_2 -receptor-mediated decrease in PGC-1 [76, 77]. However, there is also evidence that β -blockers like nebivolol (Libera, Vescovio 2010) or carvedilol (Della, Libera 2005) stop skeletal muscle wastage which is a common co-morbidity in heart failure patients.

Exercise-induced dyspnea in patients with heart failure may be related to sympathetic nervous system activation involving an increased metabo- and/or chemosensitivity [78]. Whether or not this mechanism plays a role in exercising normal subjects remains

unclear. Studies by Beloka et al. suggest that decreased aerobic exercise capacity under beta-adrenergic blockade goes along with decreased ventilation regardless of the present metabolic rate [72]. Moreover, beta-blockade was shown to reduce the sensitivity of arterial chemoceptors to potassium stimulation during exercise, thereby modulating the regulation of exercise ventilation [79], although effects could be secondary to reduced cardiac output [80]. Duration of beta-blocker treatment prior to exercise testing seems not to significantly alter exercise responses [81].

Furthermore, beta-blockers affect the pulmonary mechanical function as well as gas diffusion. The respiratory system is characterized by a significant predominance of β_2 receptors, in the airways and the alveoli [82]. Beta-receptors located in the alveoli influence gas exchange efficacy by regulating fluid reabsorption from the alveolar surface. Therefore, different β -receptor affinities result in different effects on the carbon monoxide diffusing capacity (DLCO). Long-term bisoprolol treatment leads to higher DLCO values because of changes in the active membrane transport under alveolar β_2 -receptor control [83]. In addition, carvedilol reduces hyperventilation by reducing peripheral chemoreflex sensitivity as suggested by increased PaCO₂ with normoxia [84].

Overall, beta1-selective beta-blockers are preferable to nonselective agents for patients engaged in exercise training. Furthermore, beta-blockers affect several pathways or systems that ultimately can reduce exercise capacity during training.

Perspective

Beta-blockers are an important class of drugs that are widely recommended for the management of cardiovascular disease, heart failure, and hypertension. However, pharmacological differences among beta-blockers impact use in everyday practice. The clinical rationale for using beta-blockers is to antagonize the beta-1-adrenoceptor signaling to reduce the rate and force of cardiac muscle contraction. However, the beneficial mode of action in heart failure is still not understood. In an aging society, with increasing cardiovascular morbidity and thus rising use of beta-blockers, it is important to obtain information about the degree of physical impairment by the use of these drugs. This is especially true among

young and physically active patients. In addition, people are increasingly taking more physical exercise as a consequence of the growing awareness of the importance and the lifestyle aspects of exercise.

Although highly selective beta-1-antagonists do exist, most of the beta-blockers available for clinical use are also antagonists at the beta 2-adrenoceptors, which are expressed at high levels in the airways, causing bronchospasm as a major side-effect of this therapeutic class. In a study by Baker the binding of a wide range of beta-adrenoceptor antagonists to stably expressed 1- and 2- adrenoceptors in an identical mammalian cell culture environment was measured [85]. It was shown that although great variation in the potency of clinically used beta-blockers for all receptor subtypes exist, these agents only showed little selectivity between 1 and 2 subtypes. Therefore, the clinical benefit of beta-blockers especially in heart failure is unlikely to be attributed to receptor affinity alone. Other factors, such as longevity of action at the receptors, presumably also play an important role.

Moreover, there is another controversy emerging for beta-blockers—the paradoxical hypothesis that beta-blockers might actually be beneficial in asthma is now being tested in clinical trials [86]. These studies were initiated after long-term exposure was shown to reduce lung sensitivity in a mouse model of asthma.

While research is moving forward we need to learn more about the complex pharmacology of adrenoceptors. Beta-blockers and exercise effects of it are still a matter of ongoing research. Lack of research evidence regarding possible drug and exercise interaction necessitates the close monitoring of patients undergoing pharmacotherapy. This includes tracking heart rate, rating of perceived exertion and blood pressure before, during and after exercise sessions. Overall there is clearly great potential for the development of beta-blockers with greater selectivity and less side-effects for a wide range of indications.

Conflict of Interest/Disclosure Statement

The authors have no conflicts of interest that are relevant to the content of this study.

References

- Black JW, Stephenson JS. Pharmacology of a new adrenergic beta-receptor-blocking compound (Nethalide). *Lancet* 1962;**2**:311–314.
- Mukherjee D, Fang J, Chetcuti S, Moscucci M, Kline-Rogers E, Eagle KA. Impact of combination evidence-based medical therapy on mortality in patients with acute coronary syndromes. *Circulation* 2004;**109**:745–749.
- Benovic JL, Bouvier M, Caron MG, Lefkowitz RJ. Regulation of adenylyl cyclase-coupled beta-adrenergic receptors. *Annu Rev Cell Biol* 1988;**4**:405–428.
- Bylund DB, Eikenberg DC, Hieble JP, et al. International Union of Pharmacology nomenclature of adrenoceptors. *Pharmacol Rev* 1994;**46**:121–136.
- Lopez-Sendon J, Swedberg K, McMurray J, et al. Expert consensus document on beta-adrenergic receptor blockers. *Eur Heart J* 2004;**25**:1341–1362.
- Pott C, Brixius K, Bundkirchen A, et al. The preferential beta3-adrenoceptor agonist BRL 37344 increases force via beta1-/beta2-adrenoceptors and induces endothelial nitric oxide synthase via beta3-adrenoceptors in human atrial myocardium. *Br J Pharmacol* 2003;**138**:521–529.
- Pott C, Steinritz D, Napp A, Bloch W, Schwinger RH, Brixius K. On the function of beta3-adrenoceptors in the human heart: Signal transduction, inotropic effect and therapeutic prospects. *Wien Med Wochenschr* 2006;**156**:451–458.
- Moniotte S, Kobzik L, Feron O, Trochu JN, Gauthier C, Balligand JL. Upregulation of beta(3)-adrenoceptors and altered contractile response to inotropic amines in human failing myocardium. *Circulation* 2001;**103**:1649–1655.
- Berlan M, Galitzky J, Bousquet-Melou A, Lafontan M, Montastruc JL. Beta-3 adrenoceptor-mediated increase in cutaneous blood flow in the dog. *J Pharmacol Exp Ther* 1994;**268**:1444–1451.
- Kaumann AJ, Molenaar P. Modulation of human cardiac function through 4 beta-adrenoceptor populations. *Naunyn Schmiedeberg Arch Pharmacol* 1997;**355**:667–681.
- Michel MC, Harding SE, Bond RA. Are there functional beta-adrenoceptors in the human heart? *Br J Pharmacol* 2011;**162**:817–822.
- Daaka Y, Luttrell LM, Ahn S, Della Rocca GJ, Ferguson SS, Caron MG, Lefkowitz RJ. Essential

- role for G protein-coupled receptor endocytosis in the activation of mitogen-activated protein kinase. *J Biol Chem* 1998;**273**:685–688.
13. Baker JG, Hall IP, Hill SJ. Agonist and inverse agonist actions of beta-blockers at the human beta 2-adrenoceptor provide evidence for agonist-directed signaling. *Mol Pharmacol* 2003;**64**:1357–1369.
 14. Lemoine H, Schonell H, Kaumann AJ. Contribution of beta 1- and beta 2-adrenoceptors of human atrium and ventricle to the effects of noradrenaline and adrenaline as assessed with (-)-atenolol. *Br J Pharmacol* 1988;**95**:55–66.
 15. Gaemperli O, Liga R, Spyrou N, et al. Myocardial beta-adrenoceptor down-regulation early after infarction is associated with long-term incidence of congestive heart failure. *Eur Heart J* 2010;**31**:1722–1729.
 16. Bristow MR, Ginsburg R, Umans V, et al. Beta 1- and beta 2-adrenergic-receptor subpopulations in nonfailing and failing human ventricular myocardium: Coupling of both receptor subtypes to muscle contraction and selective beta 1-receptor down-regulation in heart failure. *Circ Res* 1986;**59**:297–309.
 17. Schafers M, Dutka D, Rhodes CG, Lammertsma AA, Hermansen F, Schober O, Camici PG. Myocardial presynaptic and postsynaptic autonomic dysfunction in hypertrophic cardiomyopathy. *Circ Res* 1998;**82**:57–62.
 18. Ungerer M, Bohm M, Elce JS, Erdmann E, Lohse MJ. Altered expression of beta-adrenergic receptor kinase and beta 1-adrenergic receptors in the failing human heart. *Circulation* 1993;**87**:454–463.
 19. Varghese P, Harrison RW, Lofthouse RA, Georgakopoulos D, Berkowitz DE, Hare JM. beta(3)-adrenoceptor deficiency blocks nitric oxide-dependent inhibition of myocardial contractility. *J Clin Invest* 2000;**106**:697–703.
 20. Wood AJ. Pharmacologic differences between beta blockers. *Am Heart J* 1984;**108**:1070–1077.
 21. Covinsky JO. Esmolol: A novel cardioselective, titratable, intravenous beta-blocker with ultrashort half-life. *Drug Intell Clin Pharm* 1987;**21**:316–321.
 22. Kirch W, Gorg KG. Clinical pharmacokinetics of atenolol—a review. *Eur J Drug Metab Pharmacokinet* 1982;**7**:81–91.
 23. McDevitt DG. Comparison of pharmacokinetic properties of beta-adrenoceptor blocking drugs. *Eur Heart J* 1987;**8** Suppl M:9–14.
 24. Kalinowski L, Dobrucki LW, Szczepanska-Konkel M, Jankowski M, Martyniec L, Angielski S, Malinski T. Third-generation beta-blockers stimulate nitric oxide release from endothelial cells through ATP efflux: A novel mechanism for antihypertensive action. *Circulation* 2003;**107**:2747–2752.
 25. Yue TL, Wang X, Gu JL, Ruffolo RR, Jr., Feuerstein GZ. Carvedilol, a new vasodilating beta-adrenoceptor blocker, inhibits oxidation of low-density lipoproteins by vascular smooth muscle cells and prevents leukocyte adhesion to smooth muscle cells. *J Pharmacol Exp Ther* 1995;**273**:1442–1449.
 26. Wolf SC, Sauter G, Jobst J, Kempf VA, Risler T, Brehm BR. Major differences in gene expression in human coronary smooth muscle cells after nebivolol or metoprolol treatment. *Int J Cardiol* 2008;**125**:4–10.
 27. Beutler B, Cerami A. The biology of cachectin/TNF—a primary mediator of the host response. *Annu Rev Immunol* 1989;**7**:625–655.
 28. Hernandez-Presa MA, Martin-Ventura JL, Ortego M, et al. Atorvastatin reduces the expression of cyclooxygenase-2 in a rabbit model of atherosclerosis and in cultured vascular smooth muscle cells. *Atherosclerosis* 2002;**160**:49–58.
 29. Prichard BN. Pharmacologic aspects of intrinsic sympathomimetic activity in beta-blocking drugs. *Am J Cardiol* 1987;**59**:13F–17F.
 30. McDevitt DG. Pharmacologic aspects of cardioselectivity in a beta-blocking drug. *Am J Cardiol* 1987;**59**:10F–12F.
 31. Bristow MR. beta-adrenergic receptor blockade in chronic heart failure. *Circulation* 2000;**101**:558–569.
 32. Weber MA. The role of the new beta-blockers in treating cardiovascular disease. *Am J Hypertens* 2005;**18**:169S–176S.
 33. Van de Water A, Janssens W, Van Neuten J, et al. Pharmacological and hemodynamic profile of nebivolol, a chemically novel, potent, and selective beta 1-adrenergic antagonist. *J Cardiovasc Pharmacol* 1988;**11**:552–563.
 34. Baker JG. The selectivity of beta-adrenoceptor antagonists at the human beta1, beta2 and beta3 adrenoceptors. *Br J Pharmacol* 2005;**144**:317–322.
 35. Psaty BM, Smith NL, Siscovick DS, et al. Health outcomes associated with antihypertensive therapies used as first-line agents. A systematic review and meta-analysis. *JAMA* 1997;**277**:739–745.
 36. Freemantle N, Cleland J, Young P, Mason J, Harrison J. Beta blockade after myocardial infarction: Systematic review and meta regression analysis. *BMJ* 1999;**318**:1730–1737.
 37. Hunt SA, Baker DW, Chin MH, et al. ACC/AHA Guidelines for the Evaluation and Management of Chronic Heart Failure in the Adult: Executive Summary A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Revise the 1995 Guidelines for the Evaluation and Management of Heart Failure): Developed in Collaboration With the International Society for Heart and Lung Transplantation; Endorsed by the Heart Failure Society of America. *Circulation* 2001;**104**:2996–3007.
 38. Willenheimer R, van Veldhuisen DJ, Silke B, et al. Effect on survival and hospitalization of initiating treatment for chronic heart failure with bisoprolol followed by enalapril, as compared with the opposite sequence: Results of the randomized Cardiac Insufficiency Bisoprolol Study (CIBIS) III. *Circulation* 2005;**112**:2426–2435.
 39. Effect of metoprolol CR/XL in chronic heart failure: Metoprolol CR/XL Randomised Intervention Trial in Congestive Heart Failure (MERIT-HF). *Lancet* 1999;**353**:2001–2007.
 40. Packer M, Bristow MR, Cohn JN, Colucci WS, Fowler MB, Gilbert EM, Shusterman NH. The effect of carvedilol on morbidity and mortality in patients with chronic heart failure. U.S. Carvedilol Heart Failure Study Group. *N Engl J Med* 1996;**334**:1349–1355.
 41. Robb OJ, Webster J, Petrie JC, Harry JD, Young J. Effects of the beta 2-adrenoceptor antagonist ICI 118,551 on blood pressure in hypertensive patients known to respond to beta 1-adrenoceptor antagonists. *Br J Clin Pharmacol* 1988;**25**:433–438.
 42. Prichard BN, Ross EJ. Use of propranolol in conjunction with alpha receptor blocking drugs in pheochromocytoma. *Am J Cardiol* 1966;**18**:394–398.
 43. Cruickshank JM. Beta-blockers continue to surprise us. *Eur Heart J* 2000;**21**:354–364.
 44. Beta-Blocker Evaluation of Survival Trial Investigators. A trial of the beta-blocker bucindolol in patients with advanced chronic heart failure. *N Engl J Med* 2001;**344**:1659–1667.
 45. Kaye DM, Lefkowitz J, Jennings GL, Bergin P, Broughton A, Esler MD. Adverse consequences of high sympathetic nervous activity in the failing human heart. *J Am Coll Cardiol* 1995;**26**:1257–1263.
 46. Winniford MD, Fulton KL, Hillis LD. Long-term therapy of paroxysmal supraventricular tachycardia: A randomized, double-blind comparison of digoxin, propranolol and verapamil. *Am J Cardiol* 1984;**54**:1138–1139.
 47. Geffner DL, Hershman JM. Beta-adrenergic blockade for the treatment of hyperthyroidism. *Am J Med* 1992;**93**:61–68.
 48. Plouin PF, Gimenez-Roqueplo AP. Pheochromocytomas and secreting paragangliomas. *Orphanet J Rare Dis* 2006;**1**:49.
 49. Ziegelstein RC. Acute emotional stress and cardiac arrhythmias. *JAMA* 2007;**298**:324–329.
 50. Tacoy G, Balcioglu AS, Arslan U, Durakoglugil E, Erdem G, Ozdemir M, Cengel A. Effect of metoprolol on heart rate variability in symptomatic patients with mitral valve prolapse. *Am J Cardiol* 2007;**99**:1568–1570.
 51. Schwartz PJ, La Rovere MT, Vanoli E. Autonomic nervous system and sudden cardiac death. Experimental basis and clinical observations for post-myocardial infarction risk stratification. *Circulation* 1992;**85**:177–191.
 52. Jacob S, Rett K, Henriksen EJ. Antihypertensive therapy and insulin sensitivity: Do we have to redefine the role of beta-blocking agents? *Am J Hypertens* 1998;**11**:1258–1265.
 53. Cruickshank JM. Are we misunderstanding beta-blockers. *Int J Cardiol* 2007;**120**:10–27.
 54. Steinberg HO, Baron AD. Vascular function, insulin resistance and fatty acids. *Diabetologia* 2002;**45**:623–634.
 55. Lithell H, Pollare T, Berne C, Saltin B. The metabolic and circulatory response to beta-blockade in hypertensive men is correlated

- to muscle capillary density. *Blood Pressure* 1992;**1**:20–26.
56. Myers MG, Hope-Gill HF. Effect of d- and dl-propranolol on glucose-stimulated insulin release. *Clin Pharmacol Ther* 1979;**25**:303–308.
 57. Efficacy of atenolol and captopril in reducing risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 39. UK Prospective Diabetes Study Group. *BMJ* 1998;**317**:713–720.
 58. Caro JF. Clinical review 26: Insulin resistance in obese and nonobese man. *J Clin Endocrinol Metab* 1991;**73**:691–695.
 59. Ladage D, Reidenbach C, Rieckeheer E, Graf C, Schwinger RH, Brixius K. Nebivolol lowers blood pressure and increases weight loss in patients with hypertension and diabetes in regard to age. *J Cardiovasc Pharmacol* 2010;**56**:275–281.
 60. Nuttall SL, Routledge HC, Kendall MJ. A comparison of the beta1-selectivity of three beta1-selective beta-blockers. *J Clin Pharm Ther* 2003;**28**:179–186.
 61. Dominguez LJ, Barbagallo M, Jacober SJ, Jacobs DB, Sowers JR. Bisoprolol and captopril effects on insulin receptor tyrosine kinase activity in essential hypertension. *Am J Hypertens* 1997;**10**:1349–1355.
 62. Seguchi H, Nakamura H, Aosaki N, Homma Y, Mikami Y, Takahashi S. Effects of carvedilol on serum lipids in hypertensive and normotensive subjects. *Eur J Clin Pharmacol* 1990;**38 Suppl** 2:S139–S142.
 63. Poirier L, Cleroux J, Nadeau A, Lacourciere Y. Effects of nebivolol and atenolol on insulin sensitivity and haemodynamics in hypertensive patients. *J Hypertens* 2001;**19**:1429–1435.
 64. Chobanian AV, Bakris GL, Black HR, et al. Seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *Hypertension* 2003;**42**:1206–1252.
 65. Arch JR. beta(3)-Adrenoceptor agonists: Potential, pitfalls and progress. *Eur J Pharmacol* 2002;**440**:99–107.
 66. Brixius K, Bloch W, Pott C, et al. Mechanisms of beta 3-adrenoceptor-induced eNOS activation in right atrial and left ventricular human myocardium. *Br J Pharmacol* 2004;**143**:1014–1022.
 67. Westhoff TH, Franke N, Schmidt S, Vallbracht-Israng K, Zidek W, Dimeo F, van der Giet M. Beta-blockers do not impair the cardiovascular benefits of endurance training in hypertensives. *J Hum Hypertens* 2007;**21**:486–493.
 68. Gullestad L, Hallen J, Medbo JI, Gronnerod O, Holme I, Sejersted OM. The effect of acute vs chronic treatment with beta-adrenoceptor blockade on exercise performance, haemodynamic and metabolic parameters in healthy men and women. *Br J Clin Pharmacol* 1996;**41**:57–67.
 69. Van Baak MA. Beta-adrenoceptor blockade and exercise. An update. *Sports Med* 1988;**5**:209–225.
 70. Ades PA, Gunther PG, Meyer WL, Gibson TC, Maddalena J, Orfeo T. Cardiac and skeletal muscle adaptations to training in systemic hypertension and effect of beta blockade (metoprolol or propranolol). *Am J Cardiol* 1990;**66**:591–596.
 71. Van Bortel LM, van Baak MA. Exercise tolerance with nebivolol and atenolol. *Cardiovasc Drugs Ther* 1992;**6**:239–247.
 72. Beloka S, Gujic M, Deboeck G, et al. Beta-adrenergic blockade and metabo-chemoreflex contributions to exercise capacity. *Med Sci Sports Exerc* 2008;**40**:1932–1938.
 73. Houssiere A, Najem B, Ciarka A, Velez-Roa S, Naeije R, van de Borne P. Chemoreflex and metaboreflex control during static hypoxic exercise. *Am J Physiol Heart Circ Physiol* 2005;**288**:H1724–H1729.
 74. Cogliati C, Colombo S, Ruscone TG, et al. Acute beta-blockade increases muscle sympathetic activity and modifies its frequency distribution. *Circulation* 2004;**110**:2786–2791.
 75. Ji LL, Lennon DL, Kochan RG, Nagle FJ, Lardy HA. Enzymatic adaptation to physical training under beta-blockade in the rat. Evidence of a beta 2-adrenergic mechanism in skeletal muscle. *J Clin Invest* 1986;**78**:771–778.
 76. Miura S, Kai Y, Kamei Y, Ezaki O. Isoform-specific increases in murine skeletal muscle peroxisome proliferator-activated receptor-gamma coactivator-1alpha (PGC-1alpha) mRNA in response to beta2-adrenergic receptor activation and exercise. *Endocrinology* 2008;**149**:4527–4533.
 77. Miura S, Kawanaka K, Kai Y, Tamura M, Goto M, Shiuchi T, Minokoshi Y, Ezaki O. An increase in murine skeletal muscle peroxisome proliferator-activated receptor-gamma coactivator-1alpha (PGC-1alpha) mRNA in response to exercise is mediated by beta-adrenergic receptor activation. *Endocrinology* 2007;**148**:3441–3448.
 78. Chua TP, Clark AL, Amadi AA, Coats AJ. Relation between chemosensitivity and the ventilatory response to exercise in chronic heart failure. *J Am Coll Cardiol* 1996;**27**:650–657.
 79. Schneider DA, McEniery MT, Solomon C, Jurimae J, Wehr MS. Ventilatory and hyperkalemic responses to incremental exercise after propranolol treatment. *J Appl Physiol* 1994;**77**:1907–1912.
 80. Conway MA, Petersen ES. Effects of beta-adrenergic blockade on the ventilatory responses to hypoxic and hyperoxic exercise in man. *J Physiol* 1987;**393**:43–55.
 81. Jilka SM, Joyner MJ, Nittolo JM, Kalis JK, Taylor JA, Lohman TG, Wilmore JH. Maximal exercise responses to acute and chronic beta-adrenergic blockade in healthy male subjects. *Med Sci Sports Exerc* 1988;**20**:570–573.
 82. Mutlu GM, Koch WJ, Factor P. Alveolar epithelial beta 2-adrenergic receptors: Their role in regulation of alveolar active sodium transport. *Am J Respir Crit Care Med* 2004;**170**:1270–1275.
 83. Piepoli MF, Guazzi M, Boriani G, et al. Exercise intolerance in chronic heart failure: mechanisms and therapies. Part I. *Eur J Cardiovasc Prev Rehabil* 2010;**17**:637–642.
 84. Agostoni P, Contini M, Magini A, et al. Carvedilol reduces exercise-induced hyperventilation: A benefit in normoxia and a problem with hypoxia. *Eur J Heart Fail* 2006;**8**:729–735.
 85. Baker JG. The selectivity of beta-adrenoceptor agonists at human beta1-, beta2- and beta3-adrenoceptors. *Br J Pharmacol* 2010;**160**:1048–1061.
 86. Bond RA, Spina D, Parra S, Page CP. Getting to the heart of asthma: Can “beta blockers” be useful to treat asthma? *Pharmacol Ther* 2007;**115**:360–374.