


ADVANCING MEASUREMENT OF PATIENT-CENTERED OUTCOMES AND QUALITY METRICS WITH ELECTRONIC HEALTH RECORDS

Tina Hernandez-Boussard, PhD, MPH, MS

Director, Surgical Health Services Research Unit

Assistant Professor Surgery & Medicine (BioMedical Informatics)



Disclosures

- Research reported in this presentation was supported by the National Cancer Institute of the National Institutes of Health under Award Number R01CA183962. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health



Background


- National Initiatives to improve quality of healthcare delivered in the US - Affordable Care Act (ACA)
 - Value-based payment modifiers (Section 3007)
 - Patient-centered care initiatives (Section 10602)
 - Accountable Care Organizations (Section 3022)
 - Coordinated care; Information exchange
- Patient-Centered Outcomes Research Trust Fund
 - PCORI - \$3.5B through 9/19



Accurate quality measurement is essential – and must include patient centered outcomes


Patient-Centered Outcomes (PCOs)

- POCs are outcomes from medical care that are important to patients which can not be measured by lab values or diagnostic tests
- Current government initiatives highlight the need to include PCOs in healthcare evaluations
- Prostate Cancer Outcomes
 - Prostate cancer treatments have different risks of patient-centered, such as urinary incontinence and erectile dysfunction
 - Current treatment guidelines regarding these outcomes have been inconclusive
 - Lack of clear evidence



Patient-Centered Outcome Evaluations Limited


- Relevant data are not captured in structured formats
- Existing methods to algorithmically annotate EHR free-text are limited
 - Interoperability is a problem
- Current assessments of patient-centered outcomes research rely on
 - Manual chart review
 - Labor intensive
 - size limited
 - Administrative data
 - Lack clinical granularity
 - Lack patient centered outcomes
 - Patient Surveys
 - Select population
 - Not readily available
 - Size limiting



Electronic Health Records (EHRs)

- EHRs Offer New Opportunities for Quality Assessment
 - Improve quality, safety, efficiency, and reduce health disparities.
 - Engage patients and family
 - Improve care coordination and population and public health
- CMS has identified EHR Meaningful Use (MU)
 - Accessibility (MU-1)
 - National effort to increase EHR adoption
 - Opens new opportunities to facilitate comprehensive assessment of healthcare delivery
 - Granular Data (MU-2)
 - EHRs contain detailed information about care processes and outcomes
 - Improve healthcare delivery (MU-3)
 - *Future of EHRs*

Just beginning to tap into the wealth of information contained in EHRs!



Hurdles Using EHRs


- Comprehensive assessment of EHRs hindered due to methodological reasons
 - Data are not organized according to patient and treatment – EHRs are not intended for research
- ~80% of data are entered as free-text in clinicians' notes
- Shared algorithms to use data with EHRs do not exist
 - Site or system-specific algorithms

While some studies have assessed 1-2 quality metrics, comprehensive quality assessment using EHRs is lacking



Research Question


- How can we use EHRs to improve measurement of quality indicators and patient-centered outcomes/metrics?
 - Efficiently and accurately measure patient-centered quality metrics using data stored in EHRs
 - Use case: Urinary incontinence in localized prostate cancer patients



Methods

- Study Population
 - Male patients seen at the Stanford Medical Center, 1995-2013
- Inclusion criteria
 - ICD-9 code for prostate cancer (185)
 - Prostatectomy procedure
 - ICD-9 and CPT codes
 - Open, laparoscopic, robotic
- Exclusion criteria
 - No clinical note in their EHR (narrative text)


2252 Prostate cancer patients with EHR note



Hurdle 1: EHR-Data Organization


- Develop EHR-Based Database
 - Identify cancer patients in EHRs using ICD-9 codes
 - Confirmed in state cancer registry
 - Map structured elements of EHR to relational database
 - Tables for patient, diagnoses, procedure, etc.
 - Each patient visit contains link to clinicians' notes as free text

Integrates complex, longitudinal data from multiple electronic medical records and registries and provides a rich, validated resource for research on oncology care.




Hurdle #2: Assessing Free Text

- Processes free-text into matrices that indicate which medical terminologies were mentioned in each patient encounter
 - Ontology-based dictionaries
 - Natural language processing technologies
 - Negation, past medical history, family history, temporal sequencing
 - Standard measures to test accuracy
- Matrices have three dimensions
 - Rows are Patients
 - Columns are Medical Terms
 - “Sheets” are Time (different episodes of care)
- Matrices are used for further analyses




Hurdle #3: EHR Shared Algorithms

- Electronic Phenotypes
 - Electronic algorithms used to identify characteristics of patients within healthcare data
 - Team of clinical domain experts, bioinformaticians or NLP experts, and biostatisticians
- Electronic Quality Metric Phenotypes
 - Structured series of components that define an outcome or cohort of interest
 - Map quality metric terms to existing vocabularies and ontologies
 - Define temporal structure of elements



Quality Phenotype

- Pretreatment assessment of urinary incontinence




B. **Quality Phenotype: Pretreatment Assessment of UI**

Key Features
Incontinent
Leaky bladder
Urinary incontinence
Nocturnal enuresis
Wears diaper

Diagnosis of Prostate Cancer
NEXT
<i>ICD-9 Code BEFORE Treatment</i>
788.3 - Urinary Incontinence
OR
<i>Text Mention of Urinary Incontinence Before Treatment</i>
Urinary Incontinence (SNOMED: 165232002)
NEXT
Prostate Cancer Treatment

Linked to Diagnosis codes

Linked to Procedure codes
And term mentions




Results

- 2252 Prostate cancer patients with EHR note
- Assessment of Urinary Incontinence
 - Pre-operatively
 - Recording of risks associated with treatment
 - Post-operatively


	Open	Robotic	Other	Total
Postoperative Urinary Incontinence, Data Mining	19.7%	12.7%	29.3%	22.4%
Postoperative Urinary Incontinence, Coded Data	<1%	<1%	<1%	0.21%
Postoperative Urinary Incontinence, Total	19.7%	12.7%	29.5%	22.6%

Display of Urinary Incontinence in EHRs




Workflow performance outcome: precision: 98%; sensitivity: 97%; F1: 98%

Workflow performance discussion: precision: 82%; sensitivity: 93%; F1: 96%




Conclusions

- EHRs offer opportunities to enhance quality measurement
 - Add capability to include metrics on patient-centered outcomes
- We overcome many obstacles to use EHRs
 - Enhance accuracy of data through development of EHR-database linked to patient registries
 - Developed tools to annotate EHR free-text
 - Create electronic phenotype algorithms to standardize EHR-measurements
- EHRs are an untapped source of data
 - Patient-centered outcomes research
 - Quality assessment
 - Provider documentation and adherence to guidelines




Discussion

- Evidence of Research feasibility
 - Patient reported outcomes exist in EHRs
 - Accuracy of our workflow
- Potential Impact
 - Begin to reliably assess aspects of patient-centered care
 - Generate evidence for important comparative effectiveness research
 - Previous studies on PCOs have had limited generalizability
 - Identify value-based payments that are representative of patient values



Stanford Team Members

- Jim Brooks, Urology
- Doug Blayney, Cancer Center
- Nigam Shah, Biomedical Informatics
 - Shah Lab
 - Anna Bauer-Mehren
 - Suzanne Tamang



Thank you!

- Questions?
 - boussard@stanford.edu
- Funding support:
 - NCI: R01CA183962

