

---

# Index

## A

- $\alpha 4\beta 2$  nAChRs, 166, 168
- $\alpha 7$  nAChR receptors, 168
- $\alpha 7$  nAChR subunit knockout mice, 159
- abstainers (nonsmokers)
  - characteristics of, 22
  - inclusion *versus* exclusion of, 222–223, 235, 323
  - nicotine patches, 377
- abstinence
  - in behavioral economics effects, 420
  - deficits, 350–351
  - effects of, 405, 408
  - event-related potential, 429
- abstinence-induced craving, 438
  - effects of, 405
  - measurements, 403
  - during smoking cessation, 441
- $\alpha$ -bungarotoxin binding, 165, 167
- ACE model, 249–250
- acetaldehyde (alcohol studies), 86–87
- acoustic startle reflex. *See* startle response
- active avoidance, 435
- acute stress mimicry of withdrawal symptoms, 448
- acute tolerance, 162–163
- ADA (Americans with Disabilities Act), 49
- Add Health (National Longitudinal Study of Adolescent Health), 197–198, 261, 517
- addiction. *See also* nicotine dependence;  
substance use
  - clinical manifestations of, 79
  - as dependence, 77
  - DRD2* gene and, 32
  - versus* habituation, 24
  - models of, 293
  - versus* smoking, 24
  - stress and, 36
- additive components, 511
- adenosine knockout mice, 159
- adenosine systems, in nicotine reinforcement,  
159–160
- ADH1\*2* alleles (alcohol studies), 86–87
- ADHD. *See* attention deficit hyperactivity disorder
- adipose tissue, nicotine concentrations in, 145
- ad libitum (ad lib) self-administration, 415–416,  
419
- administration, 145–149, 418. *See also* self-  
administration
  - in drinking water, 148, 172
  - intravenous, 146, 156, 410
  - oral, 148, 154, 412
- adolescent(s)
  - alcohol use by, 516
  - cognitive control, 358, 380
  - delay discounting choices, 350
  - depression in, 351–352
  - event-related potential, 429
  - extraversion in, 349
  - first mood effects, 375–376
  - neuroticism in, 351, 352
  - nicotine deprivation learning deficits, 436
  - nicotine response in, 194–195
  - novelty seeking behavior, 27, 348
  - P300 amplitude in, 360
  - physiological changes in, 589
  - protective factors, 343
  - research limitations, 367
  - social influences on, 346–347, 517
  - substance-use vulnerability of, 195, 199, 200–  
201, 212, 233–234, 261, 292–293
  - transition to adulthood, 195
  - use of genetic information by, 46–47
- adolescent developmental trajectories, 189–235,  
592
  - age of smoking onset, 200–201
  - empirically identified, 202–214
  - example of, 223–233
  - future research directions, 233–234
  - psychopathology, 191–202, 292–293
  - statistical models, 214–223
  - substance use, 295–296 (*See also* substance-  
use comorbidity; *specific substance*)
- adolescent nicotine dependence, 191–195
  - animal models of, 155, 194–195
  - biological vulnerability for, 100, 193–195,  
200–201, 233
  - future research directions, 233
  - genetic studies of, 86, 264–266, 342
  - individual symptoms of, 192
  - measurement of, 192, 230–231, 264
  - time and exposure required for, 192–193
  - withdrawal symptoms, 192
- adolescent smoking, 371
  - antisocial behavior and, 200, 202, 211, 232
  - environmental influences on, 196–197
  - ethnic differences in, 213–214, 279

- gender differences in, 196, 199, 260, 263–264, 342
- gene-environment interactions in, 197–200, 259
- genetic research on, 195–200, 259–269
- heterogeneity in, 190, 233
- as indicator of adult nicotine dependence, 230–231
- latency between cigarettes, 371
- molecular genetic studies of, 198–199
- parental smoking and, 196–197, 200
- peer smoking and, 197
- prevalence of, 191
- twin studies of, 196, 259–262
- adolescent smoking initiation, 191
- age range in, 261–262
- heterogeneity of, 196, 201–202, 233
- progression to dependence, 341
  - genetic studies of, 263–264
  - rate of acceleration, 201–202
- psychosocial factors, 200, 202, 211
- risk profile, 211–212, 232
- shared environmental factors in, 260–261, 264, 280
- adoption studies, 196, 279
- ADRA1A* gene, 42
- adulthood, transition to, 195
- adult nicotine dependence, adolescent smoking indicators of, 230–231
- adult-onset events, 100
- adult smoking phenotype, limitations of, 190
- advertising
  - costs of, 21
  - in movies, 7, 20, 523
  - novelty seeking as response to, 348
  - protobacco, 7, 20, 30, 348
  - smoking index variable and, 30
- aerosols, nicotine, 147
- affective coping, 112–113
- affective response, 373–376
  - future research directions, 456
  - physiological measures of, 377–378, 445–446
  - regulation of, 358, 403, 443–449
- affiliation/empathy system, 362
- African Americans
  - adolescent smoking in, 213–214
  - genotypes linked to dependence, 47
  - linkage study focused on, 267
- age effects, 170–171
  - factor loadings by, 271–276
  - in smoking initiation assessment, 261–262, 279, 281, 322
  - in substance-use comorbidity, 322–323
- age-gene-environment interactions, 589
- age of onset, 26, 100, 371–372
  - developmental trajectories by, 200–201
- age-related macular degeneration, 46
- age-specific risk, measurement of, 35
- aggregate effects in complex pathways, 541
- aggression, 357
- agonists (activators), 143
- AHe mice, 435
- AIC (Akaike Information Criterion), 274, 312
- A inbred mice, 154, 165, 412, 422
- A/J mice, 435
- A/J×NMRI cross-bred mice, 154, 412
- Akaike Information Criterion (AIC), 274, 312
- alcohol use
  - adolescent, 516
  - in ATBC analysis, 496
  - Edwards's theory of, 82–83
  - Iowa gambling task as predictor, 350
  - nasal spray use correlated with, 373
  - in NHANES III analysis, 502
  - policies influencing, 294–295
  - as secondary reinforcement, 413
  - side effects, 86–87
  - tobacco use concurrent with, 98, 290, 296–298
    - empirical examples of, 307–323, 496, 502
    - health effects of, 290
    - modeling, 299–305
    - nicotine-dependence correlation, 369, 406, 411–412, 420
  - trajectories of, 295
- ALDH2\*2* alleles (alcohol studies), 86–87
- alertness, 361–362
- allele(s)
  - identical by descent (IBD), 257, 258
  - variants of, 554
- allele frequency, 48, 258
- alpha subunits, 153
- Alpha-Tocopherol, Beta-Carotene Cancer Prevention Study (ATBC), 487, 494–497
- alternative reinforcement, 417
- Americans with Disabilities Act (ADA), 49
- amphetamine, 412
- analysis of variance (ANOVA), 215
- analytic methods. *See also specific method*
  - developmental trajectories, 214–223
  - molecular genetic studies, 257–259
  - phenotypic research, 96–103, 113–118

anger, palliative effects on, 357  
 animal studies. *See also* mouse models; rat models; *specific strain or study*  
     ad lib administration in, 415  
     adolescent nicotine exposure, 194–195, 589  
     affective regulation, 443  
     dependence, 134–135  
     fetal nicotine exposure, 357  
     impulsivity research in, 449–450  
     reward studies, 372–373  
     transfer to, 350  
 ANOVA (analysis of variance), 215  
 antagonists (inhibitors)  
     CB1 receptors, 160  
     for mu opioid receptors, 160  
     muscarinic receptors, 141  
     nicotine as, 143  
     for nicotine dependence, 159  
 anti-inflammatory effects of nicotine, 148–149  
 antinociception, 162  
 antisaccade task, 359  
 antisocial behavior  
     adolescent smoking and, 200, 202, 211, 232  
     substance use and, 292–293, 304  
 antitobacco stimuli, 20  
 anxiety  
     adolescent smoking and, 201  
     nicotine linkage with, 352–353, 445  
 anxiogenic effects of nicotine, 168  
 apolipoprotein E testing, 48  
 approach, *versus* impulsivity, 378–379  
 approach-related risk, 339, 346–349, 362  
 arousal, 378  
*ASN40ASP* polymorphism, 407  
 aspartame, 412  
 association analysis, 258–259, 268–269, 280  
 assortative mating, 251, 259, 280  
 ATBC (Alpha-Tocopherol, Beta-Carotene Cancer Prevention Study), 487, 494–497  
 ATR (Australian Twin Registry), 197, 260–261, 267  
 attention, 361, 378, 432–434  
 attentional bias, 440, 456  
 attention deficit hyperactivity disorder (ADHD), 354–357  
     adolescent smoking and, 201, 211  
     as risk factor, 350, 433  
     substance use and, 292–293, 304  
 attrition biases, 224  
 Australian Twin Registry (ATR), 197, 260–261, 267  
 Automaticity subscale, 90–91

aversive mood symptoms, 443  
 avoidance-related risk, 339, 351–354, 362

## B

backcrossed mice, 151  
 bacterial contamination, during administration, 145  
 BALB/cBy mice, 164–165, 430–431, 435  
 Barratt Impulsivity Scale, 450  
 Bayes factors, 552, 570  
 Bayesian analysis, 548–549, 570, 572  
 Bayesian False Discovery Probability (BFDP), 573  
 Bayesian Information Criterion (BIC), 310, 312  
 Bayesian model averaging, 117  
 Bayes model, 543  
 BAY K 8644, 167  
 behavior  
     analysis of, 163–164  
     antisocial  
         adolescent smoking and, 200, 202, 211, 232  
         substance use and, 292–293, 304  
     drug-motivated, 404  
     measurements of, 103  
     nicotine and, 151–157  
     phenotypes, 171  
     response systems, 362  
     smoking indices, 80  
     substance-use comorbidity and, 322  
     tolerance, 168  
     traits, 344  
     undercontrol, 98–99  
 behavioral economics, 417, 420, 454  
 behavioral genetics  
     phenotypes in, 492  
     of self-administration, 153–155  
     in social context, 514–518  
 behavioral modeling  
     methodological issues with, 247–248  
     of parental smoking, 246  
 BFDP (Bayesian False Discovery Probability), 573  
 BIC (Bayesian Information Criterion), 310, 312  
 bioavailability of nicotine, 7, 20  
 biochemical indices of smoking, 80, 415  
 biochemical measures of self-administration, 97  
 biochemical pathways, 561  
 BioCyc, 561  
 biological pathways, candidate, 105  
 biological plausibility  
     affective regulation, 443–445

- attention/vigilance, 432–433
- craving, 438–439
- event-related potential, 427–428, 429
- impulsivity, 449–450
- mood effects, 374
- reinforcement, 368–369, 410–414
- resting EEG activity, 425–426
- rewards, 372–373, 421–423
- startle response, 430–431
- working memory, 434–436
- biological vulnerability, in adolescent nicotine dependence, 194–195
- biomedical ontologies, 561
- biometric factor model, 253
- biometric modeling, 37–45, 86, 88–89, 514
- BioPAX Ontology, 561
- bitter taste, 84, 148
- BKW mice, 443
- blood pressure, 448–449
- $\beta 2$  nAChRs, 153
- $\beta 2$  nAChR subunit knockout mice, 159, 166
- Bonferroni correction, 546, 573
- brain
  - nicotine concentrations in, 145
  - stimulation, 373
  - upregulation in, 144
  - of young people, 589
- brain imaging, 360
- breakpoint
  - drug use, 418
  - nicotine use, 420
  - preferring to wait, 350
- breast milk, nicotine concentrations in, 145
- breeding
  - mice, 150–151, 411
  - rats, 411
- Brown University Transdisciplinary Tobacco Use Research Center (TTURC), 89, 521, 526
- BUB/Bn mice, 165, 412
- BUB inbred mice, 154
- buzz, 374

## C

- calcium, 136, 139–140
- calcium calmodulin protein kinase II, 168
- calcium channel blockers, 162, 167, 168
- calcium signaling, 167–168
- Canadian National Longitudinal Survey of Children and Youth, 212
- cancer genetics, 50
- candidate biological pathways, 105
- candidate gene studies, 40–45
  - adolescent smoking, 342
  - discordant phenotype associations, 36
  - epigenetic differences, 37
  - linkage analysis, 32, 40–42, 267–268, 280–281, 589
  - multivariant data, 543–546
  - nicotine dependence, 42
  - prior knowledge of, 559
  - smoking association with, 24
  - SNP relationships over, 545
  - substance-use comorbidity, 325
  - trait pathways, 553
  - variants, 25, 35
- candidate neural systems, 343–346
- cannabis. *See* marijuana use
- Card Arranging Reward Responsivity Objective Test, 350–351
- CART (cocaine- and amphetamine-regulated transcript), 36
- cases, in association analysis, 258
- catechol-*O*-methyl transferase (*COMT*) gene, 269, 408–409, 430
- categorical covariates, 276–278
- causal contingent common (CCC) pathway model, 253, 263, 278–279
- causal differences between groups, 492
- causal paths
  - phenotypic research, 102, 107–110
  - sensitivity to, 105
  - smoking as, 352
  - types of, 111–113
- CBA mice, 435
- CB1 knockout mice, 160, 422
- C57BL/6 inbred mice
  - conditioned place preference in, 159, 422
  - dose-dependent effects, 165
  - fear conditioning in, 169, 432
  - five choice serial reaction time task in, 433
  - nicotine consumption, 155, 159
  - nicotine-dependence risk, 161
  - nicotine effects in, 435
  - novelty-seeking behavior in, 156
  - oral self-administration in, 412–413
  - prepulse inhibition in, 430–431
  - strain comparisons, 153–154, 164–165
  - in tolerance, 167
- C57BL/10 mice, 435

- C57BR/cd mice, 435
- CCC (causal contingent common) pathway model, 253, 263, 278–279
- CCK gene, 269
- CD-1 mice, 443
- CellDesigner, 561
- cell signaling, 161, 167–168
- Center for Antisocial Drug Dependence, 260
- Centers for Disease Control and Prevention, 498
- central nervous system (CNS), 344, 348, 353, 354
- centroid. *See* mean
- chain smoking, 37
- children
- effortful control in, 358
  - secondhand smoke from parental smoking, 246
  - self-control in, 344
  - sleep problems in, 357
  - smoking by, 371
- children of twins (COT) design, 251–252, 279, 511
- C3H inbred mice
- dose-dependence, 165
  - nicotine effects in, 431, 435
  - oral self-administration, 412
  - strain comparisons, 154, 164–165
  - tolerance in, 162
- chippers (light smokers)
- adolescent, 213, 232
  - delay discounting, 350
  - genetic factors and, 29
  - versus* heavy smokers, 43, 81, 83–84, 90, 94
  - prevalence of, 193
- choice procedure, 369–370, 416, 419–420
- cholecystokinin (CCK) gene, 269
- Christchurch, New Zealand (CNZ) study, 266
- CHRNA2 gene, 42, 376–377
- CHRNA3 gene, 376–377
- CHRNA4 gene, 268, 407–409, 412
- CHRNA5 gene, 342, 376–377
- CHRNA7 gene, 408, 409, 428
- CHRNA5-A3-B4 haplotypes, 100, 407
- CHRNA5-CHRNA3-CHRNA4 nicotinic receptor genes, 43
- CHRNA2 gene, 268, 407, 408, 409
- CHRNA3 gene, 342
- CHRNA3-CHRNA6 nicotinic receptor locus, 43
- chronic exposure, 405, 449
- chronic smoker endophenotypes, 403, 404–406
- affective regulation, 443–449
  - cognitive control, 432–438
  - craving, 438–443
  - electrophysiological measures, 425–432
  - future research directions, 452–457
  - impulse control, 449–452
  - investigation rationale, 406–410
  - motivational mechanisms, 410–424
  - summary/conclusions, 457–458
- chronic tolerance, 163
- CIDI (Composite International Diagnostic Interview), 81
- cigarette(s)
- availability of, 99
  - consumption of, 4, 20
  - design of, 7, 20
  - pricing of, 20, 21, 520
  - vendor locations, 521, 522
- Cigarette Dependence Scale, 79n
- Cigarette Evaluation Scale, 423
- cigarettes per day (CPD)
- in ATBC analysis, 496
  - with CYP2A6 gene variants, 418
  - delay discounting correlation, 452
  - as dependence measure, 79, 80
  - in factor analysis, 88–89
  - in NHANES III analysis, 502
  - nicotine metabolism and, 406
  - predictive value of, 80
  - recall reliability, 26–27
  - as smoking cessation predictor, 81, 413
  - in TUS-CPS analysis, 498, 499
- cigarettes per month, 91
- class extraction, model misspecification related to, 222
- classic dependence criteria, 84, 86
- classification bias, 493
- class I–IV phenotypes, 28–31
- class membership, for familial resemblance, 256–257
- clinical preventive services, 4
- Clinical Research Support System, 415
- cluster analysis
- developmental trajectories, 211, 217–223
  - discrete *versus* continuous phenomena in, 219
  - static *versus* dynamic, 219–220, 232
  - within-class variability in, 220–221, 233
- CNS (central nervous system), 344, 348, 353, 354
- CNZ (Christchurch, New Zealand) study, 266
- cocaine, 374, 456–457
- cocaine- and amphetamine-regulated transcript (CART), 36
- coexpression of receptors, 141, 142

- COGA (Collaborative Studies on Genetics of Alcoholism), 266
- cognition changes, 163–164
- cognitive control, 403, 424–425, 432–438
  - during adolescence, 380
  - alertness in, 361
  - electrophysiological measures, 425–432
  - as endophenotype measurements, 358, 360
  - impairment, 113
  - impulsivity and, 112, 378–379
  - physiological basis of, 381
- cognitive deficits reversal, 434–435
- cognitive measures, of craving, 440
- cohorts
  - effects of, 515
  - research models for, 519
- Collaborative Studies on Genetics of Alcoholism (COGA), 266
- colorectal polyps, 542
- commercial testing, 50
- common pathway model, 252, 270
- common-vulnerability model, 291–294, 306, 315–316
- communication of genetic findings, issues in, 45–50
- comorbidity
  - psychiatric (*See* psychiatric comorbidity)
  - substance-use (*See* substance-use comorbidity)
- complementary dimensions of dependence, 97
- complex traits
  - defining features, 31
  - genetic factors in, 35
  - genome-wide association studies for, 46
  - multiple determinations of, 22
  - replication difficulties, 45
  - sensitive genetic measures, 341
  - similarity of, 24–25
- Composite International Diagnostic Interview (CIDI), 81
- compulsive smoking, 80
- computational symbolic theory, 559–560
- COMT* gene, 269, 408–409, 430
- concurrent choice procedure, 416
- conditional independence, 256
- conditional triggers to smoke, 588
- conditioned place preference (CPP)
  - biological plausibility of, 372–374, 421
  - in mice, 157–158, 161, 422
  - in rats, 423
- conditioning, contextual, 163–164
- conduct disorder
  - adolescent smoking and, 201, 211, 232
  - definition of, 357
  - substance use and, 292–293, 304
- confounding factors, 32
- consensus, across indicators, 77
- constitutional hypothesis, 24
- construct
  - definition of, 75
  - emerging, 520
  - proximal, 522
  - refinement of, 25–27
- construct properties, 75
- construct validation, 33, 34, 35, 75–78
- consumption level
  - during adolescence, as indicator of adult dependence, 230–231
  - adolescent nicotine dependence and, 193
  - assessment of, 415, 418
- contamination, during administration, 145
- context-sensitive physiological measures, 363
- contextual conditioning, 163–164
- contingency table, smoking-drinking, 313, 315–316
- contingent factors, 247
- continuant, 541, 559
- continuous factors, 256
- continuous-level information, for behavioral modeling, 247
- continuous performance task (CPT), 362, 378, 433
- continuous phenomena, *versus* discrete phenomena, 219
- control, endophenotype measures of, 357–362
- control-related risk, 339, 354–357, 362
- controls, in association analysis, 258
- copy-number variation, 36
- core criteria, 76, 77
- core dependence dimension, 87–95
- core factors, 97, 168
- core strategy, 510
- cortisol, 447
- cosegregation of smoking behaviors, 24
- cost(s)
  - genetic testing, 50
  - smoking cessation, 593
  - tobacco use, 4, 21
- COT (children of twins) design, 251–252, 279, 511
- cotinine
  - biological activity of, 148
  - clearance factors, 39–40

as nicotine by-product, 406  
 covariance, 32, 95–96, 341–342  
   in latent growth curve models, 254–255  
   in structural equation modeling, 248, 250  
 CPD. *See* cigarettes per day  
 CPP. *See* conditioned place preference  
 CPT (continuous performance task), 362, 378, 433  
 craving, 438–443  
   abstinence-induced (*See* abstinence-induced craving)  
   in adolescents, 192  
   cue-induced (*See* cue-induced craving)  
   dopaminergic systems associated with, 111  
   effects of, 405  
   measurement of, 403, 439–440  
 Craving subscale, 90–91  
 CREB activation, 160–161  
 CREB mice, 422  
 critical constituents, 25  
 cross-cultural differences, 515–516  
 cross-sectional studies, of substance-use comorbidity, 291  
 cross-species analysis, 348  
 cue(s)  
   in animal studies, 411  
   relapse, 99  
   reward signaling, 350  
 cue-induced craving, 438–439  
   with dependence, 441–443  
   effects of, 405  
   future research directions, 456–457  
   measurements, 403  
   physiological basis of, 111, 588  
   procedures to elicit, 440–441  
   research limitations in, 456  
 cue-self-administration response, 94  
 cultural transmission  
   adolescent smoking and, 198  
   effects of, 515  
   as environmental factor, 21  
   phenotypic, 251  
   research models for, 519  
   twin studies of, 262  
 cumulative effects, of pathogens, 103  
 cumulative risk, measurement of, 35  
*CYP2A6* gene  
   adolescent smoking and, 199  
   analysis of, 554, 555  
   association analysis of, 268

coding for, 408–409  
 mood effects with, 376  
 in nicotine metabolism, 22, 39–40, 149, 342, 406  
 in smokers, 413  
 variants in, 371, 418  
*Cyp2a5* gene variants, 149, 155, 413  
*CYP2B6* gene, 39  
*CYP2D6* gene, 39  
 Cys-Cys pairs, 136  
 cytochrome P-450 (CYP) system, 268

## D

danger-alarm responses, 353, 354  
 data-mining techniques, 541–542  
 data sets  
   ATBC analysis, 495  
   NHANES III analysis, 500–501  
   smoking cessation analysis, 493–494, 494  
   TUS-CPS analysis, 498  
 DAT/SLC6A3 (dopamine transporter protein), 268  
 DBA/2 inbred mice  
   conditioned place preference, 422  
   dose-dependent effects, 165  
   IV self-administration, 156  
   nicotine consumption, 159  
   nicotine effects, 435  
   oral self-administration, 412  
   prepulse inhibition in, 430–431  
   strain comparisons, 153–154, 164–165  
   tolerance in, 162, 163  
 DDC (DOPA decarboxylase), 269  
 definition variables, in structural equation modeling, 248  
 dehydroepiandrosterone (DHEA), 447  
 delay discounting  
   definition of, 348  
   in impulsivity, 349, 449, 450–452  
   physiological basis of, 379  
   study subjects for, 350  
 deleterious effect prediction, 554  
 dependence. *See also* nicotine dependence  
   addiction as, 77  
   core features of, 592  
   definition of, 75  
   distal measures of (*See* distal measures)  
   maintenance of, 598  
   model evaluation of, 78  
   patterns, 94



- severity with comorbidity, 98
- study of, 24
- depression, 351–352
  - nicotine amelioration and, 443
  - smoking association with, 444
  - subthreshold, 352
- desensitization, 146
- developmental pathways, 100, 589
  - impact on trajectory study, 222
- developmental psychopathology, 191–202, 292–293
- developmental trajectories. *See* adolescent developmental trajectories
- DH $\beta$ E, 159, 169
- DHEA (dehydroepiandrosterone), 447
- diabetes mellitus, 46
- Diagnostic and Statistical Manual of Mental Disorders (DSM)* criteria
  - adolescent smoking, 192, 264
  - attention deficit hyperactivity disorder, 356
  - dependence, 37–38, 40, 81, 86
  - poor agreement with FTND, 25–26
  - scales in, 79
  - substance-use disorder, 291
  - as syndromal medical model, 80–81
- diagnostic criteria, heritability of, 30, 38
- diagnostic inferences, 77
- Diagnostic Interview Schedule (DIS), 501
- diary prompts/responses, 527, 528–529, 530
- Digit Span test, 433
- diltiazem, 168
- direct drug infusion, 152–153
- direct-to-consumer marketing, 50
- DIS (Diagnostic Interview Schedule), 81, 501
- disadvantaged youth, 520
- discrete phenomena, *versus* continuous phenomena, 219
- discrimination, against nicotine dependence, 46–47, 49
- discriminative validity, 77
- discussion groups (OBO Foundry project), 560
- Disease Ontology (DO), 560
- disease risk, 541, 550
- disinhibition, common trait of, 292–293
- disorder markers, 107
- dispositional tolerance
  - acquisition of, 162
  - versus* behavioral tolerance, 164
  - in tolerance, 168
- distal influence, *versus* proximal influence, 519–522
- distal measures
  - agreement among, 88
  - early *versus* mature states, 74
  - genetic mapping, 78–81, 86–87, 103–105
  - versus* proximal measures, 516
- distress tolerance, 448
- distributional assumptions
  - in growth curve models, 255
  - in structural equation modeling, 249
- DNA methylation, 36
- DNA sequences analysis, 554
- DO (Disease Ontology), 560
- DOPA decarboxylase (DDC), 269
- dopamine
  - in attention deficit hyperactivity disorder, 356
  - attention-vigilance associations with, 434
  - impulsivity linkage with, 451–452
  - inactivation of, 430
  - mesotelecephalic, 171
  - midbrain circuits, 348, 349, 350
  - in nicotine binding role, 406–407
  - in nicotine dependence, 43, 410
  - receptors, 156, 343
  - regulation of, 540
  - reinforcement role of, 374, 407
  - reward pathways, 342, 349, 352
  - signaling, 160
- dopamine  $\beta$ -hydroxylase (D $\beta$ H), 269
- dopamine hypothesis of dependence, 588
- dopaminergic neurons, in ventral tegmental area, 588
- dopaminergic systems
  - adolescent smoking and, 198–199, 201
  - association analysis of, 268–269
  - craving associated with, 111
- dopamine transporter protein (DAT/SLC6A3), 268
- dopamine transporter (*SLC6A3*) \*9-repeat allele, 441
- dose of nicotine, 161, 165
  - differences in, 367, 416
  - distribution of, 370
  - standardization, 152
- dose-response curve, 162, 444
- double variant haplotype, 545
- downregulation, 144–145, 588
- downstream processes, 81–82, 106
- DRD4 (dopamine receptor), 268
- DRD5 (dopamine receptor), 268
- DRD4* exon III polymorphism, 199



*DRD2* gene, 409  
 addiction association with, 32  
 coding for, 408  
 commercial testing for, 50  
 dependence association with, 407  
 in craving, 441  
 variants, 198  
*DRD4\*7*-repeat allele, 376  
 drinking. *See* alcohol use  
 drinking-water administration, 147–148, 172  
 drug addiction. *See* addiction; substance use;  
     *specific drug*  
 drug-motivated behavior, 404  
 drug response comparison, 490–492, 491  
 DSM. *See* *Diagnostic and Statistical Manual of*  
     *Mental Disorders* criteria  
 dual-trajectory model, of smoking-drinking  
     trajectories, 315–316  
 dynamic clustering  
     *versus* static clustering, 219–220, 232  
     within-class variability in, 220–221, 233

## E

early-emergent motive, 91  
 early-onset smokers  
     risk for persistence, 200, 212–213, 230–231  
     substance use and, 292, 296–297, 315, 320  
 early smoking experiences (ESE), 375  
 early tobacco exposure, 101, 155  
 ecological momentary assessment (EMA), 255,  
     525  
 economic deprivation, 520  
 educational attainment, adolescent smoking and,  
     227–228, 232  
 Edwards's theory of alcohol dependence syndrome,  
     82–83  
 EEA (equal environments assumption), 516–517  
 EEG (electroencephalogram), 354, 403, 425–427  
 effortful control, 358  
 elasticity of demand, 417  
 elation, 374  
 electroencephalogram (EEG), 354, 403, 425–427  
 electromyography (EMG), 431  
 electrophysiological measures, 378, 425–432  
 EMA (ecological momentary assessment), 255,  
     525  
 EMG (electromyography), 431  
 empirical-Bayes approach, 549, 551  
 empirical search strategies, 117

employment discrimination, 49  
 employment status, of hard-core smokers, 35  
 encoding prior knowledge, 571  
 endogenous cannabinoid systems, 159, 160  
 endogenous enkephalin system, 160  
 endogenous event-related potentials (ERPs),  
     427  
 endophenotypes, 5, 408–410  
     caveats, 110–111  
     characteristics of, 107–110  
     in chronic smokers (*See* chronic smoker  
         endophenotypes)  
     conceptual issues, 381–383  
     criteria, 413  
     disorders associated with, 106  
     future research directions, 455–457, 594  
     gene linking in, 347, 409  
     measurement of, 349–351, 353–354, 355,  
         357–362  
     motivational effects, 452, 454–455  
     in network models, 558  
     nicotine dependence, 409, 453  
     phenotypes associated with, 5, 33–34  
     pre-exposure risk, 340–347  
     in psychiatric genetics research, 25  
     replicability of, 27  
     transitional, 107, 108, 200, 233  
     types of, 340  
 enhanced clearance. *See* dispositional tolerance  
 entities, 541, 559, 560  
 environmental factors, 99–103. *See also* social  
     context; *specific factor*  
     adolescent smoking, 196–197  
     comorbidity, 99  
     cue-induced craving, 438–439  
     enrichment, 36  
     in experimentation, 31  
     gender differences in, 38  
     gene expression variation from, 36–37  
     genetic factors in, 35, 515  
     importance of, 29  
     linkage analysis, 258  
     measurement of, 35  
     in nicotine dependence, 22, 23  
     nicotine use, 158  
     relative contribution of, 30  
     in smoking decline, 20  
     substance use, 294–295  
     twin studies, 251, 262, 279, 280  
 environmental pathogens, 5, 25, 35

epidemiology, 31–37  
  future research in, 594  
  genetic, 257  
    extended, 250–252, 262–269, 279, 280  
  hierarchical modeling in, 570–571  
  perspectives from, 514  
  phenotypic definitions in, 493  
  public health outcomes in, 492  
  triangle, 513  
epigenetics, 36  
epistasis, 32–33  
epistemology, 74, 77  
equal environments assumption (EEA), 516–517  
equifinality, 191, 220  
ERPs (event-related potentials), 403, 427  
ESE (early smoking experiences), 375  
ethanol, 412. *See also* alcohol use  
ethnic differences, in developmental trajectories,  
  213–214, 279  
etiologial architecture, 510–511, 515–518, 519,  
  527, 530  
etiology  
  diverse, 82, 106  
  matrix of, 509  
  of phenotypic assay, 83  
  of symptoms, 78  
euphoria, 374  
Event Ontology (EVO), 560  
event-related potentials (ERPs), 403, 427  
EVO (Event Ontology), 560  
exchangeable classes, 551  
excitatory tone, 141  
executive function  
  cognitive control and, 379  
  definition of, 361  
  nicotine dependence and, 34  
Executive Order 13145, 49  
exogenous event-related potentials (ERPs), 427  
experimental design, basics of, 149–150  
experimentation  
  influences on, 31  
  progression from, 30  
exposure model, 365  
extended structural equation modeling (XSEM),  
  249  
extended twin family studies, 250–252, 262, 279,  
  280  
extra-nicotinic mechanisms, 156–157  
extraversion, 346, 348–349  
extreme group membership, 96, 97

extreme groups  
  alternatives to, 116–118  
  constructing, 114–116  
eyeblick response, 448

## F

factor analysis, 37–38, 265  
  consistency, 88  
  correlation among, 80, 87  
  *DSM-IV* correspondence with, 103  
factor loadings, 271–278  
factor mixture model (FMM), 256  
factor models, 256  
Fagerström Test for Nicotine Dependence (FTND),  
  79–80  
  as assessment tool, 405  
  dependence criteria, 37–38  
  FTQ as precursor to, 24  
  linkage analysis, 40–42, 589  
  poor agreement with *DSM*, 25–26  
  reliability and validity of, 26, 79–80  
  scales in, 79  
  visuospatial attention association with, 434  
Fagerström Tolerance Questionnaire (FTQ),  
  79–80  
  for adolescent smoking, 192, 230–231, 264  
  as physical dependence measure, 79  
  startle response inconsistency, 448  
  in susceptibility loci mapping, 24  
  test-retest reliability, 26  
false discovery rates (FDRs), 542, 546, 570, 573  
false positive reports, 570  
familial resemblance, class membership for,  
  256–257  
family-based studies  
  design of, 518  
  ecological momentary assessment in, 526–527  
  heritability documentation, 28–29  
  new methodologies in, 521  
family dysfunction scores, 33  
family environment, adolescent smoking and,  
  196–197  
family history analysis, example of, 225, 230,  
  232  
fast-ionotropic nicotinic receptors, 136  
FDRs (false discovery rates), 542, 546, 570, 573  
fear conditioning, 169, 432, 435  
fear responses, 353, 354  
feeder stream influences, 81–82, 106

female smokers. *See also* gender differences  
adolescent, 343  
blood pressure changes in, 448  
nicotine-dependence factors, 37, 38, 99  
*OPRM1* gene in, 419  
statistics on, 21  
twin studies, 515, 517  
fetal nicotine exposure, 357  
FHS (Framingham Heart Study), 266  
final common pathway, 82, 93, 106  
finite mixture model, 256  
Finnish Twin Registry, 262  
Finn Twin16-25 study, 307–323  
methods, 307–310  
results, 310–323  
first experience with smoking. *See* initial  
sensitivity  
first-stage estimates, 572  
Fisher, Ronald Aylmer, 22, 24  
five choice serial reaction time task (5CSRTT),  
432  
five-class solution, example of, 226  
fixed effects, 215–216  
flunarizine, 168  
FMM (factor mixture model), 256  
focused interaction testing framework, 542  
forced choice procedure, 416  
formal model, 541, 559  
Fosb knockout mice, 161  
Foundational Model of Anatomy, 560  
four-point Likert scale, 375  
Framingham Heart Study (FHS), 266  
F344 rats, 423  
FTND. *See* Fagerström Test for Nicotine  
Dependence  
FTQ. *See* Fagerström Tolerance Questionnaire  
future research. *See also specific topics*  
crosscutting issues, 595–596  
implications of, 588–591  
understanding, 588–596

## G

$\gamma$ -aminobutyric acid (GABA), 157, 406–407  
 $\gamma$ -aminobutyric acid receptors, 43  
GABA ( $\gamma$ -aminobutyric acid), 157, 406–407  
GABAergic interneurons, 142  
gateway theory of substance use, 292  
GAW (Genetic Analysis Workshops), 266  
gender differences. *See also* female smokers;  
male smokers  
adolescent smoking, 196, 199, 227–228, 260,  
263–264, 342  
animal studies, 155, 164  
cross-cultural, 515  
factor loadings by, 271–276  
nasal spray use, 373  
nicotine-dependence factors, 38, 99  
nicotine-dependence heritability estimates,  
279, 281  
*OPRM1* gene, 419  
smoking initiation, 267  
startle response, 378  
substance-use comorbidity, 304  
twin studies, 262, 517  
gender heterogeneity, 273–274, 276–278  
gene(s). *See also* candidate gene studies  
endophenotype linkage risks, 347, 409  
in nicotine dependence, 32, 43  
gene-environment interaction, 33, 515  
adolescent smoking, 197–200, 259, 346  
biological process initiation, 591  
environmental pathogens in, 25  
in etiology, 509  
investigation of, 546  
substance-use comorbidity, 320  
underuse of, 5  
gene expression, 36–37  
gene-gene interaction, 199, 343, 546  
gene-nicotine dependence associations, 45–46  
Gene Ontology, 560  
gene-pathogen relations, 100–101  
general growth mixture modeling (GGMM),  
308–311  
generational changes, in smoking, 515  
Genes, Environment and Health Initiative, 531  
genetically informative designs, 527  
genetically modified mice, 444  
Genetic Analysis Workshops (GAW), 266  
genetic architecture, 510  
genetic association studies, 554, 556–559  
genetic drift, 150  
genetic epidemiology, 257  
extended, 250–252, 262–269, 279, 280  
genetic factors  
acute tolerance, 162–163  
conditioned place preference, 159  
craving, 441  
detection of, 493

- in experimentation, 31
- importance of, 29
- measured, 4
- in nicotine effects, 22, 39–40, 539
- quantitative models, 512
- in reinforcement, 418
- relative contribution of, 30
- selecting for, 96–97
- genetic heterogeneity, 32
  - adolescent smoking initiation, 196, 201–202, 233
  - in developmental trajectories, 190, 233–234
  - gender, 273–274, 276–278
  - phenotypes, 341
  - population, 217
    - estimating, 218, 221–222
  - receptor, 139
- genetic heterogeneity models, difference in fit
  - between homogeneity models and, 274–275
- Genetic Information Nondiscrimination Act (GINA), 49
- genetic latent class models. *See* latent class analysis
- genetic latent growth curve models. *See* latent growth curve models
- genetic mapping, 73–75. *See also* phenotypic research
  - analytic strategies, 96–103
  - construct validation, 75–78
  - core dependence dimension, 87–95
  - covariation among measures, 95–96
  - distal measures of dependence, 78–81, 86–87, 103–105
  - multidimensional measures, 81–86
  - person factors implications in, 97–98
- genetic modeling, 245–281
  - methodological and conceptual issues, 247–248
  - statistical framework for, 248–259
- genetic polymorphism effects, 553–554
- genetic substrata, associated with tolerance, 162
- genetic testing, 46, 50
- genetic variants
  - biological processes associated with, 109
  - causal, 546
  - disease association with, 550
  - evaluation context for dependence, 76
  - phenotypes with, 102, 106, 109, 111
  - pleiotropic associations of, 47–50
  - selection of, 101
  - value of, 110
- gene-to-phenotype influence, 78
- gene-transcription cascades, 169
- genome(s)
  - candidate genes in, 24
  - data, 561
  - studies of, 589
- genome markers, linkage analysis, 257–258
- genome scan, 266–267
- genome-wide association studies (GWAS), 25, 44, 269
  - event-related potentials, 428, 430
  - FTND, 42–45
  - genetic variant findings, 589
  - genotyping technologies used in, 258–259
  - potential of, 45–46
  - results from, 342
  - susceptibility loci identification, 407–408
- genotyping
  - effects of variables on, 553
  - mouse strains, 150
  - P450, 39
  - with phase interaction, 545
  - phenotypes and, 560–562
  - technologies, 257–259
- geographic information systems (GIS), 520–521
- GGMM (general growth mixture modeling), 308–311
- GINA (Genetic Information Nondiscrimination Act), 49
- GIS (geographic information systems), 520–521
- global use, 4, 21
- GluR (glutamate receptor), 157, 169
- glutamate receptor (GluR), 157, 169
- GMM (growth mixture modeling), 202, 218, 221, 308
- go/no-go task, 358–359, 451
- government policies, 7, 20
  - effect on adolescent smoking, 193
  - substance use, 294–295
- grant funding, 45–46
- graph connectivity, 571
- grouping variables, for growth curve modeling, 216–217, 232–233
- group membership, stability of, across statistical models, 229–230
- growth curve, nonlinear, 255
- growth curve mixture modeling, 215–217, 248–249
- growth mixture modeling (GMM), 202, 218, 221, 308

growth process, random effects for, 220–221  
gum. *See* nicotine gum  
gustatory reaction to tobacco, 75, 84  
GWAS. *See* genome-wide association studies

## H

habituation, *versus* addiction, 24  
half-life of nicotine, 147  
haplotypes  
    dependence and, 100  
    disease association with, 544  
    double variant, 545  
hard-core smokers, characteristics of, 35–36  
Hardy-Weinberg equilibrium, 545  
head rush, 374  
health care access, disparities in, 47  
health effects of smoking, statistics on, 4, 21  
heart rate, 351, 354, 440, 447  
Heaviness of Smoking Index (HSI)  
    components of, 37, 413  
    predictive value of, 80, 89–90  
    scales in, 79  
    zero-order correlations in, 80  
heavy smokers, 93  
    delay discounting, 350  
    diagnostic variance in, 89  
    genetic factors in, 29  
    *versus* light smokers, 43, 81, 83–84, 90, 94  
    substance use and, 296–297, 315  
Heavy Smoking Index, 265  
hedonic impact of nicotine, 158, 372, 424  
heritability  
    adolescent nicotine dependence, 86, 342  
    antisocial scores, 33  
    anxiety, 445  
    delay aversion, 351  
    dependence, 37–38, 86, 433–434  
    diagnostic criteria, 30, 38  
    endophenotypes, 107  
    estimates, 29–30  
    event-related potential, 428, 429–430  
    factors in, 28–29  
    gender differences in, 279, 281  
    impulsivity, 451  
    neuroticism, 101  
    nicotine metabolism/clearance, 38–40  
    P450 genotype, 39  
    prepulse inhibition startle response, 431–432  
    response inhibition, 359  
    resting EEG, 426  
    smoking cessation, 406  
    smoking heaviness, 90  
    withdrawal symptoms, 30  
    working memory, 437  
heterogeneity. *See* genetic heterogeneity  
hierarchical modeling  
    estimation for, 575–576  
    with ontologies, 551–552  
    with prior knowledge, 570  
    for statistical modeling, 117  
    stochastic variable selection and, 547–549, 572, 573  
    weighting in, 570  
high-affinity nAChRs, 156, 159  
higher-order joint actions, 546  
high genetic proneness, 96  
hippocampal activity, 141, 142  
HISTONE proteins, 36  
home smoking bans, 99  
homogeneity models, 271  
    difference in fit between heterogeneity models and, 274–275  
homogeneous population  
    assumption of, 248  
    for growth curve modeling, 216–217  
Hooked on Nicotine Checklist, 26  
Horn-Russell Scale, 34  
hostility, 357, 362, 377  
Household Adult Questionnaire, 501  
HSI. *See* Heaviness of Smoking Index  
*HTR5A* gene, 42  
*5-HTT* gene, 33, 269, 441  
*5-HTTLPR* gene, 101, 199, 409  
    in adolescent girls, 343  
    in affective response, 112, 446  
    coding for, 408  
human clinical research  
    affective regulation, 444–445  
    electrophysiological measures, 426  
    event-related potential, 428  
    impulsivity, 450  
    reinforcement, 413–414  
HumanCyc database, 556  
human genome, similarity with mice genome, 134  
3-hydroxycotinine, 406  
hyperactivity, 354, 356  
hypertensive rats, 449

**I**

IBD (identical by descent), 257, 258  
ICD-10 criteria, 80–81, 192  
ICR mice, 159, 162, 163  
ICSS (intracranial self-stimulation), 372–373, 421  
identical by descent (IBD), 257, 258  
illicit substances. *See* substance use; *specific drug*  
impulse control, 449–452  
impulsivity, 348  
    in attention deficit hyperactivity disorder, 354, 356  
    clinical research, 433, 450–451  
    cognitive control and, 112, 378–379  
    delay discounting with, 349, 350, 449, 450–452  
    heritability, 451  
    measurement of, 450–451  
    neural incentive system association, 346  
    preclinical research, 449–450  
    in response inhibition, 359  
incubation effect of initial exposure, 371  
independent pathway model, 253  
Indiana University Smoking Survey, 223–233  
    data analysis, 225–226  
    discussion, 231–233  
    measures, 224–225  
    procedures, 224  
    results, 226–231  
individual pathways to mature state, 105, 106  
“infectious disease” model, 522  
inflammation masking, 145  
inhibitors. *See* antagonists  
inhibitory interneurons, 141, 142  
initial exposure response measures, 339  
initial sensitivity, 27, 363–364  
    future research directions, 380–381  
    innate sensitivity, 364–368  
    other responses, 373–380  
    reinforcement, 368–372  
    rewards, 372–373  
innate sensitivity, 364–368  
instrumental learning, 158  
insurance companies, genetic testing and, 49  
integrative model of nicotine dependence, 22, 23  
integrative theory of triadic influence, 293  
intercept, distribution of, 219, 306, 310  
intercept models, initiation-based, 322  
intermediate phenotypes, 341, 342. *See also* endophenotypes

*International Statistical Classification of Diseases and Related Health Problems, Tenth Revision* (ICD-10) criteria, 80–81, 192  
interpersonal dynamics, 517, 522–525  
interval level, 247  
intracranial self-stimulation (ICSS), 372–373, 421  
intrathecal administration, 162  
intrauterine events, 100  
intravenous administration, 146  
    humans, 415  
    rodents, 152–157, 410–412  
in utero nicotine exposure, 357  
inveterate smokers, 87  
ion-channel receptors, 143  
Iowa gambling task, 350  
IRT. *See* item response theory  
item difficulty, 270  
item response theory (IRT), 254  
    empirical example of, 269–279  
    *versus* sum score approach, 278

## J

Jarvik, Murray, 24  
journal publishing requirements, 45–46

## K

kinship model, twin studies extended to, 250–252, 262, 279, 280  
*k*-means clustering, 217  
knockout mice, 159, 160, 161, 166, 422  
Kraepelinian approach to diagnosis, 291  
Kyoto Encyclopedia of Genes and Genomes, 561

## L

laboratory-based measurements  
    attention deficit hyperactivity disorder, 356  
    consumption, 418  
    endophenotypes, 349–351, 353–354, 357–362  
late emergent symptoms, 90–91  
latency to first puff, 415  
latent class analysis, 80, 92, 98, 248–249, 256–257, 266, 280  
latent class growth analysis (LCGA), 218, 226–229, 248–249  
latent growth curve (LGC) models, 254–256, 266, 280

association data integrated into, 259  
 substance-use comorbidity, 306, 308, 321  
 latent phenotype model, 252, 270  
 latent profile modeling, 90–91  
 latent trait, 270  
 latent variables, 77, 248, 256  
     substance-use comorbidity, 306, 321, 324  
 LCGA (latent class growth analysis), 218, 226–229,  
     248–249  
 LD (linkage disequilibrium), 543, 545–546  
 learning associations, 349  
 learning differences, 158, 169  
 letter cancellation task, 433  
 level, in latent growth curve models, 254  
 Lewis rats, 411, 423  
 LGC. *See* latent growth curve models  
 liability models, of smoking behavior stages, 264,  
     280  
 lifetime regular smoking, definition of, 29  
 ligand-activated ion channels, 143  
 light smoking. *See* chippers  
 likelihood-based approaches, in cluster analysis,  
     217–218  
 likelihood ratio tests, 310, 312–313  
 linear growth, assumption of, 255  
 linear model, 116  
 linear regression, in structural equation modeling,  
     248  
 linear relations, 25  
 linkage analysis, 257–258, 266–268  
     candidate gene studies, 32, 40–42, 267–268,  
         280–281, 589  
     environmental factors, 258  
     genome markers, 24, 257–258  
     nicotine-dependence indices, 40–42  
 linkage disequilibrium (LD), 543, 545–546  
 Lister rats, 432–433, 444  
 lithium-chloride conditioned place aversion, 161  
 liver cytochrome P-450 enzyme CYP2A6. *See*  
     CYP2A6 gene  
 location of smoking, 527  
 loci segregation, 40  
 locomotor activity, 369, 422  
 logarithm of odds (LOD) score, 40–42, 258, 267  
 logistic regression curves, 91  
 Long-Evans rats, 411  
 longitudinal data  
     growth curve modeling of, 218–219, 234  
     on substance-use comorbidity, 291, 320  
 Loss of Control subscale, 90–91

low genetic proneness, 96  
*LPAAT-delta* gene, 42  
 lung, nicotine concentration in, 145  
 lung cancer, predisposition to, 44, 50

## M

macrocontextual factors, 509, 514  
     as moderators, 515–516  
 macroenvironment proximal indicators, 521–522  
 macular degeneration, age-related, 46  
 magnetic resonance imaging (MRI), 351, 439,  
     456–457  
 maintenance of dependence, 589  
 male smokers. *See also* gender differences  
     adolescent, 342, 348  
     blood pressure changes in, 449  
     nicotine-dependence factors, 37, 38, 99  
     *OPRM1* gene in, 419  
     statistics on, 21  
     twin studies, 515, 517  
 manifestations of dependence, 75  
 Mannheim Study of Risk Children, 199  
 MAO. *See* monoamine oxidase  
*MAP3K4* gene, 42  
 marginalization  
     of smoking, 4, 21  
     of social groups, 47  
 marginal nonnormality, 221  
 marijuana use  
     early pleasurable, 374  
     tobacco use concurrently with, trajectories of,  
         296–298  
     modeling, 304–305  
     trajectories of, 295–296  
 marketing direct-to-consumer, 50  
 Markov chain Monte Carlo (MCMC) methods, 542  
 masking etiology, 82  
 masking of causal factors, 74  
 maternal care, 36  
 mature subphenotypes, 82, 110–111  
 maximum acute tolerance, 162, 165  
 maximum price assessment, 418  
 McGill University Study on the National History  
     of Nicotine Dependence, 199  
 MCMC (Markov chain Monte Carlo) methods, 542  
 mean (centroid)  
     cluster analysis, 217  
     growth curve modeling, 255  
     structural equation modeling, 248–249



- measured genetic factors, 4
- measurement invariance, 247, 255, 276, 280
- mecamylamine, 159
- mediation
  - of conditioned place preference, 161
  - by endophenotypes, 107–108
  - of nicotine, 160
- memantine, 156, 171
- memory, 379, 434–438
- Mendelian randomization, 118
- mesolimbic dopaminergic system, 410, 411
- metabolic tolerance, 97
- metabotropic glutamate receptor 5 (mGluR5), 156
- methodological issues
  - assessment precision, 520
  - behavioral modeling, 247–248
  - family-based studies, 521
  - future research directions, 383–384, 594
  - genetic modeling, 247–248
  - innate sensitivity research, 366–368
  - real-time interaction, 524–525
  - research limitations, 366–368, 376, 383–384
  - substance-use comorbidity, 321–323
- methyllycaconitine citrate (MLA), 159
- mGluR5 (metabotropic glutamate receptor 5), 156
- mice. *See* mouse models
- microchip analysis, 170, 258–259
- microcontextual factors, 509, 514
  - coding of real-time interaction, 524
  - as moderators, 516–518
- microsatellites, 543, 551
- microsocial context, quantifying, 522–525
- midbrain dopamine circuits, 348, 349
- Mid-South Tobacco Family (MSTF), 267
- migration levels, 516
- Minnesota Nicotine Withdrawal Scale, 446
- Minnesota Twin Family Study (MTFS), 196, 260, 360
- mirror tracing, 448
- misleading claims, 50
- misspecification, model, 222
- mixed models, developmental trajectories, 215
- MLA (methyllycaconitine citrate), 159
- model(s)
  - clarification of, 559
  - searching, 551
  - selection of, 547
  - with stochastic variable selection, 549–550, 551–552
- model fit, evaluation of, 310, 312–313
- modeling. *See also specific types of modeling*
  - phenotype (*See* phenotype modeling)
  - with prior knowledge, 570
  - selection algorithm, 577–578
  - uncertainties in, 546, 547, 549
- model misspecification, 222
- moderation of relationships, 248
- modified pairwise interaction, 545
- modulation
  - dopamine receptors, 156
  - nicotine rewards, 160
- molecular genetic studies, 266
  - of adolescent smoking, 198–199
  - analytic framework for, 257–259
- Monitoring the Future project, 299, 304
- monoamine oxidase (MAO)
  - in anxiety disorders, 353
  - neuroticism and, 351
  - in neurotransmitter breakdown, 540
- monoamine oxidase (*MAOA/MAOB*) gene, 269
- mood effects, 373–376, 380–381
  - in ATBC analysis, 497
  - measures of, 372
  - of nicotine, 366
- Mood Form of Diener and Emmons, 374–375, 446
- morning smoking. *See* time to first cigarette
- morphine, as nicotine substitute, 153
- Morris water maze, 435
- mortality statistics, 4, 21
- motivational mechanisms, 84, 403, 408
  - reinforcement, 410–420
  - rewards, 420–424
- Mouse Genome Informatics database, 150
- mouse models, 134–135, 418
  - adolescent exposure, 194–195
  - behavioral changes, 151–157
  - future research directions, 168–172
  - nicotine administration, 145–149
  - nicotine dependence, 149–151
  - nicotinic receptors, 135
    - customizing, 141–143
    - functional diversity of, 136–141
    - molecular biology of, 136
    - nicotine as agonist/antagonist, 143–144
    - upregulation, 144–145
- reward, 157–161
- startle inconsistency, 444
- strains, 134 (*See also* strain-specific differences; *specific strain*)
  - research options with, 592
  - selection of, 150–151

tolerance, 162–168  
mouse-rat differences, 150  
movies, smoking in, 7, 20, 523  
MRI (magnetic resonance imaging), 351, 439, 456–457  
MSTF (Mid-South Tobacco Family), 267  
MTFS (Minnesota Twin Family Study), 196, 260, 360  
multidimensional measures, of nicotine dependence, 81–86  
multifinality, 191  
multilevel analysis, difficulties of, 248  
multiple trajectories, developmental, 191–202, 232–234  
multivariate analysis  
developmental trajectories, 215  
latent growth curve models, 255  
substance-use comorbidity, 315–316, 321  
twin studies, 262–263  
multivariate factor model, 252–253  
multivariate normal distribution  
in growth curve models, 255  
in structural equation modeling, 249  
multivariate normality, within-class, 221, 233  
mu opioid knockout mice, 422  
mu opioid receptors  
in conditioned place preference, 160  
in nicotine replacement therapy, 407  
reward mediation, 419  
in tolerance, 167, 168  
muscarinic acetylcholinergic systems, 166, 168  
muscarinic receptors  
in aging, 170  
blockading, 140–141  
metabotropic, 136  
muscle tension, 377–378  
mutations  
rate predictions, 554  
for tolerance, 166  
Muthén, Bengt, 218

## N

nAChRs. *See* nicotinic acetylcholine receptors  
Nagin, Daniel, 218  
naloxone, 160  
nasal spray. *See* nicotine nasal spray  
National Cancer Institute, 498  
National Center for Biomedical Ontology (NCBO), 561

National Comorbidity Study, 445  
National Health and Nutrition Examination Survey (NHANES III), 487, 494, 500–503  
National Institute of Mental Health Diagnostic Interview Schedule (DIS), 81, 501  
National Institute on Drug Abuse Genetics Consortium, 43  
National Institutes of Health, 531  
National Longitudinal Study of Adolescent Health (Add Health), 197–198, 261, 517  
National Survey on Drug Use and Health, 93  
N-back task, 436  
NCBO (National Center for Biomedical Ontology), 561  
NDSS. *See* Nicotine Dependence Syndrome Scale  
NEAD (Nonshared Environment in Adolescent Development) Project, 509, 524–525  
Netherlands Twin Register, 262  
Netherlands Twin Study of Anxious Depression (NETSAD), 266–267  
network models, 554, 556–558  
neural analysis, 349–351, 353–354, 357–362  
neural incentive system, 346  
neural networks modeling, 340–341  
neural substrata, associated with tolerance, 162  
neural systems, candidate, 343–346  
neurexin 1 (*NRXN1*) gene, 43  
neurobiological analysis, 348  
neurobiological dependence pathways, 43  
neurobiological systems, 344  
neuroendocrine response, to stress, 354  
neuroimaging, 360, 379–380  
neuropeptide systems, 159, 160  
neuroprotection, 170  
neuroticism, 101, 351–353  
neurotransmitter systems  
in chronic tolerance, 166–167  
in conditioned place preference, 159–161  
New England Family Study, 526  
NHANES III (National Health and Nutrition Examination Survey), 487, 494, 500–503  
nicotine  
administration of (*See* administration)  
age-related response differences, 194–195  
as agonist/antagonist, 143–144  
anti-inflammatory effects of, 148–149  
anxiogenic effects of, 168  
behavioral changes from, 151–157  
bioavailability of, 7, 20  
enforcement timing, 369  
in free-base form, 148

- frequency of use, 4, 21, 369–371, 413, 420
- hedonic impact of, 158, 372, 424
- neuronal activity induced by, 43
- physical changes from, 162
- pre-exposure risk (*See* pre-exposure risk)
- pretreatment, 420
- reinforcement (*See* reinforcement)
- rewards and (*See* reward)
- tolerance (*See* tolerance)
- nicotine aerosols, 147
- nicotine-binding sites, 135, 144
- nicotine choice, 369
- nicotine choice procedure, 416, 419–420
- nicotine cigarette choice paradigm, 419
- nicotine clearance
  - dispositional tolerance, 162, 164, 168
  - genetic factors in, 39–40, 539
- nicotine dependence, 20–22, 149–151
  - in adolescence (*See* adolescent nicotine dependence)
  - concurrent with substance use (*See* substance-use comorbidity)
  - construct refinement, 25–27
  - craving associated with (*See* craving)
  - crosscutting issues, 595–596
  - developmental pathways in, 589
  - distal measures of (*See* distal measures)
  - endophenotypes in, 409, 453
  - epidemiological concepts, 31–37
  - future research directions, 455–457
  - heritability of (*See* genetic factors; heritability)
  - historical perspective of, 22–25
  - inference of, 77
  - mouse models of (*See* mouse models)
  - phenotype (*See* phenotype(s))
  - progression research, 592
  - psychiatric disorders correlated with, 98
  - risk with, 375–376
  - smoking compared with, 87
  - versus* tobacco dependence, 75
  - treatment of (*See* smoking cessation)
  - understanding of, 588–596
- nicotine-dependence measures, 26, 28, 37–45, 73, 78, 79
  - adolescents, 192, 230–231, 264
  - example of, 225
  - invariance, 276, 280
- Nicotine Dependence Syndrome Scale (NDSS), 82–84
  - abbreviated, 500
  - as assessment tool, 405
  - subscales of, 90
  - in TUS-CPS analysis, 498
- nicotine deprivation memory deficits, 436
- nicotine gum
  - as consumption assessment, 415
  - effect on EEG activity, 426
  - in memory effects, 435
  - versus* placebo, 370
- nicotine metabolism
  - association analysis of, 268
  - catabolism of, 149
  - CPD variation and, 34
  - CYP2A6* gene in, 39–40, 149, 342, 406, 418
  - in dependence risk, 22, 342
  - in ontology example, 562–569
  - pathway, 556
- nicotine nasal spray
  - aversion with *DRD4*\*7-repeat allele, 376
  - as consumption assessment, 415
  - in current smokers, 374
  - memory effects of, 435–436
  - versus* placebo, 369–370
  - pleasurable responses to, 373
- nicotine patches
  - effect on EEG activity, 426
  - in memory effects, 435
  - on nonsmoking adults, 377
- Nicotine Pharmacokinetics Ontology (NPKO), 539, 561, 571
- nicotine replacement therapy (NRT), 406. *See also* smoking cessation
- nicotinic acetylcholine receptors (nAChRs), 134
  - association analysis, 268
  - as attention factor, 362
  - beta2-subunit (*CHRNA2*), 268
  - binding to, 588
  - blockading, 140–141, 143–144
  - chromosomal regions, 43, 44, 50
  - in chronic tolerance, 166–167
  - coding for, 408
  - customizing, 141–143
  - desensitization of, 146
  - fast-ionotropic, 136
  - functional diversity of, 136–141
  - high-affinity, 157
  - illustration, 409
  - as impulsivity mediator, 449
  - inactivation of, 143
  - inferences from, 152–153

initial sensitivity response with, 376–377  
 molecular biology of, 136  
 in nicotine binding role, 406–407  
 in nicotine dependence, 411  
 structure of, 136–138, 138  
 tissue-specific responses, 71, 133  
 in tolerance, 168  
 upregulation of, 135, 144–145, 588  
 nicotinic receptor subunits, 136–140  
   *CHRNA5/CHRNA3* genes, 342  
   composition of, 139  
   functional variants of, 140  
   limiting expression, 141–143  
   structure of, 136, 137  
 nimodipine, 162, 167, 168  
 nitric oxide, in conditioned place preference, 161  
 7-nitroindazole, 161  
*N*-methyl-D-aspartic acid (NMDA) glutamate  
   receptors, 156  
 NMRI outbred mice, 154, 412, 430  
 nomological net, 76  
 noncoding DNA sequences, 554  
 nonlinear growth curves, 255  
 nonnicotinic systems, 166  
 nonnormality, 221  
 Nonshared Environment in Adolescent  
   Development (NEAD) Project, 509, 524–525  
 nonsmokers (abstainers)  
   characteristics of, 22  
   inclusion *versus* exclusion of, 222–223, 235, 323  
   nicotine patches, 377  
 Northern California Twin Registry, 539, 540  
 novelty seeking, 155–156, 199, 348–349  
   by adolescents, 27  
   nasal spray use in, 373  
   neural incentive system association with, 346  
   in substance use, 292–293, 348–349  
 NPKO (Nicotine Pharmacokinetics Ontology), 539  
 NRT (nicotine replacement therapy), 406. *See*  
   *also* smoking cessation  
*NRXN1* gene, 43  
*NRXN3* gene, 43  
 nutritional cancer prevention, 497

## O

OBO (Open Biomedical Ontologies), 560  
 OBO-Edit, 561  
 OBO Foundry project, 560  
 OBO Relation Ontology, 561  
 observed variables, in structural equation  
   modeling, 248  
 occurrent, 541, 559  
 oddball stimulus, 429  
 olanzapine, 456  
 ontologies, 539–541  
   definition of, 541, 559  
   development process, 560–562  
   discussion of, 570–574  
   methods, 541–543, 550–562  
   nicotine metabolism, 562–569  
   statistical approaches, 543–550  
 Ontology Web Language (OWL), 561  
 Open Biomedical Ontologies (OBO), 560  
 Open Source software, 561  
*OPRM1* gene, 409  
   coding for, 408  
   gender differences in, 419  
   in nicotine replacement therapy, 407  
   in smokers, 423  
   support interval proximity, 42  
 oral administration, 147–149, 172, 412–413  
 oral mucosa exposure to nicotine, 148  
 ordinal data, analytical framework for, 247, 255  
 osmotic minipump, 147  
 outcomes of dependence, 75  
 outliers, controlling for, 281  
 OWL (Ontology Web Language), 561  
 oxotremorine, 166–167

## P

P3 amplitude, 294  
 panic disorder, 353  
 PANTHER Pathways databases, 556, 561  
 parent(s)  
   educational level of, 519  
   twin studies extended to, 251, 262, 279, 280  
 parental monitoring, as smoking counterforce, 22  
 parental smoking  
   adolescent smoking and, 196–197, 200, 262,  
     346  
   behavioral modeling of, 246  
   nasal spray non-response, 377  
   smoke-free home with, 523  
   socialization effects from, 517  
 parenting behavior, 342  
 partition variation, 270, 280  
 passive avoidance, 435  
 patch. *See* nicotine patches

- path diagrams, in structural equation modeling, 248–250
- pathogen modeling, 100–101, 103
- PATO (Phenotype and Trait Ontology), 560, 571
- Pavlovian learning, 158
- PBT (problem behavior theory), 198, 292–294, 322
- PDAs (personal digital assistants), 525–527
- peers
  - as influence, 100, 346, 517
  - as smoking predictors, 523
- peer smoking, effect on adolescent smoking, 197
- penetrance, incomplete, 31
- peripheral nervous system (PNS), 344, 348
- P50 ERP, 427–428
- persistence
  - drug use, 417–418
  - negative affect with, 447
  - smoking (*See* smoking persistence)
- personal digital assistants (PDAs), 525–527
- personalized health care, 49–50
- person factors, 97–99
- PET (positron emission tomography), 439
- P300 event-related potential (ERP), 359–360, 429–430
- P450 genotype, 39
- pharmacokinetics, 37, 39, 118, 149, 542, 590
- pharmacokinetics ontology, 539, 561, 571
- phenotype(s)
  - assays, 83
  - association of, 33
  - behavioral, 171, 492
  - characteristics of, 109
  - components, 341
  - developmental progression of, 74
  - effects of variables on, 553
    - as endophenotypes, 5, 33–34
  - environmental, 34–36
  - framework, 27–31
  - genetic mapping, 96–103
  - genetic variants to, 102
  - genotyping and, 560–562
  - heterogeneity, 341
  - intermediate, 341, 342 (*See also* endophenotypes)
  - of mouse strains, 150
  - pathways, 23, 590
    - as points in smoking trajectory, 490
  - research pitfalls, 489
  - stages, 109
    - substance-use disorders, 316–318, 321, 324
- Phenotype and Trait Ontology (PATO), 560, 571
- phenotype modeling, 487, 488–492
  - examples, 493–503
  - methods, 492–494
  - summary, 503–505
- phenotypic assortative mating, 251
- phenotypic cultural transmission, 251
- phenotypic research, 105–106
  - analytic strategies, 113–118
  - causal paths, 107–110, 111–113
  - caveats, 110–111
  - summary, 118–119
- phenylthiocarbamide (PTC) haplotype, 84
- phosphatase and tensin homolog (*PTEN*) gene, 269
- physical aggression, 357
- physical responses to nicotine, 169
- symptoms, 26
  - in tolerance, 162–163, 164–165
- physiological measures of response reward, 351
- physiological startle. *See* startle response
- physiology
  - of affect, 377–378
  - of behavioral traits, 344, 345, 346
- Pittsburgh Youth Study, 214
- placebo, *versus* nicotine, 369–370, 420
- placenta, 145
- plasma nicotine levels
  - with IV injection, 146
  - tissue nicotine levels compared with, 145
  - in tolerance studies, 163–164
- pleiotropy, 32, 47–50
- PNS (peripheral nervous system), 344, 348
- point mutation, 444
- policy interventions, 34–35
- polymorphisms, 553–554. *See also* single nucleotide polymorphism
  - associations with, 116
  - different priors for, 571
  - in dopamine reward pathway, 342
  - emphasis within genes, 547
  - investigation of, 546–547
  - numerous, challenges of, 546–547
  - perturbations from, 554
  - trait variation effect of, 549
- PolyPhen (polymorphism phenotyping), 554
- polysubstance use, 43, 296
- population
  - frequency of genetic factors in, 86

- for growth curve modeling, 216–217, 234
- homogeneous, assumption of, 248
- in latent class analysis, 256
- response distribution in, 370, 373, 376–380
- in structural equation modeling, 248–249
- population heterogeneity, 217
  - estimating, 218, 221–222
- Positive and Negative Affect Schedule, 374–375, 446
- positron emission tomography (PET), 439
- postural hypotension, 449
- PPI (prepulse inhibition), 378, 403, 430–432
- preclinical research. *See also* animal studies; mouse models
  - affective regulation, 443–444
  - electrophysiological measures, 425–426
  - event-related potential, 427–428
  - impulsivity, 449–450
  - reinforcement, 410
  - rewards, 421–423
- precursors, class III phenotypes as, 30
- predictive validity
  - of genetic testing, 50
  - of primary motives scales, 90–91
- pre-exposure risk, 339, 340. *See also* smoking initiation and progression risk
  - endophenotypes, 340–347
  - future research directions, 381–385
  - initial sensitivity endophenotypes (*See* initial sensitivity)
- preproenkephalin knockout mice, 160, 422
- prepulse inhibition (PPI), 378, 403, 430–432
- price-demand curve, 417
- pricing, of tobacco products, 20, 21, 520
- primary motive scales, as predictors, 90–93
- prior covariate specification, 570–571
- prior knowledge, ontologies and, 553–562
- PR (progressive ratio) measures, 417–418, 420, 454
- probability discounting, 450–451
- problem behavior theory (PBT), 198, 292–294, 322
- problem use, 196, 198, 263, 294
- Profile of Mood States, 374–375, 446
- programmed lapse procedure, 454
- progression to smoking, 491
- progressive ratio (PR) measures, 417–418, 420, 454
- Project on Human Development in Chicago Neighborhoods, 520
- “proof of concept” analyses, 493, 494
- protective factors
  - in adolescents, 343
  - versus* vulnerabilities, 87, 114
- protein sequence data, 561
- protobacco advertising, 7, 20, 30, 348
- prototypes, of nicotine-dependence research, 27–28
- proximal indicators, 509
- proximal influence, 519–522
- proximal measures
  - versus* distal measures, 516
  - of social context, 518–527
- psychiatric comorbidity, 81, 98–99
  - with *DSM-IV* dependence, 26, 81
  - empirical examples of, 496, 502
  - resolving, 115
- psychiatric genetics research, 25, 27
- psychoactive alkaloid, 22
- psychological traits
  - approach-related risk, 346–349
  - avoidance-related risk, 351
  - control-related risk, 354–357
- psychometric common factor model, 252
- psychopathology
  - developmental, 191–202, 292–293
  - indices, 32
  - physiological basis of, 344
  - substance use and, 292–293, 304
- psychophysiological responses
  - to acute stressors, 448–449
  - craving, 440
- psychosocial factors
  - in adolescent nicotine dependence, 195
  - in adolescent smoking initiation, 200, 202, 211
- PTC (phenylthiocarbamide) haplotype, 84
- PTEN* gene, 269
- public health messages, 47
- public health outcomes, 492
- public settings, smoking in, 4, 20–21
- putative endophenotypes, 341

## Q

- QSU (Questionnaire on Smoking Urges), 439
- QTL (quantitative trait locus), 258
- quantitative genetic models, 512, 522
- quantitative trait locus (QTL), 258
- quantity smoked measures, 28
- Questionnaire on Smoking Urges (QSU), 439



## R

- racial background, 439. *See also* African Americans
- racism, associated with genetic information, 47
- random effects, 216
  - within-class, 220–221, 233
- rapid-decision context, 358
- Rapid Visual Information Processing (RVIP) task, 433
- rat models
  - adolescent nicotine exposure in, 194–195
  - alcohol/nicotine correlation, 369
  - conditioned place preference in, 423
  - self-administration of electrical stimulation, 421
  - sensitivity in, 364
  - strains, 411 (*See also specific strain*)
- Reactome, 561
- real-time contexts, 526
- real-time interaction, 524–525
- recall, 379
- receptor heterogeneity, 139
- recovery, from acute tolerance, 162
- regression models, 248, 542, 543, 550–551, 572
- regular smoking
  - definition of, 29
  - genetic factors in, 29–30
- reinforcement, 151–152, 155–156, 366–372, 380
  - alternative, 417
  - in cognitive control, 349
  - enhancing, 413
  - genetic influences in, 418
  - in initial sensitivity, 368–372
  - measurement of, 414–420
  - motivational mechanisms, 410–420
  - secondary, 411, 413
- relapse
  - environmental influences in, 95
  - physiological basis of, 588
  - predictors of, 77, 413, 434, 442, 444–445
  - time to first cigarette as predictor of, 80, 89
- relations between entities, 541, 559, 560
- relative measurements, 571
- reliability. *See also* test-retest reliability
  - of developmental trajectory research, 234
  - of nicotine-dependence measures, 26, 79–80
  - recall, 26–27
- reliability coefficients, 79
- religiosity, adolescent smoking and, 198
- repeated-measures data, developmental trajectories, 215, 222
- replication, of gene-nicotine dependence associations, 45–46
- research findings, communication of, issues in, 45–50
- research limitations
  - adolescent smoking, 367
  - adult smoking phenotype, 190
  - ATBC analysis, 497, 504
  - behavioral measures, 104
  - cue-induced craving, 456
  - data, 572
  - distal measures, 103
  - DSM, 103
  - extreme groups, 114–116
  - methodology, 366–368, 376, 383–384
  - NHANES III analysis, 503–505
  - nicotine-dependence measures, 103–104
  - Nicotine Dependence Syndrome Scale, 84
  - retrospective reporting, 367
  - self-report measures, 367
  - smoking cessation research, 454–455
  - statistics modeling, 570
  - tobacco dependence assessments, 95
  - TUS-CPS analysis, 500, 504
  - twin studies, 279–280
  - Wisconsin Inventory of Smoking Dependence Motives, 85–86
- residual familial factors (F), 258
- residual item variances, 270
- respiratory sinus arrhythmia (RSA), 361
- response inhibition, 358–359
- resting EEG activity, 425–427
- retail tobacco outlets, 520, 521
- retrospective reporting limitations, 367
- reward, 157–161
  - definition of, 349
  - for depressed smokers, 352
  - future research directions, 454–455
  - immediacy over magnitude, 348, 349
  - in initial sensitivity, 372–373
  - measurement of, 351, 366, 372–373, 380, 423–424
  - modulation of, 160
  - motivational mechanisms, 420–424
  - mu opioid receptor mediation, 419
  - preclinical studies, 157–161, 372–373, 421–423
  - signaling, 350
- reward and pleasure pathways, 22
- reward-discounting tasks, 350



- rimonabant, 160
- risk. *See also* pre-exposure risk; smoking  
    initiation and progression risk  
    age-specific, measurement of, 35  
    approach-related, 339, 346–349, 362  
    avoidance-related, 339, 351–354, 362  
    control-related, 339, 354–357, 362  
    cumulative, measurement of, 35  
    disease, 541, 550  
    with nicotine dependence, 375–376  
    nicotine metabolism and, 22, 342
- risk factors  
    adolescent smoking initiation, 211–212, 232, 350, 433  
    substance use, 306, 340
- risk-taking behavior, by adolescents, 195, 199
- RNA analysis, 142
- rodent models. *See also* mouse models; rat models;  
    *specific strain*  
    adolescent nicotine exposure, 194–195  
    adolescent sensitivity, 371  
    intravenous self-administration, 410  
    nicotine effects, 368, 435  
    strain-specific differences, 418
- RSA (respiratory sinus arrhythmia), 361
- \**RS578776* subunit gene, 43
- \**RS16969968* subunit gene, 43
- Russell, M.A.H., 24
- RVIP (Rapid Visual Information Processing) task, 433
- S**
- saccharin, 148, 154, 159
- S allele, 112, 113
- sample size/followup  
    in ATBC analysis, 497  
    for developmental trajectory research, 234  
    in TUS-CPS analysis, 498
- saturated model, 547
- SBML (Systems Biology Markup Language), 561
- schizophrenia, 428, 443
- secondary criteria of nicotine dependence, 76, 77
- secondary motives scales, 92–93
- secondary reinforcement, 411, 413
- secondhand smoke, 20, 246
- second-stage mixture model, 572
- self-administration  
    ad libitum (ad lib), 415–416, 419  
    genetic effects on, 146  
    intravenous, 146, 152–157, 410–412  
    oral, 412–413
- self-control, in children, 344
- self-insuring firms, 49
- self-report measures  
    of affect, 446–447  
    components of, 79  
    of craving, 439–440  
    ecological momentary assessment in, 525  
    limitations of, 367
- SEM. *See* structural equation modeling
- semi-Bayes approach, 549
- semistructured paradigms, 524
- Sensation Seeking Scale, 377
- sensitivity  
    of measurements, 341  
    modeling, 363–364, 364  
    periodic, 96
- sensitivity analysis, 570
- sensory measures, 403
- Sensory Questionnaire, 423
- sequential process model, of substance-use  
    comorbidity, 304
- serotonin  
    association analysis, 268–269  
    genetic variation in, 343  
    metabolism of, monoamine oxidase in, 353  
    regulation of, 540  
    smoking cessation and, 407
- SES. *See* socioeconomic status
- seven-point Likert scale, 372
- shared environment effects  
    in adolescent smoking, 197, 260–261, 264, 280  
    twin studies, 251, 280
- Shiffman-Jarvik Withdrawal Scale, 440
- Shiffman Nicotine Dependence Syndrome Scale.  
    *See* Nicotine Dependence Syndrome Scale
- sibling(s)  
    IBD configurations for, 257–258  
    smoking epochs of, 531  
    as smoking predictors, 523  
    socialization studies, 517  
    twin studies extended to, 251, 262, 279, 280
- Sibling Partners Study, 526
- sickle cell discrimination, 47
- side effects  
    from drinking-water administration, 172  
    from intravenous administration, 146
- SIFT (Sorting Intolerant From Tolerant)  
    procedure, 554

- simultaneous effect of genes, 32–33
- single-factor dependence, 79n
- single-factor structure, 81
- single-group growth curve model, 219
- single nucleotide polymorphism (SNP)
  - candidate gene variants, 25
  - disease association with, 544
  - genotyping, 43, 257–259
  - nonsynonymous coding, 554
  - as reflection of underlying effects, 546
  - relationships over candidate genes, 545
  - relevance of, 570
  - in whole-genome research, 4
- situational dependence, 27
- six-class solution, example of, 227–228
- skin conductance, 354, 377–378
- skin temperature, 440
- SLC6A3* gene, 198
- SLC6A4* gene, 32, 101, 112, 113
- sleep problems, in children, 357
- slope
  - distribution of, 219
  - factor loading and, 275–276
  - in latent growth curve models, 254, 306, 310
- 129S6 mice, 430
- SMOFAM (Smoking in Families Study), 267
- smoke-free laws, 520
- smoke-free settings, 20, 30, 523
- smokeless tobacco, 147
- smokers, characteristics of, 22
- smoking
  - bans on, 99
  - decline in, 20
  - developmental phenotypes (*See* adolescent developmental trajectories)
  - first experience with (*See* initial sensitivity)
  - frequency of, 369–371, 413, 420
  - during illness, 79, 80
  - nicotine dependence compared with, 87
  - quantitative genetic model and, 512
  - status, 28
  - transition levels, 42–43, 488–490
- smoking cessation
  - age-related changes in, 170
  - barriers to, 4–5, 46
  - CHRNA2* gene in, 407
  - commercial testing, 50
  - comparison groups changes, 493
  - data sets in, 493–494, 494
  - definition of, 492
  - delay discounting factor in, 350
  - demand for, 21
  - difficulties of, 97
  - drugs for, 160
  - failures in, 44–45
  - FTND predictions of, 80
  - future research directions, 454
  - gender differences in, 38
  - heritability in, 406
  - monoamine oxidase decrease during, 351
  - research limitations in, 454–455
  - serotonin pathway and, 407
  - subthreshold pretreatments, 162
  - success predictors, 81, 89, 413, 434
  - symptoms of, 447
  - tailored, 48
- smoking index variable, 30
- Smoking in Families Study (SMOFAM), 267
- smoking initiation, 31, 42
  - adolescent (*See* adolescent smoking initiation)
  - assessment of, age effects in, 279, 322
  - definition of, 29
  - gender differences in, 267
  - linkage analysis of, 267
  - versus* persistence, 406
- smoking initiation and progression risk, 346, 491.
  - See also* pre-exposure risk
  - approach-related, 346–349
  - avoidance-related, 351–353
  - control-related, 354–357
  - endophenotypic measures, 349–351, 353–354, 357–362
  - future research directions, 362–363
- smoking level
  - measurement of, example of, 224–225
  - substance-use comorbidity and, 308, 322
- smoking pattern, factor analysis of, 265
- smoking persistence, 29, 30, 406
  - definition of, 29
  - early-onset smokers' risk for, 200, 212–213, 230–231
- smoking topography devices, 415, 418
- SNP. *See* single nucleotide polymorphism
- social context, 509, 510–511
  - adolescent smoking, 193, 198
  - behavioral genetics in, 514–518
  - future research directions, 527–532
  - proximal measures of, 518–527
  - rationale for, 511–514
  - substance use, 294–295

- social development model, 293
- socioeconomic status (SES)
  - adolescent smoking and, 232
  - distal to proximal influence, 519–522
  - of hard-core smokers, 35
  - nuanced approaches to, 519–520
- socioregional influences, 515–516
- “softening” of smoking, 193
- software
  - association analysis, 259
  - item response theory, 270
  - linkage analysis, 258
  - ontology, 561, 562
- Sorting Intolerant From Tolerant (SIFT)
  - procedure, 554
- SourceForge Web site, 562
- species-specific responses, 134
- specific-factor models, of substance use, 306–307, 315–316
- speed congenics, 151
- spinal cord minipumps, 163
- spouses, socialization effects from, 517
- Sprague-Dawley rats, 411, 431, 432–433
- stage models, developmental trajectories, 233
- startle-probe measures, 113
- startle response
  - as affective response, 447–448
  - in humans, 377
  - increases in, 444
  - prepulse inhibition of, 378, 403, 430–432
  - test-retest reliability of, 446–447
- state laws, against genetic discrimination, 49
- static clustering, *versus* dynamic clustering, 219–220, 232
- statistics
  - approaches to, 543–550
  - combining genetic studies with, 248
  - developmental phenotypes, 214–231
  - modeling, 248–259, 570
  - ontological knowledge in, 572
- ST/b inbred mice, 154, 155, 412–413
- stem cells, 151
- Sternberg Memory Task, 379, 436–437
- stigma, nicotine dependence as, 46–47
- stochastic variable selection
  - hierarchical modeling and, 547–548, 572, 573
  - model selection with, 549–550, 551–552
- stop-go task, 359
- strain-specific differences
  - aging, 170
  - conditioned place preference, 158
  - DNA markers, 151
  - five choice serial reaction time task, 432–433
  - genetic, 150
  - mouse models, 153–154, 164–165, 412–413
  - nicotine effects of, 135, 161, 169, 418
  - nicotinic acetylcholine receptors, 172
  - rat models, 411
  - tolerance, 164
  - unraveling of, 142
- strain surveys, 422
- stress, 351
  - as influence, 100
  - influences on, 36
  - neuroendocrine response to, 354
  - response mediators, 143
- Stroop interference task, 379, 440, 451, 452
- Stroop paradigm, 113
- structural equation modeling (SEM), 248–249
  - combined with latent class models, 256
  - of developmental trajectories, 215
  - linkage analysis and, 257–258
  - for twin data, 249–257
- study participants, selection of, 490–492
- subcutaneous administration, 147, 172
- subpopulation
  - in latent class analysis, 256
  - in structural equation modeling, 248–249
- subPSEC (substitution position-specific evolutionary conservation), 554
- substance use. *See also specific substance*
  - adolescent, shared environmental influences in, 261
  - age-dependent vulnerability to, 195, 198, 200–201, 212, 233–234, 292–293
  - attention deficit hyperactivity disorder in, 356
  - common-vulnerability model, 291–294, 306, 315–316
  - dopamine in, 410, 588
  - early pleasurable use, 374
  - environmental factors influencing, 294–295
  - interpersonal dynamics in, 522
  - Iowa gambling task as predictor, 350
  - modeling of, 43
  - nasal spray non-response, 377
  - novelty seeking in, 292–293, 348–349
  - prevention of, research approaches, 294–295
  - risk factors for, 340
  - specific-factor models, 306–307, 315–316

- substance-use comorbidity, 289–325
    - association between smoking trajectories and, 296–298
    - empirical examples of, 307–323, 496, 502
    - future research directions, 323–324
    - gender differences in, 304
    - gene-environment interactions in, 320
    - importance of studying, 290–292
    - informative phenotypes for, 316–318, 321, 324
    - literature review, 295–296
    - mechanisms underlying, 291–292
    - methodological issues, 321–323
    - modeling, 298–307
    - risk factors, 306
    - two-stage models of, 323
  - substance-use disorders
    - common *versus* specific liability to, 292–295
    - diagnosis of, 291
    - shared genetic risk for, 294
  - substitution position-specific evolutionary conservation (subPSEC), 554
  - subthreshold depression, 352
  - sucrose, 154–155, 412
  - support interval, 40, 42
  - Surgeon General's Report (1964), 24
  - Surgeon General's Report (1979), 24
  - susceptibility loci mapping, 24. *See also* candidate gene studies
  - sweat gland activity, 440
  - Swedish Twin Registry, 515
  - Swiss mice, 435
  - Swiss-Webster mice, 158, 161
  - switching, between trajectory groups, 255–256
  - systems biology, genetic association studies and, 554, 556–559
  - Systems Biology Markup Language (SBML), 561
- T**
- targeted treatment, 21–22
  - task performance, with nicotine deprivation, 436
  - Taste/Sensory Processes subscale, 84
  - taxon, nicotine dependence as, 93–94
  - team sports, as protective factor, 343
  - temperament-based model, 343–344
  - test performance, definition of, 75
  - test-retest reliability
    - acoustic startle reflex, 446–447
    - ad lib smoking, 415
    - diagnostic tools, 81
    - event-related potential, 429
    - mood effects tests, 375
    - nicotine-dependence measures, 26–27, 29
    - prepulse inhibition startle response, 431
  - thapsigargin, 167
  - theta rhythm (slow-wave activity), 362
  - TH* (tyrosine hydroxylase) gene, 198–199, 269
  - threshold
    - factor loading and, 275–276
    - as “smoker,” 42
  - time to first cigarette (TTFC)
    - during adolescence, as indicator of adult nicotine dependence, 230–231
    - correlations with, 90
    - factor analysis of, 88–89
    - nicotine metabolism association, 34
    - as physical dependence measure, 79
    - as quitting predictor, 80
  - time to maximum tolerance, 162
  - tissue levels of nicotine, *versus* plasma levels, 145
  - Tobacco Craving Questionnaire, 440
  - tobacco dependence, *versus* nicotine dependence, 75
  - tobacco industry, 24
  - tobacco policies, 7, 20
    - effect on adolescent smoking, 193
    - substance use policies and, 294–295
  - tobacco settlement dollars, 4
  - tobacco use. *See also* nicotine
    - frequency of, 4, 21, 369–371, 413, 420
    - history of, 27–28
    - smoke compounds, 588
  - Tobacco Use Supplement, U.S. Census Bureau' Current Population Survey (TUS-CPS), 487, 494, 498–500
  - tolerance, 162–168
    - genetic effects on, 146
    - to repeated doses, 158
  - Tolerance subscale, 90–91
  - TPH* gene, 269
  - trace fear conditioning, 435
  - transcription factor CREB, 160–161, 422
  - transcription factor Fosb, 161
  - transcription levels, alterations in, 40
  - transdisciplinary framework, 521
  - Transdisciplinary Tobacco Use Research Center (TTURC), 89, 521, 526
  - transitional endophenotypes, 107, 108, 200, 233
  - translational validation, 172

*TRPC7* gene, 43  
 129T2/SvEmsJ mice, 431  
 TTFC. *See* time to first cigarette  
 TTURC (Transdisciplinary Tobacco Use Research Center), 89, 521, 526  
 TUS-CPS (Tobacco Use Supplement, U.S. Census Bureau's Current Population Survey), 487, 494, 498–500  
 twin studies. *See also specific study*  
     adolescent smoking, 196, 259–262  
     affective regulation, 446–447  
     cross-substance concordance, 317, 319–320  
     *CYP2A6* effect, 40  
     delay aversion, 351  
     ecological momentary assessment, 527  
     epigenetic regulation, 36  
     equal environments assumption in, 516–517  
     event-related potential, 428, 429  
     extended family, 250–252, 262, 279, 280  
     factor analysis, 88–89  
     factor mixture models, 256  
     genomic studies, 589  
     heritability of dependence, 28–29, 86, 342, 406  
     impulsivity, 451  
     limitations of, 279–280  
     multivariate, 252–253, 262–263  
     P300 amplitude, 360  
     prepulse inhibition startle response, 431  
     resting EEG, 426  
     smoking habits, 22, 24  
     smoking initiation, 31  
     structural equation modeling, 249–257  
     substance-use comorbidity, 307–323, 324  
     tobacco use history, 38–39  
     working memory, 437  
 two-factor structure, 81  
 two-stage genetic models, initiation *versus* progression, 323  
 tyrosine hydroxylase (*TH*) gene, 198–199, 269

## U

univariate analysis, substance-use comorbidity, 322  
 unmeasured genetic factors, 4, 511  
 upregulation, of nicotinic receptors, 135, 144–145, 588  
 U.S. Task Force on Community Preventive Services, 4

## V

validity  
     of developmental trajectory research, 234  
     discriminative, 77  
     predictive, 50, 90–91  
 variables  
     independent, 495–496, 498–499  
     observable, 77  
     ontologies to represent knowledge about, 558–559  
     perturbations from, 554  
     in structural equation modeling, 248  
 variance  
     in latent growth curve models, 254  
     partitioning, 270, 280  
 ventral tegmental area (VTA), 152–153, 588  
 verapamil, 167, 168  
 VET (Vietnam Era Twin) Registry, 263–264  
 videotaped paradigms, 524  
 Vietnam Era Twin (VET) Registry, 263–264  
 vigilance, 432–434  
 Virginia 30,000 Study, 262  
 Virginia Twin Registry, 264, 269–279  
     measures, 269–270  
     methods, 270  
     results, 270–276  
     study conclusions, 276–279  
     subjects, 269  
 Virginia Twin Study of Adolescent Behavioral Development, 260  
 visuospatial attention, 434  
 VLMR LR (Vuong-Lo-Mendell-Rubin likelihood ratio) test, 310, 312  
 VTA (ventral tegmental area), 152–153, 588  
 vulnerability  
     of adolescents, 343  
     differences in, 22  
     modeling, 363–364  
     protective factors, 87, 114  
 Vuong-Lo-Mendell-Rubin likelihood ratio (VLMR LR) test, 310, 312

## W

Wald test, 544  
 watershed model, 78–79, 81–82  
 WCST (Wisconsin Card Sorting Test), 378, 436  
 Wechsler Adult Intelligence Test-Revised, 433  
 whole-environment scan, 530

- whole-genome association studies, 4
  - whole-genome linkage scans, 589
  - whole-genome quantitative transcript screening, 170
  - wild-type mice
    - conditioned place preference in, 159, 160, 161, 422
    - in CREB activation studies, 160
    - in nicotine reinforcement, 159
  - WinBUGS, 549
  - Wisconsin Card Sorting Test (WCST), 378, 436
  - Wisconsin Inventory of Smoking Dependence Motives (WISDM), 84–86, 85
    - as assessment tool, 405
    - subscales of, 90–93
  - Wistar rats, 425, 431, 449
  - withdrawal symptoms
    - in adolescents, 192, 194
    - cellular changes, 147
    - environmental influences in, 95
    - heritability of, 30
    - negative affect, 443
    - physical dependence inference from, 77
    - physiology of, 588
    - severity of, 97, 413, 447
    - support interval for, 40
  - within-class variability, estimation of, 220–221, 233
  - working memory, 434–438
  - World Health Organization, 80–81
  - World Mental Health Survey Initiative, 81
- 
- ## X
- XSEM (extended structural equation modeling), 249
- 
- ## Z
- zygosity, function of, in substance-use phenotypes, 316–317, 320





