



NCI Protocol #: 10323

Patient ID:

Date:

1 of 7

Patient Name:
Date Collected:
Biopsy Site: **Liver**
Sample ID:
Telephone:
Tumor Content (%): **50_69%**

Patient DOB:
Referring Physician:
Primary Diagnosis: **Small cell lung carcinoma**
MoCha ID:
Fax:

Tumor content assessment performed by Van Andel Research Institute; Grand Rapids; MI 49503

Table of Contents

Page

Variant Details	1
Biomarker Descriptions	2
Relevant Therapy Summary	3
Clinical Trials Summary	3

Report Highlights

2 Relevant Biomarkers
0 Therapies Available
13 Clinical Trials

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IIC	TP53 E285K	None	None	10
IIC	RB1 K720*	None	None	3

Public data sources included in relevant therapies: FDA¹, NCCN

Tier Reference: Li et al. *Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists.* J Mol Diagn. 2017 Jan;19(1):4-23.

Variant Details

DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect
RB1	p.(K720*)	c.2158A>T	.	chr13:49037918	71.42%	NM_000321.2	nonsense
TP53	p.(E285K)	c.853G>A	COSM10722	chr17:7577085	71.49%	NM_000546.5	missense

Comments:

N/A

Signed By: _____

Biomarker Descriptions

RB1 (RB transcriptional corepressor 1)

Background: The RB1 gene encodes the retinoblastoma protein (pRB), and is an early molecular hallmark of cancer. RB1 belongs to the family of pocket proteins that also includes p107 and p130, which play a crucial role in the cell proliferation, apoptosis, and differentiation^{1,2}. RB1 is well characterized as a tumor suppressor gene that restrains cell cycle progression from G1 phase to S phase³. Specifically, RB1 binds and represses the E2F family of transcription factors that regulate the expression of genes involved in the G1/S cell cycle regulation^{1,2,4}. Germline mutations in RB1 are associated with retinoblastoma (a rare childhood tumor) as well as other cancer types such as osteosarcoma, soft tissue sarcoma, and melanoma⁵.

Alterations and prevalence: Recurrent somatic alterations in RB1, including mutations and biallelic loss, lead to the inactivation of the RB1 protein. RB1 mutations are observed in urothelial carcinoma (approximately 16%), endometrial cancer (approximately 12%), and sarcomas (approximately 9%)⁶. Similarly, biallelic loss of RB1 is observed in sarcomas (approximately 13%), urothelial carcinoma (approximately 6%), and endometrial cancer (approximately 1%)⁶. Biallelic loss of the RB1 gene is also linked to the activation of chemotherapy-induced acute myeloid leukemia (AML) and acute lymphoblastic leukemia (ALL)^{7,8,9}.

Potential relevance: Currently, there are no therapies approved for RB1 aberrations.

TP53 (tumor protein p53)

Background: The TP53 gene encodes the p53 tumor suppressor protein that binds to DNA and activates transcription in response to diverse cellular stresses to induce cell cycle arrest, apoptosis, or DNA repair. In unstressed cells, TP53 is kept inactive by targeted degradation via MDM2, a substrate recognition factor for ubiquitin-dependent proteolysis. Alterations in TP53 is required for oncogenesis as they result in loss of protein function and gain of transforming potential¹⁰. Germline mutations in TP53 are the underlying cause of Li-Fraumeni syndrome, a complex hereditary cancer predisposition disorder associated with early-onset cancers^{11,12}.

Alterations and prevalence: TP53 is the most frequently mutated gene in the cancer genome with approximately half of all cancers experiencing TP53 mutations. Ovarian, head and neck, esophageal, and lung squamous cancers have particularly high TP53 mutation rates (60-90%)^{6,13,14,15,16,17}. Approximately two-thirds of TP53 mutations are missense mutations and several recurrent missense mutations are common including substitutions at codons R158, R175, Y220, R248, R273, and R282^{6,13}. Invariably, recurrent missense mutations in TP53 inactivate its ability to bind DNA and activate transcription of target genes^{18,19,20,21}.

Potential relevance: The small molecule p53 reactivator, PC14586, received a fast track designation (2020) by the FDA for advanced tumors harboring a TP53 Y220C mutation²². The FDA has granted fast track designation (2019) to the p53 reactivator, APR-246 alone²³ and breakthrough designation²⁴ (2020) in combination with azacitidine for myelodysplastic syndrome (MDS) and acute myeloid leukemia patients (AML) harboring a TP53 mutation. In addition to investigational therapies aimed at restoring wild-type TP53 activity, compounds that induce synthetic lethality are also under clinical evaluation^{25,26}. TP53 mutations confer poor prognosis in multiple blood cancers including AML, MDS, myeloproliferative neoplasms (MPN), chronic lymphocytic leukemia (CLL)^{27,28,29,30}. In mantle cell lymphoma, TP53 mutations are associated with poor prognosis when treated with conventional therapy including hematopoietic cell transplant³¹. Mono- and bi-allelic mutations in TP53 confer unique characteristics in MDS, with multi-hit patients also experiencing associations with complex karyotype, few co-occurring mutations, and high-risk disease presentation as well as predicted death and leukemic transformation independent of the IPSS-R staging system³².

Relevant Therapy Summary

In this cancer type

In other cancer type

In this cancer type and other cancer types

No evidence

TP53 E285K

Relevant Therapy	FDA	NCCN	Clinical Trials*
atezolizumab	×	×	● (II)
niraparib	×	×	● (II)
olaparib	×	×	● (II)
talazoparib	×	×	● (II)
eprenetapopt, pembrolizumab	×	×	● (I/II)
BAY-1895344, niraparib	×	×	● (I)
HWH-340	×	×	● (I)
talazoparib, palbociclib, axitinib, crizotinib	×	×	● (I)

RB1 K720*

Relevant Therapy	FDA	NCCN	Clinical Trials*
palbociclib	×	×	● (II)
SY-5609	×	×	● (I)

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Clinical Trials Summary

TP53 E285K

NCT ID	Title	Phase
NCT04383938	Study of APR-246 in Combination With Pembrolizumab in Subjects With Solid Tumor Malignancies	I/II
NCT02029001	A Two-period, Multicenter, Randomized, Open-label, Phase II Study Evaluating the Clinical Benefit of a Maintenance Treatment Targeting Tumor Molecular Alterations in Patients With Progressive Locally-advanced or Metastatic Solid Tumors MOST: My own specific treatment	II
NCT03297606	Canadian Profiling and Targeted Agent Utilization Trial (CAPTUR): A Phase II Basket Trial	II
NCT02401347	A Phase II Clinical Trial of the PARP Inhibitor Talazoparib in BRCA1 and BRCA2 Wild Type Patients With Advanced Triple Negative Breast Cancer and Homologous Recombination Deficiency or Advanced HER2 Negative Breast Cancer or Other Solid Tumors With a Mutation in Homologous Recombination Pathway Genes Talazoparib Beyond BRCA (TBB) Trial	II
NCT03415659	A Phase I, Open-label, Single-center, Single/Multiple-dose, Dose-escalation/Dose-expansion Clinical Study on Tolerance and Pharmacokinetics of HWH340 Tablet in Patients With Advanced Solid Tumors	I

Clinical Trials Summary (continued)

TP53 E285K (continued)

NCT ID	Title	Phase
NCT03767075	Basket of Baskets: A Modular, Open-label, Phase II, Multicentre Study To Evaluate Targeted Agents in Molecularly Selected Populations With Advanced Solid Tumours	II
NCT03207347	A Phase II Trial of the PARP Inhibitor, Niraparib, in BAP1 and Other DNA Damage Response (DDR) Pathway Deficient Neoplasms (UF-STO-ETI-001).	II
NCT03233204	NCI-COG Pediatric MATCH (Molecular Analysis for Therapy Choice)- Phase 2 Subprotocol of Olaparib in Patients With Tumors Harboring Defects in DNA Damage Repair Genes	II
NCT04267939	An Open-label Phase Ib Study to Determine the Maximum Tolerated and/or Recommended Phase II Dose of the ATR Inhibitor BAY 1895344 in Combination With PARP Inhibitor Niraparib, in Patients With Recurrent Advanced Solid Tumors and Ovarian Cancer	I
NCT04693468	Modular Phase IB Hypothesis-Testing, Biomarker-Driven, Talazoparib Combination Trial (TalaCom)	I

RB1 K720*

NCT ID	Title	Phase
NCT03155620	NCI-COG Pediatric MATCH (Molecular Analysis for Therapy Choice) Screening Protocol	II
NCT03526250	NCI-COG Pediatric MATCH (Molecular Analysis for Therapy Choice) - Phase 2 Subprotocol of Palbociclib in Patients With Tumors Harboring Activating Alterations in Cell Cycle Genes	II
NCT04247126	A Phase I Study of SY 5609, an Oral, Selective CDK7 Inhibitor, in Adult Patients With Select Advanced Solid Tumors	I

Assay Information

Methodology, Scope, and Application: The OCAv3 next-generation sequencing (NGS) assay identifies more than 3000 annotated mutations of interest (MOIs) broadly categorized into 5 mutation types: single nucleotide variants (SNVs), small insertions/deletions (Indels), large (>3 bases) insertions/deletions (Large Indels), copy number variants (CNVs), and gene fusions. The assay utilizes the Thermo Fisher Scientific Oncomine® Comprehensive Assay v3.0 (OCAv3), a next-generation sequencing (NGS) assay that utilizes a multiplex polymerase chain reaction (PCR) with DNA and RNA extracted from formalin-fixed tissue for sequencing on the Ion S5 XL platform and analyzed by the current version of Torrent Suite Software and Ion Reporter. The OCAv3 NGS Assay currently can reliably identify the presence or absence of >3000 known mutations and polymorphisms in the 161 unique genes compared to the Human Reference Genome hg19, including the genes listed below. The OCAv3 assay is a laboratory developed test designed to find gene mutations within tumors (somatic mutations).

Analytical Sensitivity and Specificity: The OCAv3 NGS Assay has been determined to be suitably analytically sensitive and specific for the various types of abnormalities within its reportable range, demonstrating 97.18% sensitivity compared with orthogonal assay results and 100% specificity for the 5 mutation types reported. 99.99% or greater reproducibility has been demonstrated. All quality measures for this assay were within defined assay parameters.

Hereditary and Germline Mutations: This assay examines tumor tissue only and does not examine normal (non-tumor) tissue. Mutations detected by the assay may be present only in the tumor, or in every cell of the body (including non-tumor cells). This test also cannot tell whether a potential germline mutation causes or will cause a hereditary cancer syndrome. If the patient's history includes any one or combination of features suggestive of a possible hereditary cancer predisposition (such as, cancer arising at a young age, especially a cancer of a type unusual in younger patients; a personal history of multiple different types of cancers; or a significant

cancer history among blood relatives, especially but not exclusively of the same type as the patient's) it is recommended that the physician arrange for the patient to meet with a genetic counselor and, if warranted, undergo the appropriate genetic test on normal (i.e. non-tumor) tissue (blood or cells brushed from the oral surface of the cheek) to check for a germline abnormality, regardless of the results of this research study.

Report Content: Thermo Fisher Scientific's Ion Torrent OncoPrint Reporter software was used in the generation of this report. Software was developed and designed internally by Thermo Fisher Scientific. The analysis was based on the current version of OncoPrint Reporter. The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. FDA information was sourced from www.fda.gov and is current as of this version. NCCN information was sourced from www.nccn.org and reflects its current version. Clinical Trials information reflects the current version. For the most up-to-date information regarding a particular trial, search www.clinicaltrials.gov by NCT ID or search local clinical trials authority website by local identifier listed in 'Other Identifiers.' Variants are reported according to HGVS nomenclature and classified following AMP/ASCO/CAP guidelines (Li et al. 2017). Based on the data sources selected, variants, therapies, and trials identified in this report are listed in order of potential clinical significance but not for predicted efficacy of the therapies.

Genes assayed for hotspot mutations:

AKT1, AKT2, AKT3, ALK, AR, ARAF, AXL, BRAF, BTK, CBL, CCND1, CDK4, CDK6, CHEK2, CSF1R, CTNNB1, DDR2, EGFR, ERBB2, ERBB3, ERBB4, ERCC2, ESR1, EZH2, FGFR1, FGFR2, FGFR3, FGFR4, FLT3, FOXL2, GATA2, GNA11, GNAQ, GNAS, H3F3A, HIST1H3B, HNF1A, HRAS, IDH1, IDH2, JAK1, JAK2, JAK3, KDR, KIT, KNSTRN, KRAS, MAGOH, MAP2K1, MAP2K2, MAP2K4, MAPK1, MAX, MDM4, MED12, MET, MTOR, MYC, MYCN, MYD88, NFE2L2, NRAS, NTRK1, NTRK2, NTRK3, PDGFRA, PDGFRB, PIK3CA, PIK3CB, PPP2R1A, PTPN11, RAC1, RAF1, RET, RHEB, RHOA, ROS1, SF3B1, SMAD4, SMO, SPOP, SRC, STAT3, TERT, TOP1, U2AF1, XPO1

Genes assayed with full exon coverage:

ARID1A, ATM, ATR, ATRX, BAP1, BRCA1, BRCA2, CDK12, CDKN1B, CDKN2A, CDKN2B, CHEK1, CREBBP, FANCA, FANCD2, FANCI, FBXW7, MLH1, MRE11, MSH2, MSH6, NBN, NF1, NF2, NOTCH1, NOTCH2, NOTCH3, PALB2, PIK3R1, PMS2, POLE, PTCH1, PTEN, RAD50, RAD51, RAD51B, RAD51C, RAD51D, RB1, RNF43, SETD2, SLX4, SMARCA4, SMARCB1, STK11, TP53, TSC1, TSC2

Genes assayed for copy number variations:

AKT1, AKT2, AKT3, ALK, AR, AXL, BRAF, CCND1, CCND2, CCND3, CCNE1, CDK2, CDK4, CDK6, EGFR, ERBB2, ESR1, FGF19, FGF3, FGFR1, FGFR2, FGFR3, FGFR4, FLT3, IGF1R, KIT, KRAS, MDM2, MDM4, MET, MYC, MYCL, MYCN, NTRK1, NTRK2, NTRK3, PDGFRA, PDGFRB, PIK3CA, PIK3CB, PPARG, RICTOR, TERT

Genes assayed for detection of fusions:

AKT2, ALK, AR, AXL, BRAF, BRCA1, BRCA2, CDKN2A, EGFR, ERBB2, ERBB4, ERG, ESR1, ETV1, ETV4, ETV5, FGFR1, FGFR2, FGFR3, FGR, FLT3, JAK2, KRAS, MDM4, MET, MYB, MYBL1, NF1, NOTCH1, NOTCH4, NRG1, NTRK1, NTRK2, NTRK3, NUTM1, PDGFRA, PDGFRB, PIK3CA, PPARG, PRKACA, PRKACB, PTEN, RAD51B, RAF1, RB1, RELA, RET, ROS1, RSP02, RSP03, TERT

DISCLAIMER

This assay is considered a Laboratory Developed Test (LDT). Its performance characteristics have been determined through extensive testing although it has not been cleared by the US Food and Drug Administration and such approval is not required for clinical implementation. Furthermore, any comments included in the report are strictly interpretive and the opinion of the reviewer. This laboratory is certified under the Clinical Laboratory Improvements Amendments (CLIA) for the performance of high-complexity testing for clinical purposes. The correlative data presented is a result of the curation of published data sources, but may not be exhaustive. The content of this report has not been evaluated or approved by the FDA or other regulatory agencies.

References

1. Korenjak et al. E2F-Rb complexes regulating transcription of genes important for differentiation and development. PMID: 16081278
2. Sachdeva et al. Understanding pRb: toward the necessary development of targeted treatments for retinoblastoma. *J. Clin. Invest.* 2012 Feb;122(2):425-34. PMID: 22293180
3. Dyson. RB1: a prototype tumor suppressor and an enigma. *Genes Dev.* 2016 Jul 1;30(13):1492-502. PMID: 27401552
4. Cobrinik. Pocket proteins and cell cycle control. *Oncogene.* 2005 Apr 18;24(17):2796-809. PMID: 15838516
5. Dommering et al. RB1 mutations and second primary malignancies after hereditary retinoblastoma. *Fam. Cancer.* 2012 Jun;11(2):225-33. PMID: 22205104
6. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
7. Anasua et al. Acute lymphoblastic leukemia as second primary tumor in a patient with retinoblastoma. . PMID: 27433042
8. Tanaka et al. Frequent allelic loss of the RB, D13S319 and D13S25 locus in myeloid malignancies with deletion/translocation at 13q14 of chromosome 13, but not in lymphoid malignancies. *Leukemia.* 1999 Sep;13(9):1367-73. PMID: 10482987
9. Gombos et al. Secondary acute myelogenous leukemia in patients with retinoblastoma: is chemotherapy a factor?. *Ophthalmology.* 2007 Jul;114(7):1378-83. PMID: 17613328
10. Muller et al. Mutant p53 in cancer: new functions and therapeutic opportunities. *Cancer Cell.* 2014 Mar 17;25(3):304-17. PMID: 24651012
11. Olivier et al. TP53 mutations in human cancers: origins, consequences, and clinical use. *Cold Spring Harb Perspect Biol.* 2010 Jan;2(1):a001008. PMID: 20182602
12. Guha et al. Inherited TP53 Mutations and the Li-Fraumeni Syndrome. *Cold Spring Harb Perspect Med.* 2017 Apr 3;7(4). PMID: 28270529
13. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
14. Peter et al. Comprehensive genomic characterization of squamous cell lung cancers. *Nature.* 2012 Sep 27;489(7417):519-25. PMID: 22960745
15. Cancer Genome Atlas Network. Comprehensive genomic characterization of head and neck squamous cell carcinomas. *Nature.* 2015 Jan 29;517(7536):576-82. PMID: 25631445
16. Campbell et al. Distinct patterns of somatic genome alterations in lung adenocarcinomas and squamous cell carcinomas. *Nat. Genet.* 2016 Jun;48(6):607-16. PMID: 27158780
17. Cancer Genome Atlas Research Network. Integrated genomic characterization of oesophageal carcinoma. *Nature.* 2017 Jan 12;541(7636):169-175. doi: 10.1038/nature20805. Epub 2017 Jan 4. PMID: 28052061
18. Olivier et al. The IARC TP53 database: new online mutation analysis and recommendations to users. *Hum. Mutat.* 2002 Jun;19(6):607-14. PMID: 12007217
19. Rivlin et al. Mutations in the p53 Tumor Suppressor Gene: Important Milestones at the Various Steps of Tumorigenesis. *Genes Cancer.* 2011 Apr;2(4):466-74. PMID: 21779514
20. Petitjean et al. TP53 mutations in human cancers: functional selection and impact on cancer prognosis and outcomes. *Oncogene.* 2007 Apr 2;26(15):2157-65. PMID: 17401424
21. Soussi et al. Recommendations for analyzing and reporting TP53 gene variants in the high-throughput sequencing era. *Hum. Mutat.* 2014 Jun;35(6):766-78. PMID: 24729566
22. <https://www.globenewswire.com/news-release/2020/10/13/2107498/0/en/PMV-Pharma-Granted-FDA-Fast-Track-Designation-of-PC14586-for-the-Treatment-of-Advanced-Cancer-Patients-that-have-Tumors-with-a-p53-Y220C-Mutation.html>
23. <https://www.aprea.com/news/aprea-therapeutics-receives-fda-fast-track-designation-and-orphan-drug-designation-for-apr-246-for-the-treatment-of-myelodysplastic-syndromes-mds/>. <https://www.aprea.com/news/aprea-therapeutics-receives-fda-fast-track-designation-and-orphan-drug-designation-for-apr-246-for-the-treatment-of-myelodysplastic-syndromes-mds/>
24. <https://ir.aprea.com/news-releases/news-release-details/aprea-therapeutics-receives-fda-fast-track-designation>
25. Parrales et al. Targeting Oncogenic Mutant p53 for Cancer Therapy. *Front Oncol.* 2015 Dec 21;5:288. doi: 10.3389/fonc.2015.00288. eCollection 2015. PMID: 26732534

References (continued)

26. Zhao et al. Molecularly targeted therapies for p53-mutant cancers. *Cell. Mol. Life Sci.* 2017 Nov;74(22):4171-4187. PMID: 28643165
27. NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2021]
28. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 3.2021]
29. NCCN Guidelines® - NCCN-Myeloproliferative Neoplasms [Version 1.2020]
30. NCCN Guidelines® - NCCN-Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma [Version 3.2021]
31. NCCN Guidelines® - NCCN-B-Cell Lymphomas [Version 3.2021]
32. Bernard et al. Implications of TP53 allelic state for genome stability, clinical presentation and outcomes in myelodysplastic syndromes. *Nat. Med.* 2020 Aug 3. PMID: 32747829