



NCI Protocol #: 10323

Patient ID:

Date:

1 of 14

Patient Name:
Date Collected:
Biopsy Site: **Skin, right flank**
Sample ID:
Telephone:
Tumor Content (%): **20_49**

Patient DOB:
Referring Physician:
Primary Diagnosis: **Melanoma**
MoCha ID:
Fax:

Preanalytics to include tumor enrichment as required is performed by Van Andel Research Institute; Grand Rapids; MI 49503

Table of Contents

Page

Variant Details	2
Biomarker Descriptions	2
Relevant Therapy Summary	4
Clinical Trials Summary	6

Report Highlights

2 Relevant Biomarkers
11 Therapies Available
51 Clinical Trials

Relevant Melanoma Findings

Gene	Finding
BRAF	BRAF V600E
KIT	None detected
NRAS	None detected
NTRK1	None detected
NTRK2	None detected
NTRK3	None detected

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	BRAF V600E	atezolizumab + cobimetinib + vemurafenib ¹ binimetinib + encorafenib ¹ cetuximab + encorafenib ¹ cobimetinib + vemurafenib ¹ dabrafenib ¹ dabrafenib + trametinib ¹ trametinib ¹ vemurafenib ¹ encorafenib	binimetinib + encorafenib ¹ cetuximab + encorafenib ¹ dabrafenib ¹ dabrafenib + trametinib ¹ trametinib ¹ cobimetinib + vemurafenib encorafenib + panitumumab selumetinib vemurafenib	50

Public data sources included in relevant therapies: FDA¹, NCCN

Tier Reference: Li et al. *Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists*. J Mol Diagn. 2017 Jan;19(1):4-23.

Shahanawaz Jiwani, MD, PhD, Laboratory Director | CLIA ID 21D2097127

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.03(005). The content of this report has not been evaluated or approved by FDA, EMA or other regulatory agencies.

Relevant Biomarkers (continued)

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
III	SETD2 R950C	None	None	1

Public data sources included in relevant therapies: FDA1, NCCN
Tier Reference: Li et al. Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists. J Mol Diagn. 2017 Jan;19(1):4-23.

Prevalent cancer biomarkers without relevant evidence based on included data sources

RB1 W99*

Variant Details

DNA Sequence Variants								
Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Coverage	Transcript	Variant Effect
SETD2	p.(R950C)	c.2848C>T	.	chr3:47163278	33.55%	1997	NM_014159.6	missense
BRAF	p.(V600E)	c.1799T>A	COSM476	chr7:140453136	44.52%	1988	NM_004333.4	missense
RB1	p.(W99*)	c.297G>A	.	chr13:48916767	49.45%	2000	NM_000321.2	nonsense

Comments:

N/A

Report Signed By

Name	Role	Date
------	------	------

Biomarker Descriptions

BRAF (B-Raf proto-oncogene, serine/threonine kinase)

Background: The BRAF gene encodes the B-Raf proto-oncogene serine/threonine kinase, a member of the RAF family of serine/threonine protein kinases which also includes ARAF and RAF1 (CRAF). BRAF is among the most commonly mutated kinases in cancer. Activation of the MAPK pathway occurs through BRAF mutations and leads to an increase in cell division, dedifferentiation, and survival^{1,2}. BRAF mutations are categorized into three distinct functional classes namely, class 1, 2, and 3, and are defined by the dependency on the RAS pathway. Class 1 and 2 BRAF mutants are RAS-independent in that they signal as active monomers (Class 1) or dimers (Class 2) and become uncoupled from RAS GTPase signaling, resulting in constitutive activation of BRAF³. Class 3 mutants are RAS dependent as the kinase domain function is impaired or dead^{3,4,5}.

Alterations and prevalence: Recurrent somatic mutations in BRAF are observed in 40-60% of melanoma and thyroid cancer, approximately 10% of colorectal cancer, and about 2% of non-small cell lung cancer (NSCLC)^{6,7,8,9,10}. Mutations at V600 belong to class 1 and include V600E, the most recurrent somatic BRAF mutation across diverse cancer types^{4,11}. Class 2 mutations include K601E/N/T, L597Q/V, G469A/V/R/, G464V/E/, and BRAF fusions⁴. Class 3 mutations include D287H, V459L, G466V/E/A, S467L, G469E, and

Biomarker Descriptions (continued)

N581S/I4. BRAF V600E is universally present in hairy cell leukemia, mature B-cell cancer, and prevalent in histiocytic neoplasms^{12,13,14}. Other recurrent BRAF somatic mutations cluster in the glycine-rich phosphate-binding loop at codons 464-469 in exon 11 as well as additional codons flanking V600 in the activation loop¹¹. In primary cancers, BRAF amplification is observed in 8% of ovarian cancer and about 1% of breast cancer^{7,10}. BRAF fusions are mutually exclusive to BRAF V600 mutations and have been described in melanoma, thyroid cancer, pilocytic astrocytoma, NSCLC, and several other cancer types^{15,16,17,18,19}. Part of the oncogenic mechanism of BRAF gene fusions is the removal of the N-terminal auto-inhibitory domain leading to constitutive kinase activation^{5,15,17}.

Potential relevance: Vemurafenib²⁰ (2011) was the first targeted therapy approved for the treatment of patients with unresectable or metastatic melanoma with a BRAF V600E mutation. BRAF class 1 mutations, including V600E, are sensitive to vemurafenib, whereas class 2 and 3 mutations are insensitive⁴. BRAF kinase inhibitors including dabrafenib²¹ (2013) and encorafenib²² (2018) are also approved for the treatment of patients with unresectable or metastatic melanoma with BRAF V600E/K mutations. Encorafenib²² is approved in combination with cetuximab²³ (2020) for the treatment of BRAF V600E mutated colorectal cancer. Due to the tight coupling of RAF and MEK signaling, several MEK inhibitors have been approved for patients harboring BRAF alterations⁴. Trametinib²⁴ (2013) and binimetinib²⁵ (2018) were approved for the treatment of metastatic melanoma with BRAF V600E/K mutations. Combination therapies of BRAF plus MEK inhibitors have been approved in melanoma and NSCLC. The combinations of dabrafenib/trametinib (2015) and vemurafenib/cobimetinib²⁶ (2015) were approved for the treatment of patients with unresectable or metastatic melanoma with a BRAF V600E/K mutation. Subsequently, the combination of dabrafenib and trametinib was approved for metastatic NSCLC (2017) with a BRAF V600E mutation. The PD-L1 antibody, atezolizumab²⁷, has also been approved in combination with cobimetinib and vemurafenib for BRAF V600 mutation-positive unresectable or metastatic melanoma. In 2018, binimetinib²⁸ was also granted breakthrough designation in combination with cetuximab and encorafenib for BRAF V600E mutant metastatic colorectal cancer. The pan-RAF kinase inhibitor, tovorafenib (DAY-101), was granted breakthrough therapy designation (2020) by the FDA for pediatric patients with advanced low-grade glioma harboring activating RAF alterations²⁹. The ERK inhibitor ulixertinib³⁰ was also granted a fast track designation in 2020 for the treatment of patients with non-colorectal solid tumors harboring BRAF mutations G469A/V, L485W, or L597Q. The FDA granted fast track designation (2022) to the pan-RAF inhibitor, KIN-2787³¹, for the treatment of BRAF class II or III alteration-positive malignant or unresectable melanoma. BRAF fusion is a suggested mechanism of resistance to BRAF targeted therapy in melanoma³². Additional mechanisms of resistance to BRAF targeted therapy include BRAF amplification and alternative splice transcripts as well as activation of PI3K signaling and activating mutations in KRAS, NRAS, and MAP2K1/2 (MEK1/2)^{33,34,35,36,37,38,39}. Clinical responses to sorafenib and trametinib in limited case studies of patients with BRAF fusions have been reported¹⁹.

RB1 (RB transcriptional corepressor 1)

Background: The RB1 gene encodes the retinoblastoma protein (pRB), and is an early molecular hallmark of cancer. RB1 belongs to the family of pocket proteins that also includes p107 and p130, which play a crucial role in the cell proliferation, apoptosis, and differentiation^{40,41}. RB1 is well characterized as a tumor suppressor gene that restrains cell cycle progression from G1 phase to S phase⁴². Specifically, RB1 binds and represses the E2F family of transcription factors that regulate the expression of genes involved in the G1/S cell cycle regulation^{40,41,43}. Germline mutations in RB1 are associated with retinoblastoma (a rare childhood tumor) as well as other cancer types such as osteosarcoma, soft tissue sarcoma, and melanoma⁴⁴.

Alterations and prevalence: Recurrent somatic alterations in RB1, including mutations and biallelic loss, lead to the inactivation of the RB1 protein. RB1 mutations are observed in urothelial carcinoma (approximately 16%), endometrial cancer (approximately 12%), and sarcomas (approximately 9%)⁷. Similarly, biallelic loss of RB1 is observed in sarcomas (approximately 13%), urothelial carcinoma (approximately 6%), and endometrial cancer (approximately 1%)⁷. Biallelic loss of the RB1 gene is also linked to the activation of chemotherapy-induced acute myeloid leukemia (AML) and acute lymphoblastic leukemia (ALL)^{45,46,47}.

Potential relevance: Currently, there are no therapies approved for RB1 aberrations.

SETD2 (SET domain containing 2, histone lysine methyltransferase)

Background: The SETD2 gene encodes the SET domain containing 2 histone lysine methyltransferase, a protein responsible for the trimethylation of lysine-36 on histone H3 (H3K36)^{48,49}. Methylation of H3K36 is a hallmark of active transcription and can be either mono-, di-, or tri-methylated where di- and tri-methylation are thought to be responsible for transcriptional regulation⁵⁰. Trimethylation of H3K36 by SETD2 promotes post-transcriptional gene silencing and prevents aberrant transcriptional initiation^{51,52}. SETD2 trimethylation activity is also observed to be involved in DNA repair through the recruitment of DNA repair machinery⁴⁹. Specifically, H3K36 tri-methylation by SETD2 has been shown to regulate mismatch repair (MMR) in vivo, wherein the loss of SETD2 results in MMR deficiency

Biomarker Descriptions (continued)

(dMMR) and consequent microsatellite instability (MSI)⁵³. Both copy number deletion and mutations resulting in SETD2 loss of function have been observed in a variety of cancers, suggesting a tumor suppressor role for SETD2^{49,54}.

Alterations and prevalence: Inactivating somatic mutations in SETD2 were first described in clear cell renal cell carcinoma (ccRCC) and are observed to be predominantly missense or truncating^{10,54,55}. Mutations at codon R1625 are observed to be the most recurrent with R1625C having been identified to result in loss of SETD2 H3K36 trimethylase activity^{10,48}. SETD2 mutation is observed in about 14% of uterine cancer, 12% of ccRCC, 9% of mesothelioma, and 6-7% of melanoma, lung adenocarcinoma, papillary renal cell carcinoma (pRCC), colorectal and bladder cancers⁴⁸. Biallelic loss of SETD2 is observed in about 6% of diffuse large B-cell lymphoma, and about 3% of ccRCC and mesothelioma⁴⁸.

Potential relevance: Currently, no therapies are approved for SETD2 aberrations. Mutations in SETD2 can be used to support diagnosis of hepatosplenic T-cell lymphoma (HSTCL)⁵⁶.

Relevant Therapy Summary

In this cancer type

In other cancer type

In this cancer type and other cancer types

No evidence

BRAF V600E			
Relevant Therapy	FDA	NCCN	Clinical Trials*
dabrafenib			
dabrafenib + trametinib			
binimetinib + encorafenib			
cetuximab + encorafenib			
trametinib			
cobimetinib + vemurafenib			(II)
vemurafenib			(II)
atezolizumab + cobimetinib + vemurafenib			(II)
encorafenib			
encorafenib + panitumumab			
selumetinib			
encorafenib, binimetinib			(III)
encorafenib, binimetinib, pembrolizumab			(III)
nivolumab			(II/III)
atezolizumab, tiragolumab			(II)
binimetinib, encorafenib			(II)

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Relevant Therapy Summary (continued)

In this cancer type

In other cancer type

In this cancer type and other cancer types

No evidence

Relevant Therapy	FDA	NCCN	Clinical Trials*
binimetinib, encorafenib, pembrolizumab, radiation therapy	×	×	● (II)
botensilimab	×	×	● (II)
dabrafenib, trametinib	×	×	● (II)
encorafenib, binimetinib, nivolumab, ipilimumab	×	×	● (II)
HL-085, vemurafenib	×	×	● (II)
ipilimumab, binimetinib, nivolumab, encorafenib	×	×	● (II)
RO-7247669	×	×	● (II)
RX208	×	×	● (II)
tocilizumab, ipilimumab, nivolumab	×	×	● (II)
trametinib, dabrafenib	×	×	● (II)
vemurafenib, cobimetinib	×	×	● (II)
vemurafenib, cobimetinib, atezolizumab	×	×	● (II)
vemurafenib, cobimetinib, ipilimumab, nivolumab	×	×	● (II)
CFT-1946, trametinib	×	×	● (I/II)
encorafenib, binimetinib, palbociclib	×	×	● (I/II)
HH-2710	×	×	● (I/II)
INCAGN-2385, INCAGN02390, retifanlimab	×	×	● (I/II)
nivolumab, tumor infiltrating lymphocytes, aldesleukin	×	×	● (I/II)
OKI-179, binimetinib	×	×	● (I/II)
RX208, trametinib	×	×	● (I/II)
tazemetostat, trametinib, dabrafenib	×	×	● (I/II)
vemurafenib, metformin hydrochloride	×	×	● (I/II)
ABM-1310	×	×	● (I)
ABM-1310, cobimetinib	×	×	● (I)
binimetinib	×	×	● (I)

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Relevant Therapy Summary (continued)

In this cancer type

In other cancer type

In this cancer type and other cancer types

No evidence

BRAF V600E (continued)			
Relevant Therapy	FDA	NCCN	Clinical Trials*
binimetinib, nivolumab, encorafenib	×	×	● (I)
cobimetinib, belvarafenib	×	×	● (I)
cobimetinib, vemurafenib	×	×	● (I)
E6201, dabrafenib	×	×	● (I)
JSI-1187	×	×	● (I)
JSI-1187, dabrafenib	×	×	● (I)
nilotinib, dabrafenib, trametinib	×	×	● (I)
PF-07799544, PF 07284890, PF-07799933	×	×	● (I)
PF-07799933, cetuximab, binimetinib	×	×	● (I)
RO-7276389, cobimetinib	×	×	● (I)

SETD2 R950C			
Relevant Therapy	FDA	NCCN	Clinical Trials*
niraparib	×	×	● (II)

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Clinical Trials Summary

BRAF V600E		
NCT ID	Title	Phase
NCT03898908	Phase II, Multicentre Clinical Trial to Evaluate the Activity of Encorafenib and Binimetinib Administered Before Local Treatment in Patients With BRAF Mutant Melanoma Metastatic to the Brain.	II
NCT04074096	Randomised Phase II Trial Testing the Addition of Upfront Stereotactic Radiosurgery to Binimetinib Encorafenib Pembrolizumab Compared to Binimetinib Encorafenib Pembrolizumab Alone in BRAFV600 Mutation-positive Melanoma With Brain Metastasis	II
NCT05263453	A Single-arm, Multi-center Phase II Trial to Evaluate the Efficacy and Safety of the Combination of HL-085 and Vemurafenib in Advanced Melanoma Patients With BRAF V600E/K Mutation	II
NCT02968303	Phase II Study With COmbination of Vemurafenib With Cobimetinib in B-RAF V600E/K Mutated Melanoma Patients to Normalize LDH and Optimize Nivolumab and Ipilimumab therapy	II

Shahanawaz Jiwani, MD, PhD, Laboratory Director | CLIA ID 21D2097127

Clinical Trials Summary (continued)

BRAF V600E (continued)

NCT ID	Title	Phase
NCT04655157	A Multi-Center Phase I/II Open Label Study to Evaluate Safety and Efficacy in Participants With Metastatic BRAF-mutant Melanoma Treated With Encorafenib With and Without Binimetinib in Combination With Nivolumab and Low-dose Ipilimumab. (QUAD 01: Quadruple Therapy in Melanoma)	I/II
NCT04557956	Phase I/II Study of an EZH2 Inhibitor (Tazemetostat) in Combination With Dual BRAF/MEK Inhibition in Patients With BRAF- Mutated Metastatic Melanoma Who Progressed on Prior BRAF/MEK Inhibitor Therapy	I/II
NCT01638676	A Phase I/II Trial of Vemurafenib and Metformin to Unresectable Stage IIIC and Stage IV BRAF.V600E+ Melanoma Patients	I/II
NCT03543969	Pilot Study of Adaptive BRAF-MEK Inhibitor Therapy for Advanced BRAF Mutant Melanoma	I
NCT04722575	A Phase II Randomized Non-Comparative Study, With Neoadjuvant Plus Adjuvant Therapy With Combination Or Sequence Of Vemurafenib, Cobimetinib, And Atezolizumab In Patients With High-Risk, Surgically Resectable BRAF Mutated And Wild-Type Melanoma	II
NCT05026983	Phase II Study of Binimetinib With Encorafenib in Patients With Metastatic Melanoma and CNS Metastases	II
NCT05299580	"Dabrafenib and Trametinib in Circulating Free DNA BRAFV600 Mutated Metastatic Melanoma Patients: a Prospective Phase II, Open Label, Multicentre Study - (Bioliquid TAILOR Study - BIOTAILOR)"	II
NCT05114603	An Open-label, Multicenter Phase II Clinical Study to Evaluate Safety, Efficacy and PK of HLX208 for Advanced Melanoma With BRAF V600 Mutation	II
NCT05668585	A Phase I/II Open-Label Multicenter Trial to Characterize the Safety, Tolerability, and Preliminary Efficacy of CFT1946 as Monotherapy and in Combination With Trametinib in Subjects With BRAF V600 Mutant Solid Tumors	I/II
NCT04720768	A Phase IB Study to Evaluate the Safety, Tolerability and Preliminary Efficacy of the Combination of Encorafenib, Binimetinib and Palbociclib in Patients With BRAF-mutant Metastatic Melanoma (The CELEBRATE Study)	I/II
NCT04198818	A First-in-Human, Open Label, Phase I/II Study to Evaluate the Safety, Tolerability and Pharmacokinetics of HH2710 in Patients With Advanced Tumors	I/II
NCT03374839	Combined Therapy of Nivolumab and Adoptive T Cell Therapy in Metastatic Melanoma Patients: Pilot Study Phase I/II	I/II
NCT04741997	A Randomized Pilot Trial of Adjuvant Therapy Based on Pathologic Response After Neoadjuvant Encorafenib and Binimetinib in Advanced Melanoma	I
NCT03284502	A Phase Ib, Open-label, Multicenter, Dose Escalation Study of the Safety, Tolerability, and Pharmacokinetics of HM95573 in Combination With Either Cobimetinib or Cetuximab in Patients With Locally Advanced or Metastatic Solid Tumors	I
NCT05388877	Phase I Study of E6201 Plus Dabrafenib for the Treatment of Central Nervous System (CNS) Metastases From BRAF V600-Mutated Metastatic Melanoma	I
NCT04418167	A Phase I Study of ERK1/2 Inhibitor JSI-1187 Administered as Monotherapy and in Combination With Dabrafenib for the Treatment of Advanced Solid Tumors With MAPK Pathway Mutations	I

Clinical Trials Summary (continued)

BRAF V600E (continued)

NCT ID	Title	Phase
NCT04903119	A Phase I Study of Nilotinib in Combination With Dabrafenib and Trametinib in BRAF V600 Mutant Metastatic Melanoma After Progression on BRAF/MEK Inhibition	I
NCT05538130	A PHASE IA/B OPEN-LABEL MASTER STUDY OF PF-07799544 AS A SINGLE-AGENT AND IN COMBINATION WITH OTHER TARGETED AGENTS IN PARTICIPANTS WITH ADVANCED SOLID TUMORS	I
NCT03554083	NeoACTIVATE: Neoadjuvant Therapy for Patients With High Risk Stage III Melanoma: A Pilot Clinical Trial	II
NCT05419388	A Randomized, Open-Label, Multicenter, Phase II Study of Multiple Doses of RO7247669 in Participants With Previously Untreated Unresectable or Metastatic Melanoma	II
NCT04940299	A Phase II Study to Assess the Safety and Efficacy of Tocilizumab in Combination With Ipilimumab and Nivolumab in Patients With Advanced Melanoma, Non-Small Cell Lung Cancer, or Urothelial Carcinoma	II
NCT04370704	A Phase I/II Study of Combination Therapy With INCMGA00012 (Anti-PD-1), INCAGN02385 (Anti-LAG-3), and INCAGN02390 (Anti-TIM-3) in Participants With Select Advanced Malignancies	I/II
NCT05270044	Adjuvant Encorafenib & Binimetinib vs. Placebo in Fully Resected Stage IIB/C BRAF V600E/K Mutated Melanoma: a Randomized Triple-blind Phase III Study in Collaboration With the EORTC Melanoma	III
NCT04657991	A Phase III, Randomized, Double-blind Study of Encorafenib and Binimetinib Plus Pembrolizumab Versus Placebo Plus Pembrolizumab in Participants With BRAF V600E/K Mutation-Positive Metastatic OR Unresectable Locally Advanced Melanoma	III
NCT04901988	Circulating Tumour DNA Guided Therapy For Stage IIB/C Melanoma After Surgical Resection (DETECTION)	II/III
NCT04551521	Continuous ReAssessment With Flexible ExTension in Rare Malignancies - CRAFT: The NCT-PMO-1602 Phase II Trial	II
NCT02693535	Targeted Agent and Profiling Utilization Registry (TAPUR) Study	II
NCT03235245	Combination of Targeted Therapy (Encorafenib and Binimetinib) Followed by Combination of Immunotherapy (Ipilimumab and Nivolumab) vs Immediate Combination of Immunotherapy in Patients With Unresectable or Metastatic Melanoma With BRAF V600 Mutation : an EORTC Randomized Phase II Study (EBIN)	II
NCT04591431	The Rome Trial From Histology to Target: the Road to Personalize Target Therapy and Immunotherapy	II
NCT04190628	A Phase I, First-In-Human, Multicenter, Open-Label Study of ABM-1310, Administered Orally in Adult Patients With Advanced Solid Tumors	I
No NCT ID	Phase I Clinical Study To Evaluate The Safety, Tolerability, Pharmacokinetics And Preliminary Efficacy of JSI-1187 Capsules In The Treatment Of MAPK Signaling Pathway-Mutated Advanced Solid Tumors	I
No NCT ID	A Phase IA/B Open-label Study to Evaluate Safety, Pharmacokinetics, and Preliminary Clinical Activity of RO7276389 alone and in Combination with Cobimetinib in Participants with BRAF-V600 Mutation-positive Advanced Solid Tumor or BRAF-V600 Mutation-positive Melanoma with Central Nervous System Metastases	I
No NCT ID	Phase I study of the safety, tolerability, pharmacokinetics and preliminary efficacy of RX208 in patients with advanced malignant solid tumors	I

Clinical Trials Summary (continued)

BRAF V600E (continued)

NCT ID	Title	Phase
NCT03297606	Canadian Profiling and Targeted Agent Utilization Trial (CAPTUR): A Phase II Basket Trial	II
No NCT ID	Dabrafenib and Trametinib in Pediatric Patients with Refractory Advanced Solid Tumors with BRAF V600 Mutations	II
NCT04511013	A Randomized Phase II Trial of Encorafenib + Binimetinib + Nivolumab vs Ipilimumab + Nivolumab in BRAF-V600 Mutant Melanoma With Brain Metastases	II
NCT04116541	MegaMOST - A Multicenter, Open-label, Biology Driven, Phase II Study Evaluating the Activity of Anti-cancer Treatments Targeting Tumor Molecular Alterations /Characteristics in Advanced / Metastatic Tumors.	II
No NCT ID	Single Arm, Open Label, Phase II Trial Of Vemurafenib And Cobimetinib In Patients With Metastatic Non-Squamous Non-Small Cell Lung Cancer And Other Tumours Harboursing BRAF V600 Mutations Detected By Comprehensive Genomic Profiling	II
NCT04965220	A Phase Ib/II Clinical Trial Evaluating the Safety, Tolerability, Pharmacokinetics, and Initial Efficacy of HLX208 (BRAF V600E Inhibitor) in Combination With Trametinib in Patients With Advanced Solid Tumors	I/II
NCT05501912	A Phase I, Open-Label, Multicenter, Dose Escalation and Expansion Study to Evaluate the Safety, Tolerability, Pharmacokinetics, and Preliminary Anti-Cancer Activity of ABM-1310 in Patients With BRAF V600-Mutant Advanced Solid Tumors	I
NCT03781219	A Phase I, Single Arm, Dose Escalation Study to Evaluate Safety, Pharmacokinetics and Preliminary Efficacy of HL-085 Plus Vemurafenib in Patients With BRAF V600 Mutant Advanced Solid Tumor	I
NCT05355701	A PHASE I OPEN-LABEL, DOSE ESCALATION AND DOSE EXPANSION STUDY TO EVALUATE THE SAFETY, TOLERABILITY, PHARMACOKINETICS, AND ANTI TUMOR ACTIVITY OF PF-07799933 (ARRY-440) AS A SINGLE AGENT AND IN COMBINATION THERAPY IN PARTICIPANTS 16 YEARS AND OLDER WITH ADVANCED SOLID TUMORS WITH BRAF ALTERATIONS	I
NCT05529316	A Multicohort, Open Label, Phase 2 Study of Botensilimab (AGEN1181) for Treatment of Advanced Melanoma Refractory to Prior Checkpoint Inhibitor Therapy	II
NCT05340621	NAUTILUS: A Phase I b/II Study of OKI-179 Plus Binimetinib in Patients With Advanced Solid Tumors and Activating Mutations in the RAS Pathway (Phase 1b) and in Patients With Advanced NRAS-Mutated Melanoma (Phase 2)	I/II
NCT04423185	Platform Study of Genotyping Guided Precision Medicine for Rare Tumors in China	II
NCT04913285	A Phase I/I b Open-label, Multicenter Study to Investigate the Safety, Tolerability, Pharmacokinetics, and Antitumor Activity of KIN-2787 in Participants With BRAF and/or NRAS Mutation-positive Solid Tumors.	I

SETD2 R950C

NCT ID	Title	Phase
NCT04992013	Genomically Guided Phase II Study to Evaluate the Clinical Benefit of Niraparib in Tumors Metastatic to the CNS	II

Assay Information

Methodology, Scope, and Application: The OCAv3 next-generation sequencing (NGS) assay identifies more than 3000 annotated mutations of interest (MOIs) broadly categorized into 5 mutation types: single nucleotide variants (SNVs), small insertions/deletions (Indels), large (>3 bases) insertions/deletions (Large Indels), copy number variants (CNVs), and gene fusions. The assay utilizes the Thermo Fisher Scientific Oncomine® Comprehensive Assay v3.0 (OCAv3), a next-generation sequencing (NGS) assay that utilizes a multiplex polymerase chain reaction (PCR) with DNA and RNA extracted from formalin-fixed tissue for sequencing on the Ion S5 XL platform and analyzed by the current version of Torrent Suite Software and Ion Reporter. The OCAv3 NGS Assay currently can reliably identify the presence or absence of >3000 known mutations and polymorphisms in the 161 unique genes compared to the Human Reference Genome hg19, including the genes listed below. The OCAv3 assay is a laboratory developed test designed to find gene mutations within tumors (somatic mutations).

Analytical Sensitivity and Specificity: The OCAv3 NGS Assay has been determined to be suitably analytically sensitive and specific for the various types of abnormalities within its reportable range, demonstrating 97.18% sensitivity compared with orthogonal assay results and 100% specificity for the 5 mutation types reported. 99.99% or greater reproducibility has been demonstrated. All quality measures for this assay were within defined assay parameters.

Hereditary and Germline Mutations: This assay examines tumor tissue only and does not examine normal (non-tumor) tissue. Mutations detected by the assay may be present only in the tumor, or in every cell of the body (including non-tumor cells). This test also cannot tell whether a potential germline mutation causes or will cause a hereditary cancer syndrome. If the patient's history includes any one or combination of features suggestive of a possible hereditary cancer predisposition (such as, cancer arising at a young age, especially a cancer of a type unusual in younger patients; a personal history of multiple different types of cancers; or a significant cancer history among blood relatives, especially but not exclusively of the same type as the patient's) it is recommended that the physician arrange for the patient to meet with a genetic counselor and, if warranted, undergo the appropriate genetic test on normal (i.e. non-tumor) tissue (blood or cells brushed from the oral surface of the cheek) to check for a germline abnormality, regardless of the results of this research study.

Report Content: Thermo Fisher Scientific's Ion Torrent Oncomine Reporter software was used in the generation of this report. Software was developed and designed internally by Thermo Fisher Scientific. The analysis was based on the current version of Oncomine Reporter. The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. FDA information was sourced from www.fda.gov and is current as of this version. NCCN information was sourced from www.nccn.org and reflects its current version. Clinical Trials information reflects the current version. For the most up-to-date information regarding a particular trial, search www.clinicaltrials.gov by NCT ID or search local clinical trials authority website by local identifier listed in 'Other Identifiers.' Variants are reported according to HGVS nomenclature and classified following AMP/ASCO/CAP guidelines (Li et al. 2017). Based on the data sources selected, variants, therapies, and trials identified in this report are listed in order of potential clinical significance but not for predicted efficacy of the therapies.

Genes assayed for hotspot mutations:

AKT1, AKT2, AKT3, ALK, AR, ARAF, AXL, BRAF, BTK, CBL, CCND1, CDK4, CDK6, CHEK2, CSF1R, CTNNB1, DDR2, EGFR, ERBB2, ERBB3, ERBB4, ERCC2, ESR1, EZH2, FGFR1, FGFR2, FGFR3, FGFR4, FLT3, FOXL2, GATA2, GNA11, GNAQ, GNAS, H3F3A, HIST1H3B, HNF1A, HRAS, IDH1, IDH2, JAK1, JAK2, JAK3, KDR, KIT, KNSTRN, KRAS, MAGOH, MAP2K1, MAP2K2, MAP2K4, MAPK1, MAX, MDM4, MED12, MET, MTOR, MYC, MYCN, MYD88, NFE2L2, NRAS, NTRK1, NTRK2, NTRK3, PDGFRA, PDGFRB, PIK3CA, PIK3CB, PPP2R1A, PTPN11, RAC1, RAF1, RET, RHEB, RHOA, ROS1, SF3B1, SMAD4, SMO, SPOP, SRC, STAT3, TERT, TOP1, U2AF1, XPO1

Genes assayed with full exon coverage:

ARID1A, ATM, ATR, ATRX, BAP1, BRCA1, BRCA2, CDK12, CDKN1B, CDKN2A, CDKN2B, CHEK1, CREBBP, FANCA, FANCD2, FANCI, FBXW7, MLH1, MRE11, MSH2, MSH6, NBN, NF1, NF2, NOTCH1, NOTCH2, NOTCH3, PALB2, PIK3R1, PMS2, POLE, PTCH1, PTEN, RAD50, RAD51, RAD51B, RAD51C, RAD51D, RB1, RNF43, SETD2, SLX4, SMARCA4, SMARCB1, STK11, TP53, TSC1, TSC2

Genes assayed for copy number variations:

AKT1, AKT2, AKT3, ALK, AR, AXL, BRAF, CCND1, CCND2, CCND3, CCNE1, CDK2, CDK4, CDK6, EGFR, ERBB2, ESR1, FGF19, FGF3, FGFR1, FGFR2, FGFR3, FGFR4, FLT3, IGF1R, KIT, KRAS, MDM2, MDM4, MET, MYC, MYCL, MYCN, NTRK1, NTRK2, NTRK3, PDGFRA, PDGFRB, PIK3CA, PIK3CB, PPARG, RICTOR, TERT

Genes assayed for detection of fusions:

Shahanawaz Jiwani, MD, PhD, Laboratory Director | CLIA ID 21D2097127

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.03(005).

AKT2, ALK, AR, AXL, BRAF, BRCA1, BRCA2, CDKN2A, EGFR, ERBB2, ERBB4, ERG, ESR1, ETV1, ETV4, ETV5, FGFR1, FGFR2, FGFR3, FGR, FLT3, JAK2, KRAS, MDM4, MET, MYB, MYBL1, NF1, NOTCH1, NOTCH4, NRG1, NTRK1, NTRK2, NTRK3, NUTM1, PDGFRA, PDGFRB, PIK3CA, PPARG, PRKACA, PRKACB, PTEN, RAD51B, RAF1, RB1, RELA, RET, ROS1, RSP02, RSP03, TERT

DISCLAIMER

This assay is considered a Laboratory Developed Test (LDT). Its performance characteristics have been determined through extensive testing although it has not been cleared by the US Food and Drug Administration and such approval is not required for clinical implementation. Furthermore, any comments included in the report are strictly interpretive and the opinion of the reviewer. This laboratory is certified under the Clinical Laboratory Improvements Amendments (CLIA) for the performance of high-complexity testing for clinical purposes. The correlative data presented is a result of the curation of published data sources, but may not be exhaustive. The content of this report has not been evaluated or approved by the FDA or other regulatory agencies.

References

1. Cheng et al. Molecular testing for BRAF mutations to inform melanoma treatment decisions: a move toward precision medicine. *Mod. Pathol.* 2018 Jan;31(1):24-38. PMID: 29148538
2. Alrabadi et al. Detection of driver mutations in BRAF can aid in diagnosis and early treatment of dedifferentiated metastatic melanoma. *Mod. Pathol.* 2019 Mar;32(3):330-337. PMID: 30315274
3. Quan et al. The association between BRAF mutation class and clinical features in BRAF-mutant Chinese non-small cell lung cancer patients. *Journal of Translational Medicine*, 29 Aug 2019, 17(1):298. PMID: 31470866
4. Yao et al. Tumours with class 3 BRAF mutants are sensitive to the inhibition of activated RAS. *Nature*. 2017 Aug 10;548(7666):234-238. PMID: 28783719
5. Bracht et al. BRAF Mutations Classes I, II, and III in NSCLC Patients Included in the SLLIP Trial: The Need for a New Pre-Clinical Treatment Rationale. *Cancers (Basel)*. 2019 Sep 17;11(9). PMID: 31533235
6. Cancer Genome Atlas Research Network. Integrated genomic characterization of papillary thyroid carcinoma. *Cell*. 2014 Oct 23;159(3):676-90. PMID: 25417114
7. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
8. Donna et al. Comprehensive molecular characterization of human colon and rectal cancer. *Nature*. 2012 Jul 18;487(7407):330-7. PMID: 22810696
9. Cancer Genome Atlas Research Network. Comprehensive molecular profiling of lung adenocarcinoma. *Nature*. 2014 Jul 31;511(7511):543-50. doi: 10.1038/nature13385. Epub 2014 Jul 9. PMID: 25079552
10. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
11. Wan et al. Mechanism of activation of the RAF-ERK signaling pathway by oncogenic mutations of B-RAF. *Cell*. 2004 Mar 19;116(6):855-67. PMID: 15035987
12. Tiacci et al. BRAF mutations in hairy-cell leukemia. *N. Engl. J. Med.* 2011 Jun 16;364(24):2305-15. PMID: 21663470
13. Diamond et al. Diverse and Targetable Kinase Alterations Drive Histiocytic Neoplasms. *Cancer Discov.* 2016 Feb;6(2):154-65. doi: 10.1158/2159-8290.CD-15-0913. Epub 2015 Nov 13. PMID: 26566875
14. Imielinski et al. Oncogenic and sorafenib-sensitive ARAF mutations in lung adenocarcinoma. *J Clin Invest.* 2014 Apr;124(4):1582-6. doi: 10.1172/JCI72763. Epub 2014 Feb 24. PMID: 24569458
15. Ciampi et al. Oncogenic AKAP9-BRAF fusion is a novel mechanism of MAPK pathway activation in thyroid cancer. *J. Clin. Invest.* 2005 Jan;115(1):94-101. PMID: 15630448
16. Palanisamy et al. Rearrangements of the RAF kinase pathway in prostate cancer, gastric cancer and melanoma. *Nat. Med.* 2010 Jul;16(7):793-8. PMID: 20526349
17. Jones et al. Tandem duplication producing a novel oncogenic BRAF fusion gene defines the majority of pilocytic astrocytomas. *Cancer Res.* 2008 Nov 1;68(21):8673-7. PMID: 18974108
18. Cin et al. Oncogenic FAM131B-BRAF fusion resulting from 7q34 deletion comprises an alternative mechanism of MAPK pathway activation in pilocytic astrocytoma. *Acta Neuropathol.* 2011 Jun;121(6):763-74. doi: 10.1007/s00401-011-0817-z. Epub 2011 Mar 20. PMID: 21424530
19. Ross et al. The distribution of BRAF gene fusions in solid tumors and response to targeted therapy. *Int. J. Cancer.* 2016 Feb 15;138(4):881-90. PMID: 26314551
20. https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/202429s019lbl.pdf
21. https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/202806s022lbl.pdf
22. https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/210496s013lbl.pdf
23. https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125084s279lbl.pdf
24. https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/204114s024lbl.pdf
25. https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/210498s001lbl.pdf
26. https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/206192s005lbl.pdf

References (continued)

27. https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/761034s0471bl.pdf
28. <https://markets.businessinsider.com/news/stocks/array-biopharma-receives-fda-breakthrough-therapy-designation-for-braftovi-in-combination-with-mektovi-and-cetuximab-for-brafv600e-mutant-metastatic-colorectal-cancer-1027437791>
29. <https://ir.dayonebio.com/news-releases/news-release-details/day-one-receives-fda-rare-pediatric-disease-designation-day101>
30. <https://biomed-valley.com/news/#press-releases>
31. <https://investors.kinnate.com/news-releases/news-release-details/kinnate-biopharma-inc-receives-fast-track-designation-us-food>
32. Kulkarni et al. BRAF Fusion as a Novel Mechanism of Acquired Resistance to Vemurafenib in BRAFV600E Mutant Melanoma. *Clin. Cancer Res.* 2017 Sep 15;23(18):5631-5638. PMID: 28539463
33. Johnson et al. Acquired BRAF inhibitor resistance: A multicenter meta-analysis of the spectrum and frequencies, clinical behaviour, and phenotypic associations of resistance mechanisms. *Eur. J. Cancer.* 2015 Dec;51(18):2792-9. PMID: 26608120
34. Nazarian et al. Melanomas acquire resistance to B-RAF(V600E) inhibition by RTK or N-RAS upregulation. *Nature.* 2010 Dec 16;468(7326):973-7. doi: 10.1038/nature09626. Epub 2010 Nov 24. PMID: 21107323
35. Rizos et al. BRAF inhibitor resistance mechanisms in metastatic melanoma: spectrum and clinical impact. *Clin. Cancer Res.* 2014 Apr 1;20(7):1965-77. PMID: 24463458
36. Shi et al. A novel AKT1 mutant amplifies an adaptive melanoma response to BRAF inhibition. *Cancer Discov.* 2014 Jan;4(1):69-79. PMID: 24265152
37. Van et al. The genetic landscape of clinical resistance to RAF inhibition in metastatic melanoma. *Cancer Discov.* 2014 Jan;4(1):94-109. doi: 10.1158/2159-8290.CD-13-0617. Epub 2013 Nov 21. PMID: 24265153
38. Villanueva et al. Concurrent MEK2 mutation and BRAF amplification confer resistance to BRAF and MEK inhibitors in melanoma. *Cell Rep.* 2013 Sep 26;4(6):1090-9. PMID: 24055054
39. Shi et al. Acquired resistance and clonal evolution in melanoma during BRAF inhibitor therapy. *Cancer Discov.* 2014 Jan;4(1):80-93. PMID: 24265155
40. Korenjak et al. E2F-Rb complexes regulating transcription of genes important for differentiation and development. *Curr Opin Genet Dev.* 2005 Oct;15(5):520-7. doi: 10.1016/j.gde.2005.07.001. PMID: 16081278
41. Sachdeva et al. Understanding pRb: toward the necessary development of targeted treatments for retinoblastoma. *J. Clin. Invest.* 2012 Feb;122(2):425-34. PMID: 22293180
42. Dyson. RB1: a prototype tumor suppressor and an enigma. *Genes Dev.* 2016 Jul 1;30(13):1492-502. PMID: 27401552
43. Cobrinik. Pocket proteins and cell cycle control. *Oncogene.* 2005 Apr 18;24(17):2796-809. PMID: 15838516
44. Dommering et al. RB1 mutations and second primary malignancies after hereditary retinoblastoma. *Fam. Cancer.* 2012 Jun;11(2):225-33. PMID: 22205104
45. Anasua et al. Acute lymphoblastic leukemia as second primary tumor in a patient with retinoblastoma. *Oman J Ophthalmol.* May-Aug 2016;9(2):116-8. PMID: 27433042
46. Tanaka et al. Frequent allelic loss of the RB, D13S319 and D13S25 locus in myeloid malignancies with deletion/translocation at 13q14 of chromosome 13, but not in lymphoid malignancies. *Leukemia.* 1999 Sep;13(9):1367-73. PMID: 10482987
47. Gombos et al. Secondary acute myelogenous leukemia in patients with retinoblastoma: is chemotherapy a factor?. *Ophthalmology.* 2007 Jul;114(7):1378-83. PMID: 17613328
48. Hacker et al. Structure/Function Analysis of Recurrent Mutations in SETD2 Protein Reveals a Critical and Conserved Role for a SET Domain Residue in Maintaining Protein Stability and Histone H3 Lys-36 Trimethylation. *J. Biol. Chem.* 2016 Sep 30;291(40):21283-21295. PMID: 27528607
49. Fahey et al. SETting the Stage for Cancer Development: SETD2 and the Consequences of Lost Methylation. *Cold Spring Harb Perspect Med.* 2017 May 1;7(5). PMID: 28159833
50. Zaghi et al. H3K36 Methylation in Neural Development and Associated Diseases. *Front Genet.* 2020 Jan 9;10:1291. doi: 10.3389/fgene.2019.01291. eCollection 2019. PMID: 31998360
51. Suzuki et al. H3K36 methylation state and associated silencing mechanisms. *Transcription.* 2017 Jan;8(1):26-31. PMID: 27723431

References (continued)

52. Sun et al. H3K36me3, Message From Chromatin to DNA Damage Repair. *Cell Biosci* . 2020 Jan 31;10:9. doi: 10.1186/s13578-020-0374-z. eCollection 2020. PMID: 32021684
53. Li et al. The Histone Mark H3K36me3 Regulates Human DNA Mismatch Repair Through Its Interaction With MutSa. *Cell*. 2013 Apr 25;153(3):590-600. PMID: 23622243
54. Duns et al. Histone methyltransferase gene SETD2 is a novel tumor suppressor gene in clear cell renal cell carcinoma. *Cancer Res*. 2010 Jun 1;70(11):4287-91. PMID: 20501857
55. Dalgliesh et al. Systematic sequencing of renal carcinoma reveals inactivation of histone modifying genes. *Nature*. 2010 Jan 21;463(7279):360-3. PMID: 20054297
56. NCCN Guidelines® - NCCN-T-Cell Lymphomas [Version 1.2023]