



NCI Protocol #: 10323

Patient ID:

Date:

1 of 4

Patient Name:

Date Collected:

Biopsy Site: **Bone Marrow Aspirate**

Sample ID:

Telephone:

Tumor Content (%): **35%**

Patient DOB:

Referring Physician:

Primary Diagnosis: **Plasma Cell Neoplasm**

MoCha ID:

Fax:

Preanalytics to include tumor enrichment as required is performed by Van Andel Research Institute; Grand Rapids; MI 49503

Table of Contents

Variant Details

Biomarker Descriptions

Page

1

2

Report Highlights

0 Relevant Biomarkers

0 Therapies Available

0 Clinical Trials

Relevant Biomarkers

No biomarkers associated with relevant evidence found in this sample

Prevalent cancer biomarkers without relevant evidence based on included data sources

NRAS Q61H

Variant Details

DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect
NRAS	p.(Q61H)	c.183A>C	COSM586	chr1:115256528	46.20%	NM_002524.4	missense

Comments:

N/A

Report Signed By

Name	Role	Date

Shahanawaz Jiwani, MD, PhD, Laboratory Director | CLIA ID 21D2097127

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2022.10(005). The content of this report has not been evaluated or approved by FDA, EMA or other regulatory agencies.

Biomarker Descriptions

NRAS (NRAS proto-oncogene, GTPase)

Background: The NRAS proto-oncogene encodes a GTPase that functions in signal transduction and is a member of the RAS superfamily which also includes KRAS and HRAS. RAS proteins mediate the transmission of growth signals from the cell surface to the nucleus via the PI3K/AKT/MTOR and RAS/RAF/MEK/ERK pathways, which regulate cell division, differentiation, and survival^{1,2,3}.

Alterations and prevalence: Recurrent mutations in RAS oncogenes cause constitutive activation and are found in 20-30% of cancers. NRAS mutations are particularly common in melanomas (up to 25%) and are observed at frequencies of 5-10% in acute myeloid leukemia, colorectal, and thyroid cancers^{4,5}. The majority of NRAS mutations consist of point mutations at G12, G13, and Q61^{4,6}. Mutations at A59, K117, and A146 have also been observed but are less frequent^{7,8}.

Potential relevance: Currently, no therapies are approved for NRAS aberrations. The EGFR antagonists, cetuximab⁹ and panitumumab¹⁰, are contraindicated for treatment of colorectal cancer patients with NRAS mutations in exon 2 (codons 12 and 13), exon 3 (codons 59 and 61), and exon 4 (codons 117 and 146)⁸. NRAS mutations are associated with poor prognosis in patients with low-risk myelodysplastic syndrome¹¹ as well as melanoma¹². In a phase III clinical trial in patients with advanced NRAS-mutant melanoma, binimetinib improved progression free survival (PFS) relative to dacarbazine with median PFS of 2.8 and 1.5 months, respectively¹³.

Assay Information

Methodology, Scope, and Application: The OCAv3 next-generation sequencing (NGS) assay identifies more than 3000 annotated mutations of interest (MOIs) broadly categorized into 5 mutation types: single nucleotide variants (SNVs), small insertions/deletions (Indels), large (>3 bases) insertions/deletions (Large Indels), copy number variants (CNVs), and gene fusions. The assay utilizes the Thermo Fisher Scientific Oncomine® Comprehensive Assay v3.0 (OCAv3), a next-generation sequencing (NGS) assay that utilizes a multiplex polymerase chain reaction (PCR) with DNA and RNA extracted from formalin-fixed tissue for sequencing on the Ion S5 XL platform and analyzed by the current version of Torrent Suite Software and Ion Reporter. The OCAv3 NGS Assay currently can reliably identify the presence or absence of >3000 known mutations and polymorphisms in the 161 unique genes compared to the Human Reference Genome hg19, including the genes listed below. The OCAv3 assay is a laboratory developed test designed to find gene mutations within tumors (somatic mutations).

Analytical Sensitivity and Specificity: The OCAv3 NGS Assay has been determined to be suitably analytically sensitive and specific for the various types of abnormalities within its reportable range, demonstrating 97.18% sensitivity compared with orthogonal assay results and 100% specificity for the 5 mutation types reported. 99.99% or greater reproducibility has been demonstrated. All quality measures for this assay were within defined assay parameters.

Hereditary and Germline Mutations: This assay examines tumor tissue only and does not examine normal (non-tumor) tissue. Mutations detected by the assay may be present only in the tumor, or in every cell of the body (including non-tumor cells). This test also cannot tell whether a potential germline mutation causes or will cause a hereditary cancer syndrome. If the patient's history includes any one or combination of features suggestive of a possible hereditary cancer predisposition (such as, cancer arising at a young age, especially a cancer of a type unusual in younger patients; a personal history of multiple different types of cancers; or a significant cancer history among blood relatives, especially but not exclusively of the same type as the patient's) it is recommended that the physician arrange for the patient to meet with a genetic counselor and, if warranted, undergo the appropriate genetic test on normal (i.e. non-tumor) tissue (blood or cells brushed from the oral surface of the cheek) to check for a germline abnormality, regardless of the results of this research study.

Report Content: Thermo Fisher Scientific's Ion Torrent Oncomine Reporter software was used in the generation of this report. Software was developed and designed internally by Thermo Fisher Scientific. The analysis was based on the current version of Oncomine Reporter. The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. FDA information was sourced from www.fda.gov and is current as of this version. NCCN information was sourced from www.nccn.org and reflects its current version. Clinical Trials information reflects the current version. For the most up-to-date information regarding a particular trials, search www.clinicaltrials.gov by NCT ID or search local clinical trials authority website by local identifier listed in 'Other Identifiers.' Variants are reported according to HGVS nomenclature and classified following AMP/ASCO/CAP guidelines (Li et al. 2017). Based on the data sources selected, variants, therapies, and trials identified in this report are listed in order of potential clinical significance but not for predicted efficacy of the therapies.

Shahanawaz Jiwani, MD, PhD, Laboratory Director | CLIA ID 21D2097127

Electronically Signed By Shahanawaz Jiwani on 11:15:56 AM

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2022.10(005).

Genes assayed for hotspot mutations:

AKT1, AKT2, AKT3, ALK, AR, ARAF, AXL, BRAF, BTK, CBL, CCND1, CDK4, CDK6, CHEK2, CSF1R, CTNNB1, DDR2, EGFR, ERBB2, ERBB3, ERBB4, ERCC2, ESR1, EZH2, FGFR1, FGFR2, FGFR3, FGFR4, FLT3, FOXL2, GATA2, GNA11, GNAQ, GNAS, H3F3A, HIST1H3B, HNF1A, HRAS, IDH1, IDH2, JAK1, JAK2, JAK3, KDR, KIT, KNSTRN, KRAS, MAGOH, MAP2K1, MAP2K2, MAP2K4, MAPK1, MAX, MDM4, MED12, MET, MTOR, MYC, MYCN, MYD88, NFE2L2, NRAS, NTRK1, NTRK2, NTRK3, PDGFRA, PDGFRB, PIK3CA, PIK3CB, PPP2R1A, PTPN11, RAC1, RAF1, RET, RHEB, RHOA, ROS1, SF3B1, SMAD4, SMO, SPOP, SRC, STAT3, TERT, TOP1, U2AF1, XPO1

Genes assayed with full exon coverage:

ARID1A, ATM, ATR, ATRX, BAP1, BRCA1, BRCA2, CDK12, CDKN1B, CDKN2A, CDKN2B, CHEK1, CREBBP, FANCA, FANCD2, FANCI, FBXW7, MLH1, MRE11, MSH2, MSH6, NBN, NF1, NF2, NOTCH1, NOTCH2, NOTCH3, PALB2, PIK3R1, PMS2, POLE, PTCH1, PTEN, RAD50, RAD51, RAD51B, RAD51C, RAD51D, RB1, RNF43, SETD2, SLX4, SMARCA4, SMARCB1, STK11, TP53, TSC1, TSC2

Genes assayed for copy number variations:

AKT1, AKT2, AKT3, ALK, AR, AXL, BRAF, CCND1, CCND2, CCND3, CCNE1, CDK2, CDK4, CDK6, EGFR, ERBB2, ESR1, FGF19, FGF3, FGFR1, FGFR2, FGFR3, FGFR4, FLT3, IGF1R, KIT, KRAS, MDM2, MDM4, MET, MYC, MYCL, MYCN, NTRK1, NTRK2, NTRK3, PDGFRA, PDGFRB, PIK3CA, PIK3CB, PPARG, RICTOR, TERT

Genes assayed for detection of fusions:

AKT2, ALK, AR, AXL, BRAF, BRCA1, BRCA2, CDKN2A, EGFR, ERBB2, ERBB4, ERG, ESR1, ETV1, ETV4, ETV5, FGFR1, FGFR2, FGFR3, FGR, FLT3, JAK2, KRAS, MDM4, MET, MYB, MYBL1, NF1, NOTCH1, NOTCH4, NRG1, NTRK1, NTRK2, NTRK3, NUTM1, PDGFRA, PDGFRB, PIK3CA, PPARG, PRKACA, PRKACB, PTEN, RAD51B, RAF1, RB1, RELA, RET, ROS1, RSP02, RSP03, TERT

DISCLAIMER

This assay is considered a Laboratory Developed Test (LDT). Its performance characteristics have been determined through extensive testing although it has not been cleared by the US Food and Drug Administration and such approval is not required for clinical implementation. Furthermore, any comments included in the report are strictly interpretive and the opinion of the reviewer. This laboratory is certified under the Clinical Laboratory Improvements Amendments (CLIA) for the performance of high-complexity testing for clinical purposes. The correlative data presented is a result of the curation of published data sources, but may not be exhaustive. The content of this report has not been evaluated or approved by the FDA or other regulatory agencies.

References

1. Pylayeva-Gupta et al. RAS oncogenes: weaving a tumorigenic web. *Nat. Rev. Cancer*. 2011 Oct 13;11(11):761-74. PMID: 21993244
2. Karnoub et al. Ras oncogenes: split personalities. *Nat. Rev. Mol. Cell Biol.* 2008 Jul;9(7):517-31. PMID: 18568040
3. Scott et al. Therapeutic Approaches to RAS Mutation. *Cancer J.* 2016 May-Jun;22(3):165-74. doi: 10.1097/PPO.000000000000187. PMID: 27341593
4. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
5. Janku et al. PIK3CA mutations frequently coexist with RAS and BRAF mutations in patients with advanced cancers. *PLoS ONE*. 2011;6(7):e22769. PMID: 21829508
6. Ohashi et al. Characteristics of lung cancers harboring NRAS mutations. *Clin. Cancer Res.* 2013 May 1;19(9):2584-91. PMID: 23515407
7. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
8. Allegra et al. Extended RAS Gene Mutation Testing in Metastatic Colorectal Carcinoma to Predict Response to Anti-Epidermal Growth Factor Receptor Monoclonal Antibody Therapy: American Society of Clinical Oncology Provisional Clinical Opinion Update 2015. *J. Clin. Oncol.* 2016 Jan 10;34(2):179-85. PMID: 26438111
9. https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125084s279lbl.pdf
10. https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125147s210lbl.pdf
11. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 3.2022]
12. Johnson et al. Treatment of NRAS-Mutant Melanoma. *Curr Treat Options Oncol.* 2015 Apr;16(4):15. doi: 10.1007/s11864-015-0330-z. PMID: 25796376
13. Dummer et al. Binimetinib versus dacarbazine in patients with advanced NRAS-mutant melanoma (NEMO): a multicentre, open-label, randomised, phase 3 trial. *Lancet Oncol.* 2017 Apr;18(4):435-445. PMID: 28284557