

CHEMISTRY 456 / INTERNATIONAL WORKSHOP

Fundamentals of Biological Macromolecules

Andy Howard

Associate Professor of Biology & Physics,
Illinois Tech

Monday 28 June 2021


WHY BIOCHEMISTRY MATTERS

- ◆ Biochemists point out how chemical processes occur in biological systems, inside and between cells.
- ◆ These remarks are intended to introduce you to the chemistry of biopolymers and their biological significance.

WHAT WE'LL DISCUSS

- ◆ Unity and diversity
- ◆ Thermodynamics
- ◆ Organic chemistry
- ◆ Small biomolecules
- ◆ Polymers
- ◆ Proteins
 - ◆ Components
 - ◆ Structure
 - ◆ Functions
- ◆ Enzymes
- ◆ Nucleic acids
 - ◆ Components
 - ◆ Polymers
 - ◆ Structure
 - ◆ Function
- ◆ Carbohydrates

BIOCHEMICAL UNITY




- There is a remarkable unity to biochemical processes all across eubacteria, archaea, and eukaryota
- The genetic code is almost uniform throughout all 3 superkingdoms
- Specific proteins look the same across wide gaps of evolutionary history

BIOCHEMICAL DIVERSITY


◆ The morphological and functional differences between you and an elephant, or between you and a flowering plant, or between you and a bacterium, arise from differences in biochemical phenomena

- Some classes of proteins are present only in certain kinds of organisms
- Structural components in vertebrates differ significantly from those in arthropods




THERMODYNAMICS AND KINETICS

- ◆ Systems of molecules tend toward an equilibrium state in which the minimum free energy is obtained
- ◆ In order to go from an initial state R to a final, lower-energy state P , a higher-energy transition state T may have to be overcome
- ◆ Free energy has both enthalpic (heat) and entropic (organizational) components (Gibbs & Boltzmann): $\Delta G = \Delta H - T\Delta S$




BIOORGANIC CHEMISTRY

- Most biological reactions involve carbon-containing compounds
- Therefore organic chemistry (the chemistry of compounds that contain C-C or C-H bonds) explains a large fraction of what's happening below the surface in biochemical systems



WE CAN IGNORE A LOT OF ORGANIC REACTION MECHANISMS!

- Only a subset of all possible organic reactions occur in biochemistry:
 - They're almost all aqueous
 - Generally occur at temperatures between 3°C and 50°C
 - Most occur between pH 5 and pH 9 (exception in humans: the stomach, which is highly acidic)
 - Entire classes of organic reactions are unrepresented




Claisen
condensation

WHAT REACTIONS *DO* OCCUR?

- Mostly nucleophilic substitutions (S_N1 and S_N2)
 - These are two-electron transformations, as we've discussed in previous lectures
 - These take place under enzymatic control
 - Therefore they can proceed at measurable rates even though the nucleophiles aren't very powerful by organic lab standards
- Some free radical reactions, as we've said
 - There are enzymatic free-radical reactions
 - But there are non-enzymatic ones too


CHIRALITY



- ◆ Many biomolecules are chiral:
 - At least one carbon atom has 4 distinct substituents
 - Therefore the mirror image of the molecule is different from the original molecule
- ◆ Specific classes of biomolecules are chiral
- ◆ Standard amino acids are L- and so are their polymers
- ◆ Most sugars are D- and so are their polymers
- ◆ Lipids are usually not chiral
- ◆ Nucleic acid bases: achiral; but ribose is chiral

SMALL BIOMOLECULES

- Most common small molecule in biological systems is water
- There are other important small molecules and ions:
 - CO_2 , O_2 , glyceraldehyde, glucose
 - HCO_3^- , PO_4^{3-} , Na^+ , Cl^- , K^+ , Ca^{2+} , Fe^{2+} , Mg^{2+}
 - Palmitate, oleate, stearate, choline, sphingomyelin
 - Amino acids (20 ribosomal + others)
 - Nucleic acids (A, C, G, U, dA, dC, dG, dT, ...)



BIOPOLYMERS

- Most complex activities in biology involve polymers
- Biology achieves catalysis, reproduction, specificity, and many other things by making polymers, not by building complex monomeric molecules
- Biological polymers are built up by formal (and actual) elimination of water ($M_1 + M_2 \rightarrow M_{1-2} + H_2O$)
 - Protein: polymer of amino acids
 - RNA: polymer of ribonucleotides
 - DNA: polymer of deoxyribonucleotides
 - Polysaccharides: polymers of sugars
- Except for polysaccharides, they're unbranched

MAJOR CLASSES OF BIOPOLYMERS





Category	Types of monomers	<mol wt/ monomer >	# mono- mers per polymer	Branched?
Protein	20	110	65-5000	no
RNA	4-10	150-300	10-15K	no
DNA	4	130-270	50-10 ⁶	no
Polysac- charide	~10	162	2-10 ⁵	Some- times

PROTEINS

- ◆ Proteins are polymers of L-amino acids
- ◆ There are 20 amino acids in proteins from 99% of all organisms; a few use 2 others
- ◆ Variety of functions:
 - Enzymes (catalysts)
 - Structural proteins
 - Storage and transport proteins
 - Hormones
 - Receptors
 - Nucleic-acid binding proteins
 - Molecular machines
 - Specialized functions (e.g. sweetness)

ALPHA-AMINO ACIDS

- These are the building blocks from which proteins are made
- Alpha carbon covalently bonded to a carboxylate and an amine
- Polymerization makes peptide bonds between the α -amino acids



PROTEIN STRUCTURE

- Most (>80%) proteins fold into a specific 3-D conformation at room temperature
- This isn't intuitively obvious: many polymers are much floppier than that
- A few proteins are “intrinsically unstructured”
- Held together by ionic interactions (H-bonds, salt bridges) and nonpolar interactions (van der Waals and solvent exclusion)



Human DHFR
PDB 1KMV, 1.05Å


LEVELS OF PROTEIN STRUCTURE

- Primary structure: linear sequence of amino acids, e.g. val-gly-ser-leu-asn-cys-ile-val-ala-val-ser-gln-asn-met-gly ...
- Secondary structure: short-range ordering provided by main-chain hydrogen bonds between amine nitrogens and carbonyl oxygens
- Tertiary structure: three-dimensional arrangements of all atoms in a single polypeptide
- Quaternary structure: arrangement of individual polypeptides into a complete multimeric structure

HOW DO PROTEINS PERFORM SO MANY FUNCTIONS?

- ◆ Amino acid side chains have a wide variety of reactivities
- ◆ Scaffolding surrounding active or binding site can significantly change reactivity of specific moieties within the protein
- ◆ Side chains that would normally be hard to ionize become easy to ionize because the ion is stabilized by interactions with neighboring groups
- ◆ Regions that would be hydrophilic if exposed become hydrophobic when buried within protein

IS THAT ALL?



- ◆ No; Some proteins have non-amino-acid components bound, loosely or tightly, to the amino acid chain(s); these entities are *cofactors*
- ◆ Cofactors can be inorganic (e.g. Mg^{2+} ions) or organic (coenzymes)
- ◆ These provide functionalities that would otherwise be unavailable to the polypeptide
- ◆ Many coenzymes are derived from vitamins, particularly the B-complex vitamins


WHY ARE PROTEINS SO BIG?




- ◆ Typical enzyme active site or carrier-protein binding site involves < 5% of the amino acids in the protein
- ◆ So what are all those other amino acids doing?
 - Creating environment perfectly suited to the electrostatics and hydrophobicity or hydrophilicity required for proper function
 - Guarantees that the active site or binding site *keeps* its shape and electrostatic properties over time
 - Enables binding of cofactors and other effectors
 - Allows for appropriate interactions with other macromolecules

DOES EVERY PROTEIN HAVE EXACTLY ONE FUNCTION?

- ◆ No.
- ◆ Some proteins that have evolved for one purpose become pressed into service in a different guise
- ◆ Example: some of the proteins that make up the eye lens are actually enzymes, reused as structural proteins



ENZYMES



James
Sumner

- Enzymes are defined as biological catalysts
- Biochemists figured out in the 1920's that all enzymes were proteins, although it took until the 1930's for that notion to be definitively accepted
- In the 1970's this assertion had to be modified: some enzymes (biocatalysts) are actually RNA molecules
- 1990's: Recognition that the catalytic step in protein synthesis is performed by a specific adenine base in ribosomal RNA

WHAT PROPERTIES MUST AN ENZYME HAVE?

- ◆Catalysis:

 - must lower the activation energy of a reaction

- ◆Specificity:

 - must act preferentially on certain substrates

- ◆Regulation:

 - must be subject to some sort of regulation

HOW IS CATALYSIS ACCOMPLISHED?

◆ Multiple mechanisms, often acting in concert:

- Acid-base catalysis (specific amino acid side-chains participate temporarily in reactions by acting as nucleophiles or electrophiles)
- Proximity effect (enzyme contains channels with appropriate shape and electrostatic properties to cause substrates to travel down them toward one another)
- Induced fit (enzyme reshapes itself to bind substrate, but substrate gets reshaped as well)
- Transition state stabilization


HOW EFFECTIVE ARE ENZYMES, AND HOW ARE THEY REGULATED?

- The best can speed up a reaction by a factor of 10^{12} or more
- They don't change equilibrium: they just make the reaction go faster.
- Regulation by:
 - Control of transcription
 - Control of translation
 - Interference with activity via inhibitors

INHERENT PROPERTIES OF ENZYMES

- ◆ K_m is an inherent property of an enzyme
 - Can be measured for a specific substrate, independent of enzyme and substrate concentration
 - Measures binding affinity of enzyme for substrate
- ◆ V_{max} clearly depends on how much enzyme we put in: the more we put in, the faster the reaction
- ◆ But $k_{cat} = V_{max}/[E]_{tot}$ is an inherent property
 - Measures turnover rate
 - k_{cat} = number of substrate molecules converted to product per unit time by a single enzyme molecule






ENZYME INHIBITION



- ◆ Various small molecules can bind to an enzyme and slow it down
- ◆ Competitive inhibitors compete with the substrate for the active site and raise K_m
- ◆ Noncompetitive inhibitors interfere with catalysis and decrease V_{max}
- ◆ Uncompetitive inhibitors decrease both V_{max} and K_m , but do so in a way that slows the reaction

NUCLEIC ACID BUILDING-BLOCKS

- All nucleotides are based on *nucleobases*, which are either purines (9 ring-atoms, double-ring structures) or pyrimidines (6 ring-atoms)
- The nucleobase is attached to a ribose or deoxyribose sugar by an N-C bond
- A phosphate is attached to the 5'-position of the sugar




6/28/21

Biological Macromolecules

p. 28 of 38

NUCLEOSIDES AND NUCLEOTIDES

- Attach a nitrogen on a nucleobase to carbon 1 of ribose to make a nucleoside
- Attach a phosphate to the 5' position on the nucleoside to make a nucleotide




POLYNUCLEOTIDES

- Connect the 5' phosphate of one nucleotide to the 3' position on the previous nucleotide and continue
- A DNA molecule can comprise millions of nucleotides
- RNA is somewhat smaller but still substantial

NUCLEIC ACID STRUCTURE

- DNA is generally double-helical
- Many RNA molecules are partially double-stranded too
- DNA is unstructured above the double-helix level during G1 but becomes structured during S within spool-like structures called nucleosomes
- Higher levels of organizations occur too





TYPES OF RNA

- We generally recognize four major classes of RNA, differing in size, cellular role, and replacement rate

<i>Type</i>	<i>% RNA</i>	<i>%turn-over</i>	<i>Size, bases DS?</i>	<i>Partly Role</i>
mRNA	3	25	50-10 ⁴ no	protein template
tRNA	15	21	55-90 yes	aa activation
rRNA	80	50	10 ² -10 ⁴	yes transl. catalysis & scaffolding
sRNA	2	4	15-10 ³ yes	various

CARBOHYDRATES

- ◆ Monomers are three-carbon to seven-carbon polyalcohols with one carbonyl group at either C1 or C2. They're very soluble.
- ◆ Simplest: glyceraldehyde & dihydroxyacetone - C₃
- ◆ Most important other ones:
 - Ribose, xylose (C5)
 - Glucose, fructose, galactose (C6)
- ◆ Monomers are fuel, signals, intermediaries (especially when phosphorylated)




SUGAR DERIVATIVES

- ◆ Sugar monomers that have been modified
 - Deoxyribose
 - Phosphorylated sugars (especially at C1 and C5 or C6)
 - Aminated sugars (mostly C6 sugars)
 - N-acetyl sugars
 - Sugar acids
 - Sugar alcohols (carbonyl converted to alcohol)
- ◆ These derivatives are often the active agents in intermediary metabolism

DISACCHARIDES

- Compounds made up of two sugar monomers
- Glucose + glucose \rightarrow Maltose + H_2O
- Glucose + fructose \rightarrow Sucrose + H_2O
- Glucose + galactose \rightarrow Lactose + H_2O
- These are instances of *oligomers* (few rather than many building blocks)




sucrose

STORAGE POLYSACCHARIDES

- ◆ Starch: storage form of glucose in plants
 - Amylose: 1-4 linkages only
 - Amylopectin: mostly 1-4 linkages, a few 1-6
- ◆ Glycogen: storage form of glucose in animals, some bacteria
 - mostly 1-4 linkages, some 1-6
 - Glycogen is primary short-term fuel in humans; long-term fuel is usually fats (more efficient)

GLYCOGEN STRUCTURE

- ◆ Single glucose units broken off & phosphorylated to use as fuel
- ◆ Amylases cleave the 1-4 linkages;
- ◆ Other enzymes deal with the 1-6 linkages



SUGAR-PEPTIDE COMPLEXES

- ◆ Sugars are often complexed either to peptides (oligomers of amino acids) or proteins (polymers of amino acids)
- ◆ Smaller-peptide complexes make up cell walls
- ◆ Full-size proteins that have been decorated with a few sugar molecules are ubiquitous in eukaryotes and often alter the properties of the protein or enable it to be involved in cell-cell communication

